

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
21 November 2002 (21.11.2002)

PCT

(10) International Publication Number
WO 02/092841 A2

(25) International Patent Classification⁷: **C12Q**
(21) International Application Number: PCT/US02/15332
(22) International Filing Date: 14 May 2002 (14.05.2002)
(25) Filing Language: English
(26) Publication Language: English
(30) Priority Data:
60/291,001 14 May 2001 (14.05.2001) US

LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:
US 60/291,001 (CIP)
Filed on 14 May 2001 (14.05.2001)

(71) Applicant (for all designated States except US): **CORVAS INTERNATIONAL, INC.** [US/US]; 3030 Science Park Road, San Diego, CA 92121 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MADISON, Edwin, L.** [US/US]; 11005 Cedarcrest Way, San Diego, CA 92121 (US). **YEH, Jiunn-Chern** [—/US]; 11629 Welsview Parkway, San Diego, CA 92126 (US).

(74) Agents: **SEIDMAN, Stephanie, L.** et al.; Heller Ehrman White & McAuliffe LLP, 4350 La Jolla Village Drive, San Diego, CA 92122-1246 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

— as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/092841 A2

(54) Title: NUCLEIC ACID MOLECULES ENCODING A TRANSMEMBRANE SERINE PROTEASE 10, THE ENCODED POLYPEPTIDES AND METHODS BASED THEREON

(57) Abstract: Provided herein are type II transmembrane serine protease 10 (MTSP10) polypeptides. Zymogen and activated forms of these polypeptides as well as single and two chain forms of the protease domain are also provided. Methods using the polypeptides to identify compounds that modulate the activity of an MTSP10 are provided.

-1-

**NUCLEIC ACID MOLECULES ENCODING A TRANSMEMBRANE SERINE
PROTEASE 10, THE ENCODED POLYPEPTIDES AND METHODS BASED
THEREON**

RELATED APPLICATIONS

- 5 Benefit of priority under 35 U.S.C. § 119(e) is claimed to U.S. provisional application Serial No. 60/291,001, filed May 14, 2001, Edwin L. Madison and Yeh, Jiunn-Chern, entitled "NUCLEIC ACID MOLECULES ENCODING TRANSMEMBRANE SERINE PROTEASE 10, THE ENCODED PROTEINS AND METHODS BASED THEREON." The subject matter of this application is
10 incorporated in its entirety by reference thereto.

FIELD OF INVENTION

- Nucleic acid molecules that encode proteases and portions thereof, particularly protease domains are provided. Also provided are prognostic; diagnostic and therapeutic methods using the proteases and domains thereof and
15 the encoding nucleic acid molecules.

BACKGROUND OF THE INVENTION AND OBJECTS THEREOF

- Cancer, which is a leading cause of death in the United States, is characterized by an increase in the number of abnormal neoplastic cells, which proliferate to form a tumor mass, the invasion of adjacent tissues by these
20 neoplastic tumor cells, and the generation of malignant cells that metastasize via the blood or lymphatic system to regional lymph nodes and to distant sites. Among the hallmarks of cancer is a breakdown in the communication among tumor cells and their environment. Normal cells do not divide in the absence of stimulatory signals and cease dividing in the presence of inhibitory signals.
25 Growth-stimulatory and growth-inhibitory signals, are routinely exchanged between cells within a tissue. In a cancerous, or neoplastic, state, a cell acquires the ability to "override" these signals and to proliferate under conditions in which normal cells do not grow.

- In order to proliferate tumor cells acquire a number of distinct aberrant
30 traits reflecting genetic alterations. The genomes of certain well-studied tumors carry several different independently altered genes, including activated oncogenes and inactivated tumor suppressor genes. Each of these genetic

-2-

changes appears to be responsible for imparting some of the traits that, in the aggregate, represent the full neoplastic phenotype.

A variety of biochemical factors have been associated with different phases of metastasis. Cell surface receptors for collagen, glycoproteins such as laminin, and proteoglycans, facilitate tumor cell attachment, an important step in invasion and metastases. Attachment triggers the release of degradative enzymes which facilitate the penetration of tumor cells through tissue barriers. Once the tumor cells have entered the target tissue, specific growth factors are required for further proliferation. Tumor invasion and progression involve a complex series of events, in which tumor cells detach from the primary tumor, break down the normal tissue surrounding it, and migrate into a blood or lymphatic vessel to be carried to a distant site. The breaking down of normal tissue barriers is accomplished by the elaboration of specific enzymes that degrade the proteins of the extracellular matrix that make up basement membranes and stromal components of tissues.

A class of extracellular matrix degrading enzymes has been implicated in tumor invasion. Among these are the matrix metalloproteinases (MMP). For example, the production of the matrix metalloproteinase stromelysin is associated with malignant tumors with metastatic potential (see, *e.g.*, McDonnell *et al.* (1990) *Smnrs. in Cancer Biology* 1:107-115; McDonnell *et al.* (1990) *Cancer and Metastasis Reviews* 9:309-319).

The capacity of cancer cells to metastasize and invade tissue is facilitated by degradation of the basement membrane. Several proteinase enzymes, including the MMPs, have been reported to facilitate the process of invasion of tumor cells. MMPs are reported to enhance degradation of the basement membrane, which thereby permits tumorous cells to invade tissues. For example, two major metalloproteinases having molecular weights of about 70 kDa and 92 kDa appear to enhance ability of tumor cells to metastasize.

Type II Transmembrane Serine Proteases

In addition to the MMPs, serine proteases have been implicated in neoplastic disease progression. Most serine proteases, which are either secreted enzymes or are sequestered in cytoplasmic storage organelles, have roles in blood coagulation, wound healing, digestion, immune responses and tumor invasion and metastasis. A class of cell surface proteins designated type II transmembrane serine proteases, which are membrane-anchored proteins with additional extracellular domains, has been identified. As cell surface proteins, they are positioned to play a role in intracellular signal transduction and in mediating cell surface proteolytic events.

Cell surface proteolysis is a mechanism for the generation of biologically active proteins that mediate a variety of cellular functions. Membrane-associated proteases include membrane-type metalloproteinases (MT-MMP), ADAMs (proteases that contain disintegrin-like and metalloproteinase domains) and the transmembrane serine proteases. In mammals, at least 17 members of the transmembrane serine protease family are known, including seven in humans (see, Hooper *et al.* (2001) *J. Biol. Chem.* 276:857-860). These include: corin (accession nos. AF133845 and AB013874; see, Yan *et al.* (1999) *J. Biol. Chem.* 274:14926-14938; Tomia *et al.* (1998) *J. Biochem.* 124:784-789; Uan *et al.* (2000) *Proc. Natl. Acad. Sci. U.S.A.* 97:8525-8529); enteropeptidase (also designated enterokinase; accession no. U09860 for the human protein; see, Kitamoto *et al.* (1995) *Biochem.* 27: 4562-4568; Yahagi *et al.* (1996) *Biochem. Biophys. Res. Commun.* 219:806-812; Kitamoto *et al.* (1994) *Proc. Natl. Acad. Sci. U.S.A.* 91:7588-7592; Matsushima *et al.* (1994) *J. Biol. Chem.* 269:19976-19982); human airway trypsin-like protease (HAT; accession no. AB002134; see Yamaoka *et al.* *J. Biol. Chem.* 273:11894-11901); MTSP1 and matriptase (also called TADG-15; see SEQ ID Nos. 1 and 2; accession nos. AF133086/AF118224, AF04280022; Takeuchi *et al.* (1999) *Proc. Natl. Acad. Sci. U.S.A.* 96:11054-1161; Lin *et al.* (1999) *J. Biol. Chem.* 274:18231-18236; Takeuchi *et al.* (2000) *J. Biol. Chem.* 275:26333-26342; and Kim *et al.* (1999) *Immunogenetics* 49:420-429); hepsin (see, accession nos. M18930, AF030065, X70900; Leytus *et al.* (1988) *Biochem.* 27: 11895-11901; Vu *et al.* (1997) *J.*

-4-

Biol. Chem. 272:31315-31320; and Farley *et al.* (1993) *Biochem. Biophys. Acta* 1173:350-352; and see, U.S. Patent No. 5,972,616); TMPRS2 (see, Accession Nos. U75329 and AF113596; Paoloni-Giacobino *et al.* (1997) *Genomics* 44:309-320; and Jacquinet *et al.* (2000) *FEBS Lett.* 468: 93-100); and TMPRSS4 (see, 5 Accession No. NM 016425; Wallrapp *et al.* (2000) *Cancer* 60:2602-2606).

Serine proteases, including transmembrane serine proteases and secreted proteases, have been implicated in processes involved in neoplastic development and progression. While the precise, detailed mechanism by which these proteases promote tumor growth and progression has not been elaborated, 10 serine proteases and inhibitors thereof are involved in the control of many intra- and extracellular physiological processes, including degradative actions in cancer cell invasion, metastatic spread, and neovascularization of tumors, that are involved in tumor progression. It is believed that proteases are involved in the degradation of extracellular matrix (ECM) and contribute to tissue remodeling, 15 and are necessary for cancer invasion and metastasis. The activity and/or expression of some proteases have been shown to correlate with tumor progression and development.

For example, a membrane-type serine protease MTSP1 (also called matriptase; see SEQ ID Nos. 1 and 2 from U.S. Patent No. 5,972,616; and 20 GenBank Accession No. AF118224; (1999) *J. Biol. Chem.* 274:18231-18236; U.S. Patent No. 5,792,616; see, also Takeuchi (1999) *Proc. Natl. Acad. Sci. U.S.A.* 96:11054-1161) that is expressed in epithelial cancer and normal tissue (Takeuchi *et al.* (1999) *Proc. Natl. Acad. Sci. USA* 96:11054-61) has been identified. Matriptase was originally identified in human breast cancer cells as a 25 major gelatinase (see, U.S. Patent No. 5,482,848) and was initially believed to be a type of matrix metalloprotease (MMP). It has been proposed that it plays a role in the metastasis of breast cancer. Matriptase also is expressed in a variety of epithelial tissues with high levels of activity and/or expression in the human gastrointestinal tract and the prostate. MTSPs, designated MTSP3, MTSP4, 30 MTSP6 have been described in published International PCT application No. WO 01/57194, based in International PCT application No. PCT/US01/03471.

RECTIFIED SHEET (RULE 91)

-5-

Prostate-specific antigen (PSA), a kallikrein-like serine protease, degrades extracellular matrix glycoproteins fibronectin and laminin, and, has been postulated to facilitate invasion by prostate cancer cells (Webber *et al.* (1995) *Clin. Cancer Res.*, 1(10):1089-94). Blocking PSA proteolytic activity with

5 PSA-specific monoclonal antibodies results in a dose-dependent decrease *in vitro* in the invasion of the reconstituted basement membrane Matrigel by LNCaP human prostate carcinoma cells which secrete high levels of PSA.

Hepsin, a cell surface serine protease identified in hepatoma cells, is overexpressed in ovarian cancer (Tanimoto *et al.* (1997) *Cancer Res.*, 57(14):2884-7). The hepsin transcript appears to be abundant in carcinoma

10 tissue and is almost never expressed in normal adult tissue, including normal ovary. It has been suggested that hepsin is frequently overexpressed in ovarian tumors and therefore can be a candidate protease in the invasive process and growth capacity of ovarian tumor cells.

15 A serine protease-like gene, designated normal epithelial cell-specific 1 (NES1) (Liu *et al.*, *Cancer Res.*, 56(14):3371-9 (1996)) has been identified. Although expression of the NES1 mRNA is observed in all normal and immortalized nontumorigenic epithelial cell lines, the majority of human breast cancer cell lines show a drastic reduction or a complete lack of its expression.

20 The structural similarity of NES1 to polypeptides known to regulate growth factor activity and a negative correlation of NES1 expression with breast oncogenesis suggest a direct or indirect role for this protease-like gene product in the suppression of tumorigenesis.

Hence transmembrane serine proteases appear to be involved in the

25 etiology and pathogenesis of tumors. There is a need to further elucidate their role in these processes and to identify additional transmembrane proteases. Therefore, it is an object herein to provide transmembrane serine protease (MTSP) proteins and nucleic acids encoding such MTSP proteases that are involved in the regulation of or participate in tumorigenesis and/or

30 carcinogenesis. It is also an object herein to provide prognostic, diagnostic and therapeutic screening methods using such proteases and the nucleic acids encoding such proteases.

RECTIFIED SHEET (RULE 91)

SUMMARY

Provided herein is the protease domain of a protein designated herein as MTSP10. The protease domain and full-length protein, including the zymogen and activated forms, and uses thereof are also provided. Proteins encoded by
5 splice variants are also provided. Hence, provided herein is a family proteins designated MTSP10, and functional domains, including one or more of a transmembrane (TM) domain, two CUB domains, three LDL receptor type a domains and a serine protease catalytic domain, especially protease (or catalytic) domains thereof. Also provided are muteins and other derivatives and analogs
10 thereof. Also provided herein are nucleic acids encoding the MTSP10s.

The protease domains provided herein include, but are not limited to, the single chain region having an N-terminus at the cleavage site for activation of the zymogen, through the C-terminus, or C-terminal truncated portions thereof that exhibit proteolytic activity as a single-chain polypeptide in *in vitro* proteolysis
15 assays of MTSP10, from a mammal, including a human, that, for example, displays functional activity in tumor cells that is different from its activity non-tumor cells.

Nucleic acid molecules encoding the proteins and protease domains are also provided. Nucleic acid molecules that encode a single-chain protease
20 domain or catalytically active portion thereof and also those that encode the full-length MTSP10 or portions thereof are provided. In one embodiment, a nucleic acid that encodes a MTSP, designated MTSP10 is provided. The nucleic acid molecule includes the sequence of nucleotides set forth in SEQ ID No. 5 or SEQ ID No. 22 or a portion thereof (see, also EXAMPLE 1) that encodes a
25 catalytically active polypeptide or a domain thereof.

Also provided are nucleic acid molecules that encode all or a portion of a catalytically active polypeptide, or a nucleic acid molecule that encodes the protease domain or a larger polypeptide that can include up to the full length polypeptide and that hybridizes to such MTSP10-encoding nucleic acid along
30 their full-length or along at least about 70%, 80% or 90% of the full-length and encode the protease domain or portion thereof are provided.

-7-

Hybridization is generally effected under conditions of at least low, generally at least moderate, and often high stringency.

The isolated nucleic acid fragment is DNA, including genomic or cDNA, or is RNA, or can include other components, such as protein nucleic acid or other
5 nucleotide analogs. The isolated nucleic acid may include additional components, such as heterologous or native promoters, and other transcriptional and translational regulatory sequences, these genes may be linked to other genes, such as reporter genes or other indicator genes or genes that encode indicators.

10 Also provided is an isolated nucleic acid molecule that includes the sequence of molecules that is complementary to the nucleotide sequence encoding MTSP10 or the portion thereof.

Also provided are nucleic acid molecules that hybridize under conditions of at least low stringency, generally moderate stringency, more typically high
15 stringency to the sequence of nucleotides set forth in SEQ ID No. 5 or SEQ ID No. 22 or degenerates thereof. In one embodiment, the isolated nucleic acid fragment hybridizes to a nucleic acid molecule containing the nucleotide sequence set forth in SEQ ID No. 5 or SEQ ID No. 22 (or degenerates thereof) under high stringency conditions. In one embodiment, it contains the sequence
20 of nucleotides set forth in SEQ ID No. 5. A full-length MTSP10 polypeptide includes the sequence of amino acids set forth in SEQ ID No. 6 or SEQ ID No. 23, and is encoded by a sequence of nucleotides set forth in SEQ ID No. 5 or SEQ ID No. 22 or degenerates thereof. Methods for isolating nucleic acid encoding other MTSP10s, including nucleic acid molecules encoding full-length
25 molecules and splice variants and MTSPs from species, such as cows, sheep, goats, pigs, horses, primates, including chimpanzees and gorillas, rodents, dogs, cats and other species of interest, such as domesticated animals, farm and zoo animals are also provided. The nucleic acid molecules provided herein, including those set forth in SEQ ID Nos. 5 and 23 can be used to obtain nucleic acid
30 molecules encoding full-length MTSP10 polypeptides from human sources or from other species, such as by screening appropriate libraries using the nucleic acid molecules or selected primers or probes based thereon.

-8-

Also provided are fragments thereof or oligonucleotides that can be used as probes or primers and that contain at least about 10, 14, 16 nucleotides, generally less than 1000 or less than or equal to 100, set forth in SEQ ID No. 5 or SEQ ID No. 22 (or the complement thereof); or contain at least about 30
5 nucleotides (or the complement thereof) or contain oligonucleotides that hybridize along their full-length (or at least about 70, 80 or 90% thereof) to any such fragments or oligonucleotides. The length of the fragments are a function of the purpose for which they are used and/or the complexity of the genome of interest. Generally probes and primers contain less than about 30, 50, 150 or
10 500 nucleotides.

Also provided are plasmids containing any of the nucleic acid molecules provided herein. Cells containing the plasmids are also provided. Such cells include, but are not limited to, bacterial cells, yeast cells, fungal cells, plant cells, insect cells and animal cells.

15 Methods of expressing the encoded MTSP10 polypeptide and portions thereof using the cells are also provided, as are cells that express MTSP10 on the cell surface. Such cells are used in methods of identifying candidate therapeutic compounds.

MTSP10, particularly the protease domain thereof, can be produced by
20 growing the above-described cells under conditions whereby the MTSP10 is expressed by the cells, and recovering the expressed MTSP10 polypeptide.

Also provided are cells, generally eukaryotic cells, such as mammalian cells and yeast cells, in which the MTSP10 polypeptide is expressed on the surface of the cells. Such cells are used in drug screening assays to identify
25 compounds that modulate the activity of the MTSP10 polypeptide. These assays, including *in vitro* binding assays, and transcription based assays in which signal transduction is mediated directly or indirectly, such as via activation of pro-growth factors, by the MTSP10 is assessed.

Also provided are peptides that are encoded by such nucleic acid
30 molecules. Included among those polypeptides are the MTSP10 protease domain or a polypeptide with amino acid changes such that the specificity and/or protease activity remains substantially unchanged. In particular, a substantially

RECTIFIED SHEET (RULE 91)

-9-

purified mammalian MTSP10 polypeptide is provided that includes a serine protease catalytic domain and may additionally include other domains. The MTSP10 can form homodimers and can also form heterodimers with some other protein, such as a membrane-bound protein. Also provided is a substantially
5 purified protein including a sequence of amino acids that has at least 60%, 70%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% identity to the MTSP10 where the percentage identity is determined using standard algorithms and gap penalties that maximize the percentage identity. A human MTSP10 polypeptide is
10 exemplified, although other mammalian MTSP10 polypeptides are contemplated. Splice variants of the MTSP10, particularly those with a proteolytically active protease domain, are contemplated herein.

In other embodiments, substantially purified polypeptides that include a protease domain of a MTSP10 polypeptide or a catalytically active portion
15 thereof are provided. Among these are polypeptides that include a sequence of amino acids that has at least 60%, 70%, 80%, 85%, 90%, 95% or 100% sequence identity to SEQ ID No. 6 or SEQ ID No. 23 or to a portion thereof that includes a catalytically active polypeptide or a CUB domain or an LDL receptor domain.

20 Also provided are muteins of the single chain protease domain of MTSP10 particularly muteins in which the Cys residue in the protease domain that is free (*i.e.*, does not form disulfide linkages with any other Cys residue in the protease domain) is substituted with another amino acid substitution, typically, although not necessarily, with a conservative amino acid substitution
25 or a substitution that does not eliminate the activity, and muteins in which a glycosylation site(s) is eliminated.

Hence muteins in which one or more of the Cys residues, particularly, a residue that is paired in the activated two form, but unpaired in the protease domain alone (*i.e.*, the Cys a residue position 26 (see SEQ ID Nos. 5 and 6) in the
30 protease domain), is/are replaced with any amino acid, typically, although not necessarily, a conservative amino acid residue, such as Ser, are contemplated. Muteins of MTSP10, particularly those in which Cys residues, such as the

RECTIFIED SHEET (RULE 91)

-10-

unpaired Cys in the single chain protease domain, is replaced with another amino acid that does not eliminate the activity, are provided. Muteins in which other conservative or non-conservative amino acid substitutions in which catalytic activity is retained are also contemplated (see, *e.g.*, Table 1, for exemplary amino acid substitutions).

MTSP10 polypeptides, including, but not limited to splice variants thereof, and nucleic acids encoding MTSPs, and domains, derivatives and analogs thereof are provided herein. Single chain protease domains that have an N-terminus functionally equivalent to that generated by activation of the zymogen form of MTSP10 are also provided. The cleavage site for the protease domain of MTSP10 is between amino acid R and amino acid I (R↓IIGGT) (see SEQ ID NO. 6, residues 1-5; residues 462-467 SEQ ID No. 23). There are potential glycosylation sites at N₆₇, N₂₇₂, N₃₃₆, N₃₈₃, N₄₀₉ and N₄₁₈ SEQ ID No. 23). The catalytic triad of the MTSP10 in SEQ ID No. 23 is H₅₀₃, D₅₅₁ and S₆₄₇.

There are potential disulfide bonds as follows: C₄₈₈-C₅₀₄, C₅₈₇-C₆₅₃; C₆₁₉-C₆₃₂; C₆₄₃-C₆₇₃ (see SEQ ID Nos. 22 and 23) (chymotrypsin numbering 42 to 58; 136-201; 168-182 and 191-220). Disulfide bonds form between the Cys residues C₅₇₃-C₂₉₆ to link the protease domain to another domain so that upon activation cleavage (between residues R₄₆₂ and I₄₆₃ of SEQ ID No. 23) the resulting polypeptide is a two chain molecule. The C₅₇₃ (SEQ ID NO. 23, is a free Cys in the single chain form of the protease domain. As noted the protease also can be provided as a two chain molecule. Single chain and two chain forms are proteolytically active. A two chain form of the polypeptide set forth in SEQ ID No. 23 is provided; smaller catalytically active two chain forms are also provided. A two chain form is produced by bonding, typically between the C₅₇₃ and a Cys outside the protease domain, such as Cys₂₉₆. Upon activation cleavage the bond remains resulting in a two chain polypeptide. The size of chain "A" is a function the starting length of the polypeptide prior to activation cleavage between the R₄₆₂ and I₄₆₃. Any length polypeptide that includes the protease domain (residues 463-692 of SEQ ID No. 23) or catalytically active fragments thereof, is contemplated herein. Two chain forms include at least the protease domain of a polypeptide from C₂₉₆ up to and including C₅₇₃.

RECTIFIED SHEET (RULE 91)

-11-

MTSPs are expressed or are activated in certain tumor or cancer cells such as lung, prostate, colon and breast, ovarian, pancreatic, lung in other tumors. MTSP10 is of interest because it is expressed or is active in tumor cells. In particular, it is shown herein, that MTSP10 is, for example, expressed
5 in esophageal tumor tissues, in lung carcinoma, prostate cancers, pancreatic and breast cancers and in cell lines as well as in certain normal cells and tissues (see *e.g.*, EXAMPLES for tissue-specific expression profile). The level of activated MTSP10 can be diagnostic of prostate, uterine, lung esophagus, or colon cancer or leukemia or other cancer. The expression and/or activation of MTSP10 on or
10 in the vicinity of a cell or in a bodily fluid in a subject can be a marker for breast, prostate, lung, colon and other cancers.

Hence the MTSPs provided herein can serve as diagnostic markers for certain tumors. In certain embodiments, the MTSP10 polypeptide is detectable in a body fluid at a level that differs from its level in body fluids in a subject not
15 having a tumor. In other embodiments, the polypeptide is present in a tumor; and a substrate or cofactor for the polypeptide is expressed at levels that differ from its level of expression in a non-tumor cell in the same type of tissue. In other embodiments, the level of expression and/or activity of the MTSP10 polypeptide in tumor cells differs from its level of expression and/or activity in
20 non-tumor cells. In other embodiments, the MTSP10 is present in a tumor; and a substrate or cofactor for the MTSP10 is expressed at levels that differ from its level of expression in a non-tumor cell in the same type of tissue.

Assays for identifying effectors, such as compounds, including small molecules, and conditions, such pH, temperature and ionic strength, that
25 modulate the activation, expression or activity of MTSP10 are also provided herein. In exemplary assays, the effects of test compounds on the ability of a protease domain of MTSP10 to proteolytically cleave a known substrate, typically a fluorescently, chromogenically or otherwise detectably labeled substrate, are assessed. Agents, generally compounds, particularly small
30 molecules, that modulate the activity of the protease domain are candidate compounds for modulating the activity of the MTSP10. The protease domains can also be used to produce protease-specific antibodies.

-12-

Also provided are methods for screening for compounds that modulate the activity of MTSP10. The compounds are identified by contacting them with the MTSP10 or protease domain thereof and a substrate for the MTSP10. A change in the amount of substrate cleaved in the presence of the compounds
5 compared to that in the absence of the compound indicates that the compound modulates the activity of the MTSP10. Such compounds are selected for further analyses or for use to modulate the activity of the MTSP10, such as inhibitors or agonists. The compounds can also be identified by contacting the substrates with a cell that expresses the MTSP10 or the extracellular domain or
10 proteolytically active portion thereof.

Also provided herein are methods of modulating the activity of the MTSP10 and screening for compounds that modulate, including inhibit, antagonize, agonize or otherwise alter the activity of the MTSP10. Of particular interest is the extracellular domain of MTSP10 that includes the proteolytic
15 (catalytic) portion of the protein.

Additionally provided herein are antibodies that specifically bind to single and two chains forms of MTSP10, cells, combinations, kits and articles of manufacture that contain the antibodies. Antibodies that specifically bind to the MTSP10, particularly the single-chain protease domain, the two-chain form of
20 the protease domain, the zymogen and activated form of MTSP10 and other fragments thereof. Neutralizing antibodies that inhibit a biological activity, particularly protease activity are also provided.

Further provided herein are prognostic, diagnostic, therapeutic screening methods using MTSP10 and the nucleic acids encoding MTSP10. In particular,
25 the prognostic, diagnostic and therapeutic screening methods are used for preventing, treating, or for finding agents useful in preventing or treating, tumors or cancers such as lung carcinoma, colon adenocarcinoma and ovarian carcinoma.

Also provided herein are modulators of the activity of MTSP10, especially
30 the modulators obtained according to the screening methods provide herein. Such modulators can have use in treating cancerous conditions.

-13-

Methods of diagnosing a disease or disorder characterized by detecting an aberrant level of an MTSP10 in a subject is provided. The method can be practiced by measuring the level of the DNA, RNA, protein or functional activity of the MTSP10. An increase or decrease in the level of the DNA, RNA, protein or functional activity of the MTSP, relative to the level of the DNA, RNA, protein or functional activity found in an analogous sample not having the disease or disorder (or other suitable control) is indicative of the presence of the disease or disorder in the subject or other relative any other suitable control.

Also provided are methods of identifying a compound that binds to the single-chain and/or two-chain form of MTSP10, by contacting a test compound with a both forms; determining to which form the compound binds; and if it binds to a form of MTSP10, further determining whether the compound has at least one of the following properties:

- (i) inhibits activation of the single-chain zymogen form of MTSP10;
- (ii) inhibits activity of the two-chain or single-chain form; and
- (iii) inhibits dimerization of the protein.

The forms can be full length or truncated forms, including but not limited to, the protease domain resulting from cleavage at the activation cleavage site (between amino acids R₄₆₂ and I₄₆₃ SEQ ID No. 23); or from expression of the protease domain or catalytically active portions thereof.

Pharmaceutical composition containing the protease domain and/or full-length or other domain of an MTSP10 polypeptide are provided herein in a pharmaceutically acceptable carrier or excipient are provided herein.

Also provided are articles of manufacture that contain MTSP10 polypeptide and protease domains of MTSP10 in single chain forms or activated forms. The articles contain a) packaging material; b) the polypeptide (or encoding nucleic acid), particularly the single chain protease domain thereof; and c) a label indicating that the article is for use in assays for identifying modulators of the activities of an MTSP10 polypeptide and are provided herein.

Conjugates containing a) an MTSP10 polypeptide or protease domain in a single or two chain form; and b) a targeting agent linked to the MTSP directly or via a linker, wherein the agent facilitates: i) affinity isolation or purification of the

RECTIFIED SHEET (RULE 91)

-14-

conjugate; ii) attachment of the conjugate to a surface; iii) detection of the conjugate; or iv) targeted delivery to a selected tissue or cell, are provided herein. The conjugate can contain a plurality of agents linked thereto. The conjugate can be a chemical conjugate; and it can be a fusion protein. The
5 targeting agent can be a protein or peptide fragment. The protein or peptide fragment can include a protein binding sequence, a nucleic acid binding sequence, a lipid binding sequence, a polysaccharide binding sequence, or a metal binding sequence.

Combinations, kits and articles of manufacture containing the MTSP10
10 polypeptides, domains thereof, or encoding nucleic acids are also provided herein. For example, combinations are provided herein. The combination can include: a) an inhibitor of the activity of an MTSP10; and b) an anti-cancer treatment or agent. The MTSP inhibitor and the anti-cancer agent can be formulated in a single pharmaceutical composition or each is formulated in a
15 separate pharmaceutical composition. The MTSP10 inhibitor can be an antibody or a fragment or binding portion thereof made against the MTSP10, such as an antibody that specifically binds to the protease domain, an inhibitor of MTSP10 production, or an inhibitor of MTSP10 membrane-localization or an inhibitor of MTSP10 activation. Other MTSP10 inhibitors include, but are not limited to, an
20 antisense nucleic acid or double-stranded RNA (dsRNA), such as RNAi, encoding the MTSP10, particularly a portion of the protease domain; a nucleic acid encoding at least a portion of a gene encoding the MTSP10 with a heterologous nucleotide sequence inserted therein such that the heterologous sequence inactivates the biological activity of MTSP10 or the gene encoding it. For
25 example, the portion of the gene encoding the MTSP10 can flank the heterologous sequence to promote homologous recombination with a genomic gene encoding the MTSP10.

Also provided are methods for treating or preventing a tumor or cancer in a mammal by administering to a mammal an effective amount of an inhibitor of
30 an MTSP10, whereby the tumor or cancer is treated or prevented. The MTSP10 inhibitor used in the treatment or for prophylaxis is administered with a pharmaceutically acceptable carrier or excipient. The mammal treated can be a

RECTIFIED SHEET (RULE 91)

-15-

human. The treatment or prevention method can additionally include administering an anti-cancer treatment or agent simultaneously with, subsequent to, or before administration of the MTSP10 inhibitor.

Also provided are transgenic non-human animals bearing inactivated
5 genes encoding the MTSP and bearing the genes encoding the MTSP10 under non-native promotor control are provided. Such animals are useful in animal models of tumor initiation, growth and/or progression models. Transgenic non-human animals containing heterologous nucleic acid MTSP10 under native, non-native promotor control or on an exogenous element, such as a plasmid or
10 artificial chromosome, are additionally provided herein. In particular, recombinant non-human animals are provided herein, where the gene of an MTSP10 is under control of a promoter that is not the native promoter of the gene or that is not the native promoter of the gene in the non-human animal or where the nucleic acid encoding the MTSP10 is heterologous to the non-human
15 animal and the promoter is the native or a non-native promoter or the MTSP10 is on an extrachromosomal element, such as a plasmid or artificial chromosome. Recombinant and transgenic animals can be produced by homologous recombination and non-homologous recombination methods.

Methods of gene therapy are provided. Such methods can be effected
20 administering *in vivo* or *ex vivo* an inactivating form of the MTSP10 or by administering an MTSP-encoding nucleic acid molecule are also provided.

Also provided are methods of treatment of tumors by administering a prodrug that is activated by MTSP10 that is expressed or active in tumor cells, particularly those in which its functional activity in tumor cells is greater than in
25 non-tumor cells. The prodrug is administered and, upon administration, active MTSP10 expressed on cells cleaves the prodrug and releases active drug in the vicinity of the tumor cells. The active anti-cancer drug accumulates in the vicinity of the tumor. This is particularly useful in instances in which MTSP10 is

RECTIFIED SHEET (RULE 91)

-16-

expressed or active in greater quantity, higher level or predominantly in tumor cells compared to other cells.

Also provided are methods of diagnosing the presence of a pre-malignant lesion, a malignancy, or other pathologic condition in a subject, by obtaining a biological sample from the subject; and exposing it to a detectable agent that binds to a two-chain and/or single-chain form of MTSP10, where the pathological condition is characterized by the presence or absence of the two-chain and/or single-chain form.

Methods of inhibiting tumor invasion or metastasis or treating a malignant or pre-malignant condition by administering an agent that inhibits activation of the zymogen form of MTSP10 or an activity of the activated form are provided. The conditions include, but are not limited to, a condition, such as a tumor, of the breast, cervix, prostate, lung, ovary or colon.

DETAILED DESCRIPTION

A. DEFINITIONS

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of skill in the art to which the invention(s) belong. All patents, patent applications, published applications and publications, Genbank sequences, websites and other published materials referred to throughout the entire disclosure herein, unless noted otherwise, are incorporated by reference in their entirety. In the event that there are a plurality of definitions for terms herein, those in this section prevail. Where reference is made to a URL or other such identifier or address, it is understood that such identifiers can change and particular information on the internet can come and go, but equivalent information can be found by searching the internet. Reference thereto evidences the availability and public dissemination of such information.

As used herein, the abbreviations for any protective groups, amino acids and other compounds, are, unless indicated otherwise, in accord with their common usage, recognized abbreviations, or the IUPAC-IUB Commission on Biochemical Nomenclature (see, (1972) *Biochem.* 11:942-944).

RECTIFIED SHEET (RULE 91)

-17-

As used herein, serine protease refers to a diverse family of proteases wherein a serine residue is involved in the hydrolysis of proteins or peptides. The serine residue can be part of the catalytic triad mechanism, which includes a serine, a histidine and an aspartic acid in the catalysis, or be part of the hydroxyl/ ϵ -amine or hydroxyl/ α -amine catalytic dyad mechanism, which involves a serine and a lysine in the catalysis. Of particular interest are SPs of mammalian, including human, origin. Those of skill in this art recognize that, in general, single amino acid substitutions in non-essential regions of a polypeptide do not substantially alter biological activity (see, *e.g.*, Watson *et al.* (1987) *Molecular Biology of the Gene*, 4th Edition, The Benjamin/Cummings Pub. co., p.224).

As used herein, "transmembrane serine protease (MTSP)" refers to a family of transmembrane serine proteases that share common structural features as described herein (see, also Hooper *et al.* (2001) *J. Biol. Chem.* 276:857-860). Thus, reference, for example, to "MTSP" encompasses all proteins encoded by the MTSP gene family, including but are not limited to: MTSP3, MTSP4, MTSP6, MTSP7 or an equivalent molecule obtained from any other source or that has been prepared synthetically or that exhibits the same activity. Other MTSPs include, but are not limited to, corin, enteropeptidase, human airway trypsin-like protease (HAT), MTSP1, TMPRSS2 and TMPRSS4. Sequences of encoding nucleic acid molecules and the encoded amino acid sequences of exemplary MTSPs and/or domains thereof are set forth, for example in U.S. application Serial No. 09/776,191 (SEQ ID Nos. 1-12, 49, 50 and 61-72 therein, published as International PCT application No. WO 01/57194). The term also encompass MTSPs with amino acid substitutions that do not substantially alter activity of each member and also encompasses splice variants thereof. Suitable substitutions, including, although not necessarily, conservative substitutions of amino acids, are known to those of skill in this art and can be made without eliminating the biological activity, such as the catalytic activity, of the resulting molecule.

As used herein an MTSP10, whenever referenced herein, includes at least one or all of or any combination of:

RECTIFIED SHEET (RULE 91)

-18-

a polypeptide encoded by the sequence of nucleotides set forth in SEQ ID No. 5 or SEQ ID No. 22 or by a sequence of nucleotides that includes nucleotides that encode the sequence of amino acids set forth in SEQ ID No. 6 or SEQ ID No. 23;

5 a polypeptide encoded by a sequence of nucleotides that hybridizes under conditions of low, moderate or high stringency to the sequence of nucleotides set forth in is set forth as SEQ ID No. 5 or SEQ ID No. 22;

a polypeptide that includes the sequence of amino acids set forth in SEQ ID No. 6 or SEQ ID No. 23 or a catalytically active portion thereof;

10 a polypeptide that includes a sequence of amino acids having at least about 60%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% sequence identity with the sequence of amino acids set forth in SEQ ID No. 6 or SEQ ID No. 23; and/or

15 a polypeptide encoded by a splice variant of the MTSP10 that includes the sequence of amino acids set forth in SEQ ID No. 6 or SEQ ID No. 23.

In particular, MTSP10 polypeptides, with a protease domain as indicated in SEQ ID Nos. 5 and 6, is provided. The polypeptide is a single or two chain
20 polypeptide. Smaller portions thereof that retain protease activity are also provided. The protease domains from MTSPs vary in size and constitution, including insertions and deletions in surface loops. They retain conserved structure, including at least one of the active site triad, primary specificity pocket, oxyanion hole and/or other features of serine protease domains of
25 proteases. Thus, for purposes herein, the protease domain is a portion of an MTSP, as defined herein, and is homologous to a domain of other MTSPs, such as corin, enteropeptidase, human airway trypsin-like protease (HAT), MTSP1, TMPRSS2, and TMPRSS4, which have been previously identified; it was not recognized, however, that an isolated single chain form of the protease domain
30 could function proteolytically in *in vitro* assays. As with the larger class of enzymes of the chymotrypsin (S1) fold (see, *e.g.*, Internet accessible MEROPS data base), the MTSPs protease domains share a high degree of amino acid

RECTIFIED SHEET (RULE 91)

-19-

sequence identity. The His, Asp and Ser residues necessary for activity are present in conserved motifs. The activation site, which results in the N-terminus of the second chain in the two chain form is located in a conserved motif and readily can be identified. In the exemplified MTSP10, it is between residues R₄₆₂ and I₄₆₃ of SEQ ID No. 23.

The MTSP10 can be from any animal, particularly a mammal, and includes but are not limited to, primates, including humans, rodents, fowl, ruminants and other animals. The full-length zymogen or two-chain activated form is contemplated or any domain thereof, including the protease domain, which can be a two-chain activated form, or a single chain form.

As used herein, a "protease domain of an MTSP" refers to an extracellular protease domain of an MTSP that exhibits proteolytic activity and shares homology and structural features with the chymotrypsin/trypsin family protease domains. Hence it is at least the minimal portion of the domain that exhibits proteolytic activity as assessed by standard *in vitro* assays. Contemplated herein are such protease domains and catalytically active portions thereof. Also provided are truncated forms of the protease domain that include the smallest fragment thereof that acts catalytically as a single chain form.

A protease domain of an MTSP10, whenever referenced herein, includes at least one or all of or any combination of or a catalytically active portion of:

- a) a polypeptide that includes the sequence of amino acids set forth in SEQ ID No. 6, particularly amino acids 1-230 thereof;
- b) a polypeptide that includes the sequence of amino acids set forth in SEQ ID No. 23;
- c) a polypeptide encoded by a sequence of nucleotides that hybridizes under conditions of low, moderate or high stringency to the sequence of nucleotides set forth in SEQ ID No. 5 or to SEQ ID No. 22;
- d) a polypeptide that includes the sequence of amino acids set forth in SEQ ID No. 6 or SEQ ID No. 23 or a catalytically active portion thereof;
- e) a polypeptide that includes a sequence of amino acids having at least about 60%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99%

-20-

sequence identity with the sequence of amino acids set forth in SEQ ID No. 6 or SEQ ID No. 23; and/or

f) a protease domain of a polypeptide encoded by a splice variant of a sequence of nucleotides that encodes an MTSP10 of any of a)-e).

5 The protease domains of MTSPs vary in size and constitution, including insertions and deletions in surface loops. They retain conserved structure, including at least one of the active site triad, primary specificity pocket, oxyanion hole and/or other features of serine protease domains of proteases. Thus, for purposes herein, the protease domain is a portion of an MTSP, as
10 defined herein, and is homologous to a domain of other MTSP. As with the larger class of enzymes of the chymotrypsin (S1) fold (see, *e.g.*, Internet accessible MEROPS data base), the MTSP protease domains share a high degree of amino acid sequence identity. The His, Asp and Ser residues necessary for activity are present in conserved motifs. The activation site, whose cleavage
15 creates the N-terminus of the protease domain in the two-chain forms is located in a conserved motif and readily can be identified.

By active form is meant a form active *in vivo* and/or *in vitro*. As described herein, the protease domain also can exist as a two-chain form. It is shown herein that, at least *in vitro*, the single chain forms of the SPs and the
20 catalytic domains or proteolytically active portions thereof (typically C-terminal truncations) exhibit protease activity. Hence provided herein are isolated single chain forms of the protease domains of SPs and their use in *in vitro* drug screening assays for identification of agents that modulate the activity thereof.

As used herein, the catalytically active domain of an MTSP refers to the
25 protease domain. Reference to the protease domain of an MTSP generally refers to the single chain form of the protein. If the two-chain form or both forms is intended, it is so-specified. The zymogen form of each protein is a single chain, which is converted to the active two chain form by activation cleavage.

As used herein, activation cleavage refers to the cleavage of the protease
30 at the N-terminus of the protease domain (generally between an R and I in the full-length protein, which includes amino acid residue 1 of SEQ ID No. 6, residue 463-492 of SEQ ID No. 23). By virtue of the Cys-Cys pairing between a Cys

-21-

outside the protease domain and a Cys in the protease domain (in this instance Cys₅₇₃ SEQ ID No. 23, upon cleavage the resulting polypeptide has two chains ("A" chain and the "B" chain, which is the protease domain). Cleavage can be effected by another protease or autocatalytically.

5 As used herein, a two-chain form of the protease domain refers to a two-chain form that is formed from the two-chain form of the protease in which the Cys pairing between, in this instance, a Cys outside the protease domain and Cys₅₇₃ (SEQ ID No. 23), which links the protease domain to the remainder of the polypeptide, the "A" chain. A two chain protease domain form refers to any
10 form in which the "remainder of the polypeptide", *i.e.*, "A" chain, is shortened and includes from at the Cys outside the protease domain. For example a two chain form of an MTSP10 includes from Cys₂₉₆ up to and including Cys₅₇₃ of SEQ ID No. 23 where the A chain includes Cys₂₉₆ to R₄₆₂ and the B chain includes I₄₆₃ to at least Cys₅₇₃.

15 MTSPs of interest include those that are activated and/or expressed in tumor cells different, typically higher, from those in non-tumor cells; and those from cells in which substrates therefor differ in tumor cells from non-tumor cells or differ with respect to the substrates, co-factors or receptors, or otherwise alter the activity or specificity of the MTSP.

20 As used herein, a human protein is one encoded by nucleic acid, such as DNA, present in the genome of a human, including all allelic variants and conservative variations as long as they are not variants found in other mammals.

 As used herein, a "nucleic acid encoding a protease domain or catalytically active portion of a SP" shall be construed as referring to a nucleic
25 acid encoding only the recited single chain protease domain or active portion thereof, and not the other contiguous portions of the SP as a continuous sequence.

 As used herein, catalytic activity refers to the activity of the SP as a serine protease. Function of the SP refers to its function in tumor biology,
30 including promotion of or involvement in initiation, growth or progression of tumors, and also roles in signal transduction. Catalytic activity refers to the

-22-

activity of the SP as a protease as assessed in *in vitro* proteolytic assays that detect proteolysis of a selected substrate.

As used herein, a CUB domain is a motif that mediates protein-protein interactions in complement components C1r/C1s and has also been identified in various proteins involved in developmental processes.

As used herein, "LDLR" refers to a low density lipoprotein receptor domain, which mediates binding to an LDL receptor.

As used herein, a zymogen is an inactive precursor of a proteolytic enzyme. Such precursors are generally larger, although not necessarily larger than the active form. With reference to serine proteases, zymogens are converted to active enzymes by specific cleavage, including catalytic and autocatalytic cleavage, or by binding of an activating co-factor, which generates an active enzyme. A zymogen, thus, is an enzymatically inactive protein that is converted to a proteolytic enzyme by the action of an activator.

As used herein, "disease or disorder" refers to a pathological condition in an organism resulting from, *e.g.*, infection or genetic defect, and characterized by identifiable symptoms.

As used herein, neoplasm (neoplasia) refers to abnormal new growth, and thus means the same as *tumor*, which can be benign or malignant. Unlike *hyperplasia*, neoplastic proliferation persists even in the absence of the original stimulus.

As used herein, neoplastic disease refers to any disorder involving cancer, including tumor development, growth, metastasis and progression.

As used herein, cancer refers to a general term for diseases caused by any type of malignant tumor.

As used herein, malignant, as applies to tumors, refers to primary tumors that have the capacity of *metastasis* with loss of *growth control* and *positional control*.

As used herein, an anti-cancer agent (used interchangeably with "anti-tumor or anti-neoplastic agent") refers to any agents used in the anti-cancer treatment. These include any agents, when used alone or in combination with other compounds, that can alleviate, reduce, ameliorate, prevent, or place or

RECTIFIED SHEET (RULE 91)

-23-

maintain in a state of remission of clinical symptoms or diagnostic markers associated with neoplastic disease, tumor and cancer, and can be used in methods, combinations and compositions provided herein. Non-limiting examples of anti-neoplastic agents include anti-angiogenic agents, alkylating
5 agents, antimetabolites, certain natural products, platinum coordination complexes, anthracenediones, substituted ureas, methylhydrazine derivatives, adrenocortical suppressants, certain hormones, antagonists and anti-cancer polysaccharides.

As used herein, a splice variant refers to a variant produced by differential
10 processing of a primary transcript of genomic nucleic acid, such as DNA, that results in more than one type of mRNA. Splice variants of SPs are provided herein.

As used herein, angiogenesis is intended to broadly encompass the totality of processes directly or indirectly involved in the establishment and
15 maintenance of new vasculature (neovascularization), including, but not limited to, neovascularization associated with tumors.

As used herein, anti-angiogenic treatment or agent refers to any therapeutic regimen and compound, when used alone or in combination with other treatment or compounds, that can alleviate, reduce, ameliorate, prevent, or
20 place or maintain in a state of remission of clinical symptoms or diagnostic markers associated with undesired and/or uncontrolled angiogenesis. Thus, for purposes herein an anti-angiogenic agent refers to an agent that inhibits the establishment or maintenance of vasculature. Such agents include, but are not limited to, anti-tumor agents, and agents for treatments of other disorders
25 associated with undesirable angiogenesis, such as diabetic retinopathies, restenosis, hyperproliferative disorders and others.

As used herein, non-anti-angiogenic anti-tumor agents refer to anti-tumor agents that do not act primarily by inhibiting angiogenesis.

As used herein, pro-angiogenic agents are agents that promote the
30 establishment or maintenance of the vasculature. Such agents include agents for treating cardiovascular disorders, including heart attacks and strokes.

RECTIFIED SHEET (RULE 91)

-24-

As used herein, undesired and/or uncontrolled angiogenesis refers to pathological angiogenesis wherein the influence of angiogenesis stimulators outweighs the influence of angiogenesis inhibitors. As used herein, deficient angiogenesis refers to pathological angiogenesis associated with disorders where
5 there is a defect in normal angiogenesis resulting in aberrant angiogenesis or an absence or substantial reduction in angiogenesis.

As used herein, the protease domain of an SP protein refers to the protease domain of an SP that exhibits proteolytic activity. Hence it is at least the minimal portion of the protein that exhibits proteolytic activity as assessed
10 by standard assays *in vitro*. It refers, herein, to a single chain form and also the two chain activated form (where the two chain form is intended it will be so-noted). Exemplary protease domains include at least a sufficient portion of sequences of amino acids set forth in SEQ ID No. 6 (amino acids 1-230, encoded by nucleotides in SEQ ID No. 5) to exhibit protease activity.

Also contemplated are nucleic acid molecules that encode a polypeptide
15 that has proteolytic activity in an *in vitro* proteolysis assay and that have at least 60%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% sequence identity with the full-length or a protease domain of an MTSP10 polypeptide or
20 other domain thereof, or that hybridize along their full-length or along at least about 70%, 80% or 90% of the full-length to a nucleic acids that encode a protease domain or other domain, particularly under conditions of moderate, generally high, stringency.

For the protease domains, residues at the N-terminus can be critical for
25 activity. It is shown herein that the protease domain of the single chain form of the MTSP10 protease is catalytically active. Hence the protease domain generally requires the N-terminal amino acids thereof for activity; the C-terminus portion can be truncated. The amount that can be removed can be determined empirically by testing the polypeptide for protease activity in an *in vitro* assay
30 that assesses catalytic cleavage.

Hence smaller portions of the protease domains, particularly the single chain domains, thereof that retain protease activity are contemplated. Such

RECTIFIED SHEET (RULE 91)

-25-

smaller versions generally are C-terminal truncated versions of the protease domains. Such domains exhibit conserved structure, including at least one structural feature, such as the active site triad, primary specificity pocket, oxyanion hole and/or other features of serine protease domains of proteases.

5 Thus, for purposes herein, the protease domain is a single chain portion of an MTSP10, as defined herein, but is homologous in its structural features and retention of sequence of similarity or homology the protease domain of chymotrypsin or trypsin. The polypeptide exhibits proteolytic activity as a single chain.

10 As used herein, "homologous" means about greater than 25% nucleic acid sequence identity, such as 25%, 40%, 60%, 70%, 80%, 90% or 95%. If necessary the percentage homology will be specified. The terms "homology" and "identity" are often used interchangeably. In general, sequences are aligned so that the highest order match is obtained (see, e.g.: *Computational Molecular*
15 *Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data, Part I*, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press, 1987; and
20 *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; Carillo *et al.* (1988) *SIAM J Applied Math* 48:1073). By sequence identity, the number of conserved amino acids are determined by standard alignment algorithms programs, and are used with default gap penalties established by each supplier. Substantially homologous nucleic acid molecules
25 would hybridize typically at moderate stringency or at high stringency all along the length of the nucleic acid or along at least about 70%, 80% or 90% of the full-length nucleic acid molecule of interest. Also contemplated are nucleic acid molecules that contain degenerate codons in place of codons in the hybridizing nucleic acid molecule.

30 Whether any two nucleic acid molecules have nucleotide sequences that are at least, for example, 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% "identical" can be determined using known computer algorithms such as the

RECTIFIED SHEET (RULE 91)

-26-

"FAST A" program, using for example, the default parameters as in Pearson *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:2444 (other programs include the GCG program package (Devereux, J., *et al.*, *Nucleic Acids Research* 12(1):387 (1984)), BLASTP, BLASTN, FASTA (Atschul, S.F., *et al.*, *J Molec Biol* 215:403 (1990); Guide to Huge Computers, Martin J. Bishop, ed., Academic Press, San Diego, 1994, and Carillo *et al.* (1988) *SIAM J Applied Math* 48:1073). For example, the BLAST function of the National Center for Biotechnology Information database can be used to determine identity. Other commercially or publicly available programs include, DNASTar "MegAlign" program (Madison, WI) and the University of Wisconsin Genetics Computer Group (UWG) "Gap" program (Madison WI). Percent homology or identity of proteins and/or nucleic acid molecules can be determined, for example, by comparing sequence information using a GAP computer program (*e.g.*, Needleman *et al.* (1970) *J. Mol. Biol.* 48:443, as revised by Smith and Waterman ((1981) *Adv. Appl. Math.* 2:482). Briefly, the GAP program defines similarity as the number of aligned symbols (*i.e.*, nucleotides or amino acids) which are similar, divided by the total number of symbols in the shorter of the two sequences. Default parameters for the GAP program can include: (1) a unary comparison matrix (containing a value of 1 for identities and 0 for non-identities) and the weighted comparison matrix of Gribskov *et al.* (1986) *Nucl. Acids Res.* 14:6745, as described by Schwartz and Dayhoff, eds., *ATLAS OF PROTEIN SEQUENCE AND STRUCTURE*, National Biomedical Research Foundation, pp. 353-358 (1979); (2) a penalty of 3.0 for each gap and an additional 0.10 penalty for each symbol in each gap; and (3) no penalty for end gaps. Therefore, as used herein, the term "identity" represents a comparison between a test and a reference polypeptide or polynucleotide.

As used herein, the term "at least 90% identical to" refers to percent identities from 90 to 99.99 relative to the reference polypeptides. Identity at a level of 90% or more is indicative of the fact that, assuming for exemplification purposes a test and reference polynucleotide length of 100 amino acids are compared, no more than 10% (*i.e.*, 10 out of 100) of amino acids in the test polypeptide differs from that of the reference polypeptides. Similar comparisons can be made between a test and reference polynucleotides. Such differences

RECTIFIED SHEET (RULE 91)

-27-

can be represented as point mutations randomly distributed over the entire length of an amino acid sequence or they can be clustered in one or more locations of varying length up to the maximum allowable, e.g. 10/100 amino acid difference (approximately 90% identity). Differences are defined as nucleic acid or amino acid substitutions, or deletions. At the level of homologies or identities above about 85-90%, the result should be independent of the program and gap parameters set; such high levels of identity can be assessed readily, often without relying on software.

As used herein, primer refers to an oligonucleotide containing two or more deoxyribonucleotides or ribonucleotides, typically more than three, from which synthesis of a primer extension product can be initiated. Experimental conditions conducive to synthesis include the presence of nucleoside triphosphates and an agent for polymerization and extension, such as DNA polymerase, and a suitable buffer, temperature and pH.

As used herein, animals include any animal, such as, but are not limited to, goats, cows, deer, sheep, rodents, pigs and humans. Non-human animals exclude humans as the contemplated animal. The SPs provided herein are from any source, animal, plant, prokaryotic and fungal. Most MTSP10s are of animal origin, including mammalian origin.

As used herein, genetic therapy involves the transfer of heterologous nucleic acid, such as DNA, into certain cells, target cells, of a mammal, particularly a human, with a disorder or conditions for which such therapy is sought. The nucleic acid, such as DNA, is introduced into the selected target cells in a manner such that the heterologous nucleic acid, such as DNA, is expressed and a therapeutic product encoded thereby is produced. Alternatively, the heterologous nucleic acid, such as DNA, can in some manner mediate expression of DNA that encodes the therapeutic product, or it can encode a product, such as a peptide or RNA that in some manner mediates, directly or indirectly, expression of a therapeutic product. Genetic therapy can also be used to deliver nucleic acid encoding a gene product that replaces a defective gene or supplements a gene product produced by the mammal or the cell in which it is introduced. The introduced nucleic acid can encode a

RECTIFIED SHEET (RULE 91)

-28-

therapeutic compound, such as a growth factor inhibitor thereof, or a tumor necrosis factor or inhibitor thereof, such as a receptor therefor, that is not normally produced in the mammalian host or that is not produced in therapeutically effective amounts or at a therapeutically useful time. The

5 heterologous nucleic acid, such as DNA, encoding the therapeutic product can be modified prior to introduction into the cells of the afflicted host in order to enhance or otherwise alter the product or expression thereof. Genetic therapy can also involve delivery of an inhibitor or repressor or other modulator of gene expression.

10 As used herein, heterologous nucleic acid is nucleic acid that (if DNA encodes RNA) encodes proteins that are not normally produced *in vivo* by the cell in which it is expressed or that mediates or encodes mediators that alter expression of endogenous nucleic acid, such as DNA, by affecting transcription, translation, or other regulatable biochemical processes. Heterologous nucleic
15 acid, such as DNA, can also be referred to as foreign nucleic acid, such as DNA. Any nucleic acid, such as DNA, that one of skill in the art would recognize or consider as heterologous or foreign to the cell in which the nucleic acid is expressed is herein encompassed by heterologous nucleic acid; heterologous nucleic acid includes exogenously added nucleic acid that is also expressed
20 endogenously. Examples of heterologous nucleic acid include, but are not limited to, nucleic acid that encodes traceable marker proteins, such as a protein that confers drug resistance, nucleic acid that encodes therapeutically effective substances, such as anti-cancer agents, enzymes and hormones, and nucleic acid, such as DNA, that encodes other types of proteins, such as antibodies.
25 Antibodies that are encoded by heterologous nucleic acid can be secreted or expressed on the surface of the cell in which the heterologous nucleic acid has been introduced. Heterologous nucleic acid is generally not endogenous to the cell into which it is introduced, but has been obtained from another cell or prepared synthetically. Generally, although not necessarily, such nucleic acid
30 encodes RNA and proteins that are not normally produced by the cell in which it is now expressed.

RECTIFIED SHEET (RULE 91)

-29-

introduction of the nucleic acid into a host, a product is expressed that ameliorates or eliminates the symptoms, manifestations of an inherited or acquired disease or that cures the disease. Also included are biologically active nucleic acid molecules, such as RNAi and antisense.

5 As used herein, recitation that a polypeptide consists essentially of the protease domain means that the only SP portion of the polypeptide is a protease domain or a catalytically active portion thereof. The polypeptide can optionally, and generally will, include additional non-SP-derived sequences of amino acids.

10 As used herein, cancer or tumor treatment or agent refers to any therapeutic regimen and/or compound that, when used alone or in combination with other treatments or compounds, can alleviate, reduce, ameliorate, prevent, or place or maintain in a state of remission of clinical symptoms or diagnostic markers associated with deficient angiogenesis.

15 As used herein, domain refers to a portion of a molecule, *e.g.*, protein or the encoding nucleic acid, that is structurally and/or functionally distinct from other portions of the molecule.

20 As used herein, protease refers to an enzyme catalyzing hydrolysis of proteins or peptides. It includes the zymogen form and activated forms thereof. For clarity, reference to protease refers to all forms, and particular forms will be specifically designated. For purposes herein, the protease domain includes single and two chain forms of the protease domain of an SP protein. For MTSP10 the protease domain also includes single and two chain forms of the protease domain.

25 As used herein, nucleic acids include DNA, RNA and analogs thereof, including peptide nucleic acids (PNA) and mixtures thereof. Nucleic acids can be single or double-stranded. When referring to probes or primers, optionally labeled, with a detectable label, such as a fluorescent or radiolabel, single-stranded molecules are contemplated. Such molecules are typically of a length such that their target is statistically unique or of low copy number (typically less
30 than 5, generally less than 3) for probing or priming a library. Generally a probe or primer contains at least 14, 16 or 30 contiguous of sequence complementary

RECTIFIED SHEET (RULE 91)

-30-

to or identical a gene of interest. Probes and primers can be 10, 20, 30, 50, 100 or more nucleic acids long.

As used herein, nucleic acid encoding a fragment or portion of an SP refers to a nucleic acid encoding only the recited fragment or portion of SP, and
5 not the other contiguous portions of the SP.

As used herein, operative linkage of heterologous nucleic acids to regulatory and effector sequences of nucleotides, such as promoters, enhancers, transcriptional and translational stop sites, and other signal sequences refers to the relationship between such nucleic acid, such as DNA, and such sequences of
10 nucleotides. Thus, operatively linked or operationally associated refers to the functional relationship of nucleic acid, such as DNA, with regulatory and effector sequences of nucleotides, such as promoters, enhancers, transcriptional and translational stop sites, and other signal sequences. For example, operative linkage of DNA to a promoter refers to the physical and functional relationship
15 between the DNA and the promoter such that the transcription of such DNA is initiated from the promoter by an RNA polymerase that specifically recognizes, binds to and transcribes the DNA. In order to optimize expression and/or *in vitro* transcription, it can be necessary to remove, add or alter 5' untranslated portions of the clones to eliminate extra, potential inappropriate alternative translation
20 initiation (*i.e.*, start) codons or other sequences that can interfere with or reduce expression, either at the level of transcription or translation. Alternatively, consensus ribosome binding sites (see, *e.g.*, Kozak *J. Biol. Chem.* 266:19867-19870 (1991)) can be inserted immediately 5' of the start codon and can enhance expression. The desirability of (or need for) such modification can be
25 empirically determined.

As used herein, a sequence complementary to at least a portion of an RNA, with reference to antisense oligonucleotides, means a sequence having sufficient complementarity to be able to hybridize with the RNA, generally under moderate or high stringency conditions, forming a stable duplex; in the case of
30 double-stranded SP antisense nucleic acids, a single strand of the duplex DNA (or dsRNA) can thus be tested, or triplex formation can be assayed. The ability to hybridize depends on the degree of complementarity and the length of the

-31-

antisense nucleic acid. Generally, the longer the hybridizing nucleic acid, the more base mismatches with a SP encoding RNA it can contain and still form a stable duplex (or triplex, as the case can be). One skilled in the art can ascertain a tolerable degree of mismatch by use of standard procedures to determine the melting point of the hybridized complex.

For purposes herein, amino acid substitutions can be made in any of SPs and protease domains thereof provided that the resulting protein exhibits protease activity. Amino acid substitutions contemplated include conservative substitutions, such as those set forth in Table 1, which do not eliminate proteolytic activity. As described herein, substitutions that alter properties of the proteins, such as removal of cleavage sites and other such sites are also contemplated; such substitutions are generally non-conservative, but can be readily effected by those of skill in the art.

Suitable conservative substitutions of amino acids are known to those of skill in this art and can be made generally without altering the biological activity, for example enzymatic activity, of the resulting molecule. Those of skill in this art recognize that, in general, single amino acid substitutions in non-essential regions of a polypeptide do not substantially alter biological activity (see, *e.g.*, Watson *et al. Molecular Biology of the Gene*, 4th Edition, 1987, The Bejacmin/Cummings Pub. co., p.224). Also included within the definition, is the catalytically active fragment of an SP, particularly a single chain protease portion. Conservative amino acid substitutions are made, for example, in accordance with those set forth in TABLE 1 as follows:

TABLE 1

	Original residue	Conservative substitution
25	Ala (A)	Gly; Ser, Abu
	Arg (R)	Lys, orn
	Asn (N)	Gln; His
	Cys (C)	Ser
30	Gln (Q)	Asn
	Glu (E)	Asp
	Gly (G)	Ala; Pro
	His (H)	Asn; Gln
	Ile (I)	Leu; Val; Met; Nle; Nva
35	Leu (L)	Ile; Val; Met; Nle; Nv
	Lys (K)	Arg; Gln; Glu

-32-

Original residue	Conservative substitution
Met (M)	Leu; Tyr; Ile; NLe Val
Ornithine	Lys; Arg
Phe (F)	Met; Leu; Tyr
Ser (S)	Thr
Thr (T)	Ser
Trp (W)	Tyr
Tyr (Y)	Trp; Phe
Val (V)	Ile; Leu; Met; Nle; Nv

Other substitutions are also permissible and can be determined empirically or in
 10 accord with known conservative substitutions.

As used herein, Abu is 2-aminobutyric acid; Orn is ornithine.

As used herein, the amino acids, which occur in the various amino acid
 sequences appearing herein, are identified according to their well-known, three-
 letter or one-letter abbreviations. The nucleotides, which occur in the various
 15 DNA fragments, are designated with the standard single-letter designations used
 routinely in the art.

As used herein, a probe or primer based on a nucleotide sequence
 disclosed herein, includes at least 10, 14, typically at least 16 contiguous
 sequence of nucleotides of SEQ ID No. 5 or SEQ ID No. 22, and probes of at
 20 least 30, 50 or 100 contiguous sequence of nucleotides of SEQ ID No. 5 or SEQ
 ID No. 22. The length of the probe or primer for unique hybridization is a
 function of the complexity of the genome of interest.

As used herein, amelioration of the symptoms of a particular disorder by
 administration of a particular pharmaceutical composition refers to any lessening,
 25 whether permanent or temporary, lasting or transient that can be attributed to or
 associated with administration of the composition.

As used herein, antisense polynucleotides refer to synthetic sequences of
 nucleotide bases complementary to mRNA or the sense strand of double-
 stranded DNA. Admixture of sense and antisense polynucleotides under
 30 appropriate conditions leads to the binding of the two molecules, or
 hybridization. When these polynucleotides bind to (hybridize with) mRNA,
 inhibition of protein synthesis (translation) occurs. When these polynucleotides
 bind to double-stranded DNA, inhibition of RNA synthesis (transcription) occurs.
 The resulting inhibition of translation and/or transcription leads to an inhibition of

RECTIFIED SHEET (RULE 91)

-33-

the synthesis of the protein encoded by the sense strand. Antisense nucleic acid molecules typically contain a sufficient number of nucleotides to specifically bind to a target nucleic acid, generally at least 5 contiguous nucleotides, often at least 14 or 16 or 30 contiguous nucleotides or modified nucleotides

- 5 complementary to the coding portion of a nucleic acid molecule that encodes a gene of interest, for example, nucleic acid encoding a single chain protease domain of an SP.

As used herein, an array refers to a collection of elements, such as antibodies, containing three or more members. An addressable array is one in
10 which the members of the array are identifiable, typically by position on a solid phase support. Hence, in general the members of the array are immobilized on discrete identifiable loci on the surface of a solid phase.

As used herein, antibody refers to an immunoglobulin, whether natural or partially or wholly synthetically produced, including any derivative thereof that
15 retains the specific binding ability the antibody. Hence antibody includes any protein having a binding domain that is homologous or substantially homologous to an immunoglobulin binding domain. Antibodies include members of any immunoglobulin claims, including IgG, IgM, IgA, IgD and IgE.

As used herein, antibody fragment refers to any derivative of an antibody
20 that is less than full-length, retaining at least a portion of the full-length antibody's specific binding ability. Examples of antibody fragments include, but are not limited to, Fab, Fab', F(ab)₂, single-chain Fvs (scFV), FV, dsFV diabody and Fd fragments. The fragment can include multiple chains linked together, such as by disulfide bridges. An antibody fragment generally contains at least
25 about 50 amino acids and typically at least 200 amino acids.

As used herein, a Fv antibody fragment is composed of one variable heavy domain (V_H) and one variable light domain linked by noncovalent interactions.

As used herein, a dsFV refers to a Fv with an engineered intermolecular
30 disulfide bond, which stabilizes the V_H-V_L pair.

RECTIFIED SHEET (RULE 91)

-34-

As used herein, a $F(ab)_2$ fragment is an antibody fragment that results from digestion of an immunoglobulin with pepsin at pH 4.0-4.5; it can be recombinantly expressed to produce the equivalent fragment.

As used herein, Fab fragments are antibody fragments that result from digestion of an immunoglobulin with papain; they can be recombinantly expressed to produce the equivalent fragment.

As used herein, scFVs refer to antibody fragments that contain a variable light chain (V_L) and variable heavy chain (V_H) covalently connected by a polypeptide linker in any order. The linker is of a length such that the two variable domains are bridged without substantial interference. Included linkers are $(Gly-Ser)_n$ residues with some Glu or Lys residues dispersed throughout to increase solubility.

As used herein, humanized antibodies refer to antibodies that are modified to include human sequences of amino acids so that administration to a human does not provoke an immune response. Methods for preparation of such antibodies are known. For example, to produce such antibodies, the encoding nucleic acid in the hybridoma or other prokaryotic or eukaryotic cell, such as an *E. coli* or a CHO cell, that expresses the monoclonal antibody is altered by recombinant nucleic acid techniques to express an antibody in which the amino acid composition of the non-variable region is based on human antibodies. Computer programs have been designed to identify such non-variable regions.

As used herein, diabodies are dimeric scFV; diabodies typically have shorter peptide linkers than scFvs, and they generally dimerize.

As used herein, production by recombinant means by using recombinant DNA methods means the use of the well known methods of molecular biology for expressing proteins encoded by cloned DNA.

As used herein the term assessing is intended to include quantitative and qualitative determination in the sense of obtaining an absolute value for the activity of an SP, or a domain thereof, present in the sample, and also of obtaining an index, ratio, percentage, visual or other value indicative of the level of the activity. Assessment can be direct or indirect and the chemical species actually detected need not of course be the proteolysis

RECTIFIED SHEET (RULE 91)

-35-

product itself but can for example be a derivative thereof or some further substance.

As used herein, biological activity refers to the *in vivo* activities of a compound or physiological responses that result upon *in vivo* administration of a compound, composition or other mixture. Biological activity, thus, encompasses therapeutic effects and pharmaceutical activity of such compounds, compositions and mixtures. Biological activities can be observed in *in vitro* systems designed to test or use such activities. Thus, for purposes herein the biological activity of a luciferase is its oxygenase activity whereby, upon oxidation of a substrate, light is produced.

As used herein, functional activity refers to a polypeptide or portion thereof that displays one or more activities associated with a full-length protein. Functional activities include, but are not limited to, biological activity, catalytic or enzymatic activity, antigenicity (ability to bind to or compete with a polypeptide for binding to an anti-polypeptide antibody), immunogenicity, ability to form multimers, and the ability to specifically bind to a receptor or ligand for the polypeptide.

As used herein, a conjugate refers to the compounds provided herein that include one or more SPs, including an MTSP10, particularly single chain protease domains thereof, and one or more targeting agents. These conjugates include those produced by recombinant means as fusion proteins, those produced by chemical means, such as by chemical coupling, through, for example, coupling to sulfhydryl groups, and those produced by any other method whereby at least one SP, or a domain thereof, is linked, directly or indirectly via linker(s) to a targeting agent.

As used herein, a targeting agent is any moiety, such as a protein or effective portion thereof, that provides specific binding of the conjugate to a cell surface receptor, which, can internalize the conjugate or SP portion thereof. A targeting agent can also be one that promotes or facilitates, for example, affinity isolation or purification of the conjugate; attachment of the conjugate to a surface; or detection of the conjugate or complexes containing the conjugate.

RECTIFIED SHEET (RULE 91)

-36-

As used herein, an antibody conjugate refers to a conjugate in which the targeting agent is an antibody.

As used herein, derivative or analog of a molecule refers to a portion derived from or a modified version of the molecule.

5 As used herein, an effective amount of a compound for treating a particular disease is an amount that is sufficient to ameliorate, or in some manner reduce the symptoms associated with the disease. Such an amount can be administered as a single dosage or can be administered according to a regimen, whereby it is effective. The amount can cure the disease but, typically,
10 is administered in order to ameliorate the symptoms of the disease. Repeated administration can be required to achieve the desired amelioration of symptoms.

As used herein, equivalent, when referring to two sequences of nucleic acids, means that the two sequences in question encode the same sequence of amino acids or equivalent proteins. When equivalent is used in referring to two
15 proteins or peptides, it means that the two proteins or peptides have substantially the same amino acid sequence with only amino acid substitutions (such as, but not limited to, conservative changes such as those set forth in Table 1 above) that do not substantially alter the activity or function of the protein or peptide. When equivalent refers to a property, the property does not
20 need to be present to the same extent (e.g., two peptides can exhibit different rates of the same type of enzymatic activity), but the activities are usually substantially the same. Complementary, when referring to two nucleotide sequences, means that the two sequences of nucleotides are capable of hybridizing, typically with less than 25%, 15%, 5% or 0% mismatches between
25 opposed nucleotides. If necessary, the percentage of complementarity will be specified. Typically the two molecules are selected such that they will hybridize under conditions of high stringency.

As used herein, an agent that modulates the activity of a protein or expression of a gene or nucleic acid either decreases or increases or otherwise
30 alters the activity of the protein or, in some manner, up- or down-regulates or otherwise alters expression of the nucleic acid in a cell.

RECTIFIED SHEET (RULE 91)

-37-

As used herein, inhibitor of the activity of an SP encompasses any substance that prohibits or decrease production, post-translational modification(s), maturation, or membrane localization of the SP or any substance that interferes with or decreases the proteolytic efficacy of thereof, particularly
5 of a single chain form in an *in vitro* screening assay.

As used herein, a method for treating or preventing neoplastic disease means that any of the symptoms, such as the tumor, metastasis thereof, the vascularization of the tumors or other parameters by which the disease is characterized are reduced, ameliorated, prevented, placed in a state of remission,
10 or maintained in a state of remission. It also means that the hallmarks of neoplastic disease and metastasis can be eliminated, reduced or prevented by the treatment. Non-limiting examples of the hallmarks include uncontrolled degradation of the basement membrane and proximal extracellular matrix, migration, division, and organization of the endothelial cells into new functioning
15 capillaries, and the persistence of such functioning capillaries.

As used herein, pharmaceutically acceptable salts, esters or other derivatives of the conjugates include any salts, esters or derivatives that can be readily prepared by those of skill in this art using known methods for such derivatization and that produce compounds that can be administered to animals
20 or humans without substantial toxic effects and that either are pharmaceutically active or are prodrugs.

As used herein, a prodrug is a compound that, upon *in vivo* administration, is metabolized or otherwise converted to the biologically, pharmaceutically or therapeutically active form of the compound. To produce a
25 prodrug, the pharmaceutically active compound is modified such that the active compound is regenerated by metabolic processes. The prodrug can be designed to alter the metabolic stability or the transport characteristics of a drug, to mask side effects or toxicity, to improve the flavor of a drug or to alter other characteristics or properties of a drug. By virtue of knowledge of
30 pharmacodynamic processes and drug metabolism *in vivo*, those of skill in this art, once a pharmaceutically active compound is known, can design prodrugs of

RECTIFIED SHEET (RULE 91)

-38-

the compound (see, e.g., Nogrady (1985) *Medicinal Chemistry A Biochemical Approach*, Oxford University Press, New York, pages 388-392).

As used herein, a drug identified by the screening methods provided herein refers to any compound that is a candidate for use as a therapeutic or as
5 a lead compound for the design of a therapeutic. Such compounds can be small molecules, including small organic molecules, peptides, peptide mimetics, antisense molecules or dsRNA, such as RNAi, antibodies, fragments of antibodies, recombinant antibodies and other such compounds that can serve as drug candidates or lead compounds.

10 As used herein, a peptidomimetic is a compound that mimics the conformation and certain stereochemical features of the biologically active form of a particular peptide. In general, peptidomimetics are designed to mimic certain desirable properties of a compound, but not the undesirable properties, such as flexibility, that lead to a loss of a biologically active conformation and
15 bond breakdown. Peptidomimetics may be prepared from biologically active compounds by replacing certain groups or bonds that contribute to the undesirable properties with bioisosteres. Bioisosteres are known to those of skill in the art. For example the methylene bioisostere CH_2S has been used as an amide replacement in enkephalin analogs (see, e.g., Spatola (1983) pp. 267-357
20 in Chemistry and Biochemistry of Amino Acids, Peptides, and Proteins, Weistein, Ed. volume 7, Marcel Dekker, New York). Morphine, which can be administered orally, is a compound that is a peptidomimetic of the peptide endorphin. For purposes herein, cyclic peptides are included among peptidomimetics.

As used herein, a promoter region or promoter element refers to a
25 segment of DNA or RNA that controls transcription of the DNA or RNA to which it is operatively linked. The promoter region includes specific sequences that are sufficient for RNA polymerase recognition, binding and transcription initiation. This portion of the promoter region is referred to as the promoter. In addition, the promoter region includes sequences that modulate this recognition, binding
30 and transcription initiation activity of RNA polymerase. These sequences can be *cis* acting or can be responsive to *trans* acting factors. Promoters, depending upon the nature of the regulation, can be constitutive or regulated. Exemplary

RECTIFIED SHEET (RULE 91)

-39-

promoters contemplated for use in prokaryotes include the bacteriophage T7 and T3 promoters.

As used herein, a receptor refers to a molecule that has an affinity for a given ligand. Receptors can be naturally-occurring or synthetic molecules.

- 5 Receptors can also be referred to in the art as anti-ligands. As used herein, the receptor and anti-ligand are interchangeable. Receptors can be used in their unaltered state or as aggregates with other species. Receptors can be attached to, covalently or noncovalently, or in physical contact with, a binding member, either directly or indirectly via a specific binding substance or linker. Examples
- 10 of receptors, include, but are not limited to: antibodies, cell membrane receptors surface receptors and internalizing receptors, monoclonal antibodies and antisera reactive with specific antigenic determinants [such as on viruses, cells, or other materials], drugs, polynucleotides, nucleic acids, peptides, cofactors, lectins, sugars, polysaccharides, cells, cellular membranes, and organelles.

- 15 Examples of receptors and applications using such receptors, include but are not restricted to:

a) enzymes: specific transport proteins or enzymes essential to survival of microorganisms, which could serve as targets for antibiotic [ligand] selection;

- b) antibodies: identification of a ligand-binding site on the antibody
- 20 molecule that combines with the epitope of an antigen of interest can be investigated; determination of a sequence that mimics an antigenic epitope can lead to the development of vaccines of which the immunogen is based on one or more of such sequences or lead to the development of related diagnostic agents or compounds useful in therapeutic treatments such as for auto-immune diseases

- 25 c) nucleic acids: identification of ligand, such as protein or RNA, binding sites;

- d) catalytic polypeptides: polymers, including polypeptides, that are capable of promoting a chemical reaction involving the conversion of one or more reactants to one or more products; such polypeptides generally include a
- 30 binding site specific for at least one reactant or reaction intermediate and an active functionality proximate to the binding site, in which the functionality is

-40-

capable of chemically modifying the bound reactant (see, e.g., U.S. Patent No. 5,215,899);

- e) hormone receptors: determination of the ligands that bind with high affinity to a receptor is useful in the development of hormone replacement therapies; for example, identification of ligands that bind to such receptors can lead to the development of drugs to control blood pressure; and
- f) opiate receptors: determination of ligands that bind to the opiate receptors in the brain is useful in the development of less-addictive replacements for morphine and related drugs.

10 As used herein, sample refers to anything that contains an analyte for which an analyte assay is desired. The sample can be a biological sample, such as a biological fluid or a biological tissue. Examples of biological fluids include urine, blood, plasma, serum, saliva, semen, stool, sputum, cerebral spinal fluid, tears, mucus, sperm, amniotic fluid or the like. Biological tissues are aggregates
15 of cells, usually of a particular kind together with their intercellular substance that form one of the structural materials of a human, animal, plant, bacterial, fungal or viral structure, including connective, epithelium, muscle and nerve tissues. Examples of biological tissues also include organs, tumors, lymph nodes, arteries and individual cell(s).

20 As used herein: stringency of hybridization in determining percentage mismatch is as follows:

- 1) high stringency: 0.1 x SSPE, 0.1% SDS, 65°C
- 2) medium stringency: 0.2 x SSPE, 0.1% SDS, 50°C
- 3) low stringency: 1.0 x SSPE, 0.1% SDS, 50°C

25 Those of skill in this art know that the washing step selects for stable hybrids and also know the ingredients of SSPE (see, e.g., Sambrook, E.F. Fritsch, T. Maniatis, in: *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Laboratory Press (1989), vol. 3, p. B.13, see, also, numerous catalogs that describe commonly used laboratory solutions). SSPE is pH 7.4 phosphate-
30 buffered 0.18 M NaCl. Further, those of skill in the art recognize that the stability of hybrids is determined by T_m , which is a function of the sodium ion concentration and temperature ($T_m = 81.5^\circ \text{C} - 16.6(\log_{10}[\text{Na}^+]) + 0.41(\% \text{G} + \text{C}) -$

RECTIFIED SHEET (RULE 91)

-41-

600/l)), so that the only parameters in the wash conditions critical to hybrid stability are sodium ion concentration in the SSPE (or SSC) and temperature.

It is understood that equivalent stringencies can be achieved using alternative buffers, salts and temperatures. By way of example and not

5 limitation, procedures using conditions of low stringency are as follows (see also Shilo and Weinberg, *Proc. Natl. Acad. Sci. USA* 78:6789-6792 (1981)): Filters containing DNA are pretreated for 6 hours at 40°C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH 7.5), 5 mM EDTA, 0.1% PVP, 0.1% Ficoll, 1% BSA, and 500 µg/ml denatured salmon sperm DNA (10X SSC is 1.5
10 M sodium chloride, and 0.15 M sodium citrate, adjusted to a pH of 7).

Hybridizations are carried out in the same solution with the following modifications: 0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100 µg/ml salmon sperm DNA, 10% (wt/vol) dextran sulfate, and 5-20 X 10⁶ cpm ³²P-labeled probe is used. Filters are incubated in hybridization mixture for 18-20 hours at 40°C,
15 and then washed for 1.5 hours at 55°C in a solution containing 2X SSC, 25 mM Tris-HCl (pH 7.4), 5 mM EDTA, and 0.1% SDS. The wash solution is replaced with fresh solution and incubated an additional 1.5 hours at 60°C. Filters are blotted dry and exposed for autoradiography. If necessary, filters are washed for a third time at 65-68°C and reexposed to film. Other conditions of low
20 stringency which can be used are well known in the art (*e.g.*, as employed for cross-species hybridizations).

By way of example and not way of limitation, procedures using conditions of moderate stringency include, for example, but are not limited to, procedures using such conditions of moderate stringency are as follows: Filters
25 containing DNA are pretreated for 6 hours at 55°C in a solution containing 6X SSC, 5X Denhart's solution, 0.5% SDS and 100 µg/ml denatured salmon sperm DNA. Hybridizations are carried out in the same solution and 5-20 X 10⁶ cpm ³²P-labeled probe is used. Filters are incubated in hybridization mixture for 18-20 hours at 55°C, and then washed twice for 30 minutes at 60°C in a solution
30 containing 1X SSC and 0.1% SDS. Filters are blotted dry and exposed for autoradiography. Other conditions of moderate stringency which can be used

-42-

are well-known in the art. Washing of filters is done at 37°C for 1 hour in a solution containing 2X SSC, 0.1% SDS.

By way of example and not way of limitation, procedures using conditions of high stringency are as follows: Prehybridization of filters containing DNA is carried out for 8 hours to overnight at 65°C in buffer composed of 6X SSC, 50 mM Tris-HCl (pH 7.5), 1 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.02% BSA, and 500 µg/ml denatured salmon sperm DNA. Filters are hybridized for 48 hours at 65°C in prehybridization mixture containing 100 µg/ml denatured salmon sperm DNA and 5-20 X 10⁶ cpm of ³²P-labeled probe. Washing of filters is done at 37°C for 1 hour in a solution containing 2X SSC, 0.01% PVP, 0.01% Ficoll, and 0.01% BSA. This is followed by a wash in 0.1X SSC at 50°C for 45 minutes before autoradiography. Other conditions of high stringency which can be used are well known in the art.

The term substantially identical or substantially homologous or similar varies with the context as understood by those skilled in the relevant art and generally means at least 60% or 70%, preferably means at least 80%, 85% or more preferably at least 90%, and most preferably at least 95% identity.

As used herein, substantially identical to a product means sufficiently similar so that the property of interest is sufficiently unchanged so that the substantially identical product can be used in place of the product.

As used herein, substantially pure means sufficiently homogeneous to appear free of readily detectable impurities as determined by standard methods of analysis, such as thin layer chromatography (TLC), gel electrophoresis and high performance liquid chromatography (HPLC), used by those of skill in the art to assess such purity, or sufficiently pure such that further purification would not detectably alter the physical and chemical properties, such as enzymatic and biological activities, of the substance. Methods for purification of the compounds to produce substantially chemically pure compounds are known to those of skill in the art. A substantially chemically pure compound can, however, be a mixture of stereoisomers or isomers. In such instances, further purification might increase the specific activity of the compound.

As used herein, target cell refers to a cell that expresses an SP *in vivo*.

-43-

As used herein, test substance (or test compound) refers to a chemically defined compound (*e.g.*, organic molecules, inorganic molecules, organic/inorganic molecules, proteins, peptides, nucleic acids, oligonucleotides, lipids, polysaccharides, saccharides, or hybrids among these molecules such as glycoproteins, etc.) or mixtures of compounds (*e.g.*, a library of test compounds, natural extracts or culture supernatants, etc.) whose effect on an SP, particularly a single chain form that includes the protease domain or a sufficient portion thereof for activity, as determined by an *in vitro* method, such as the assays provided herein, is tested.

As used herein, the terms a therapeutic agent, therapeutic regimen, radioprotectant or chemotherapeutic mean conventional drugs and drug therapies, including vaccines, which are known to those skilled in the art. Radiotherapeutic agents are well known in the art.

As used herein, treatment means any manner in which the symptoms of a condition, disorder or disease are ameliorated or otherwise beneficially altered. Treatment also encompasses any pharmaceutical use of the compositions herein.

As used herein, vector (or plasmid) refers to discrete elements that are used to introduce heterologous nucleic acid into cells for either expression or replication thereof. The vectors typically remain episomal, but can be designed to effect integration of a gene or portion thereof into a chromosome of the genome. Also contemplated are vectors that are artificial chromosomes, such as yeast artificial chromosomes and mammalian artificial chromosomes. Selection and use of such vehicles are well known to those of skill in the art. An expression vector includes vectors capable of expressing DNA that is operatively linked with regulatory sequences, such as promoter regions, that are capable of effecting expression of such DNA fragments. Thus, an expression vector refers to a recombinant DNA or RNA construct, such as a plasmid, a phage, recombinant virus or other vector that, upon introduction into an appropriate host cell, results in expression of the cloned DNA. Appropriate expression vectors are well known to those of skill in the art and include those that are replicable in eukaryotic cells and/or prokaryotic cells and those that remain episomal or those which integrate into the host cell genome.

RECTIFIED SHEET (RULE 91)

-44-

As used herein, protein binding sequence refers to a protein or peptide sequence or a portion of other macromolecules that is capable of specific binding to protein or peptide sequences generally, to a set of protein or peptide sequences or to a particular protein or peptide sequence.

5 As used herein, epitope tag refers to a short stretch of amino acid residues corresponding to an epitope to facilitate subsequent biochemical and immunological analysis of the epitope tagged protein or peptide. Epitope tagging is achieved by including the sequence of the epitope tag to the protein-encoding sequence in an appropriate expression vector. Epitope tagged proteins can be
10 affinity purified using highly specific antibodies raised against the tags.

As used herein, metal binding sequence refers to a protein or peptide sequence that is capable of specific binding to metal ions generally, to a set of metal ions or to a particular metal ion.

15 As used herein, a combination refers to any association between two or among more items.

As used herein, a composition refers to any mixture. It can be a solution, a suspension, liquid, powder, a paste, aqueous, non-aqueous or any combination thereof.

20 As used herein, fluid refers to any composition that can flow. Fluids thus encompass compositions that are in the form of semi-solids, pastes, solutions, aqueous mixtures, gels, lotions, creams and other such compositions.

As used herein, a cellular extract refers to a preparation or fraction which is made from a lysed or disrupted cell.

25 As used herein, an agent is said to be randomly selected when the agent is chosen randomly without considering the specific sequences involved in the association of a protein alone or with its associated substrates, binding partners, etc. An example of randomly selected agents is the use a chemical library or a peptide combinatorial library, or a growth broth of an organism or conditioned medium.

30 As used herein, an agent is said to be rationally selected or designed when the agent is chosen on a non-random basis which takes into account the sequence of the target site and/or its conformation in connection with the

RECTIFIED SHEET (RULE 91)

-45-

agent's action. As described in the Examples, there are proposed binding sites for serine protease and (catalytic) sites in the protein having SEQ ID NO:2 or SEQ ID NO:4. Agents can be rationally selected or rationally designed by utilizing the peptide sequences that make up these sites. For example, a

5 rationally selected peptide agent can be a peptide whose amino acid sequence is identical to the ATP or calmodulin binding sites or domains.

For clarity of disclosure, and not by way of limitation, the detailed description is divided into the subsections that follow.

B. MTSP10 polypeptides, muteins, derivatives and analogs thereof

10 **MTSPs**

The MTSPs are a family of transmembrane serine proteases that are found in mammals and also other species. MTSPs are of interest because they appear to be expressed and/or activated at different levels in tumor cells from normal cells, or have functional activity that is different in tumor cells from

15 normal cells, such as by an alteration in a substrate therefor, or a cofactor or a receptor.

The MTSPs share a number of common structural features including: a proteolytic extracellular C-terminal domain; a transmembrane domain, with a hydrophobic domain near the N-terminus; a short cytoplasmic domain; and a

20 variable length stem region that may contain additional modular domains. The proteolytic domains share sequence homology including conserved His, Asp, and Ser residues necessary for catalytic activity that are present in conserved motifs. The MTSPs are normally synthesized as zymogens and can be activated to two-chain forms by cleavage. It is shown herein that a single chain proteolytic

25 domain can function *in vitro* and, hence is useful in *in vitro* assays for identifying agents that modulate the activity of members of this family.

For purposes herein, the protease domain of the MTSP does not have to result from activation cleavage, which produces a two chain activated product, but rather includes single chain polypeptides where the N-termini include the

30 consensus sequence ↓VVGG, ↓IVGG, ↓VGLL, ↓ILGG, ↓IVQG or ↓IVNG ↓IASG or other such motif. Such polypeptides, although not the result of activation cleavage and not two-chain forms, exhibit proteolytic (catalytic) activity. These

RECTIFIED SHEET (RULE 91)

-46-

protease domain polypeptides are used in assays to screen for agents that modulate the activity of the MTSP10.

The MTSP family is a target for therapeutic intervention and also some members can serve as diagnostic markers for tumor development, growth and/or progression. As discussed, the members of this family are involved in proteolytic processes that are implicated in tumor development, growth and/or progression. This implication is based upon their functions as proteolytic enzymes in processes related to ECM degradation and/or remodeling and activation of pro-growth factors, pro-hormones or pro-angiogenic compounds. In addition, their levels of expression or level of activation or their apparent activity resulting from substrate levels or alterations in substrates and levels thereof differs in tumor cells and non-tumor cells in the same tissue. Similarly the level of co-factors or receptors for these proteases can vary between tumor and non-tumor cells. Hence, protocols and treatments that alter their activity, such as their proteolytic activities and roles in signal transduction, and/or their expression, such as by contacting them with a compound that modulates their activity and/or expression, could impact tumor development, growth and/or progression. Also, in some instances, the level of activation and/or expression can be altered in tumors, such as lung carcinoma, colon adenocarcinoma and ovarian carcinoma.

MTSP10

MTSP10 is of interest because it is expressed or is active in tumor cells. The MTSP provided herein can serve as a diagnostic marker for particular tumors, by virtue of a level of activity and/or expression or function in a subject (i.e. a mammal, particularly a human) with neoplastic disease, compared to a subject or subjects that do not have the neoplastic disease. In addition, detection of activity (and/or expression) in a particular tissue can be indicative of neoplastic disease. It is shown herein, that MTSP10s provided herein are expressed and/or activated in certain tumors; hence their activation or expression can serve as a diagnostic marker for tumor development, growth and/or progression. In other instances, the MTSP polypeptide can exhibit altered activity by virtue of a change in activity or expression of a co-factor, a substrate

-47-

or a receptor. In addition, in some instances, these MTSPs and/or variants thereof can be shed from cell surfaces. Detection of the shed MTSPs, particularly the extracellular protease domains, in body fluids, such as serum, blood, saliva, cerebral spinal fluid, synovial fluid and interstitial fluids, urine, sweat and other such fluids and secretions, can serve as a diagnostic tumor marker. In particular, detection of higher levels of such shed polypeptides in a subject compared to a subject known not to have any neoplastic disease or compared to earlier samples from the same subject, can be indicative of neoplastic disease in the subject.

10 Polypeptides and muteins

Provided herein are isolated substantially pure single chain and two chain polypeptides that contain the protease domain of an MTSP10. The polypeptides also can include other non-MTSP sequences of amino acids, but includes the protease domain or a sufficient portion thereof to exhibit catalytic activity in any *in vitro* assay that assess such protease activity, such as any provided herein.

MTSP10 polypeptides provided herein are expressed or activated by or in tumor cells, typically at a level that differs from the level in which they are expressed by or activated in a non-tumor cell of the same type. Hence, for example, if the MTSP is expressed in an cervical tumor cell, it is expressed or active at a different level from the level in non-tumor cervical cells. MTSP10 expression or activation can be indicative of cervical, lung, esophageal, colon, prostate, uterine, pancreatic, breast and other tumors.

Isolated, substantially pure proteases that include protease domains or a catalytically active portion thereof are provided. Provided are single chain forms and two chain forms of the MTSP10. The protease domains can be included in a longer protein, and such longer protein is optionally the MTSP10 zymogen. Exemplary MTSP10-encoding nucleic acid and protein sequences of a protease domain are set forth in SEQ ID Nos. 5 and 6. Full-length MTSP10-encoding nucleic acid molecules that contain the sequence set forth in SEQ ID No. 5 or SEQ ID No. 22 and polypeptides that include the sequence of amino acids set forth as residues 1-230 in SEQ ID No. 6 or SEQ ID No. 23 or catalytically active portions thereof are also provided herein. Thus, an MTSP10 polypeptide

-48-

includes the sequence of amino acids set forth as residues 1-230 SEQ ID No. 6 or residues 463-492 of SEQ ID No. 23 are provided. Smaller portions thereof that retain protease activity are contemplated.

Substantially purified MTSP10 protease is encoded by a nucleic acid that
 5 hybridizes to a nucleic acid molecule containing the protease domain encoded by the sequence of nucleotides that encodes residues 1-230 set forth in SEQ. ID No. 6 under at least moderate, generally high, stringency conditions, such that the protease domain encoding nucleic acid thereof hybridizes along its full-length or at least 70%, 80% or 90% of the full-length. In certain
 10 embodiments the substantially purified MTSP protease is a single chain polypeptide that includes substantially the sequence of amino acids set forth as residues 1-230 in SEQ ID No. 6 or a catalytically active portion thereof.

Also included are substantially purified MTSP10 zymogens, activated two chain forms, single chain protease domains and two chain protease domains.
 15 These polypeptides are encoded by nucleic acids that include sequences encoding a protease domain that exhibits proteolytic activity and that hybridizes to a nucleic acid molecule having a nucleotide sequence set forth in SEQ ID No. 5 or SEQ ID No. 22, typically under moderate, generally under high stringency, conditions and generally along the full-length or along at least about 70%, 80%
 20 or 90% of the full-length (or substantially the full-length) of the protease domain. Splice variants are also contemplated herein.

Structural features

The catalytic triad of the MTSP10 in SEQ ID No. 23 is H₅₀₃, D₅₅₁ and S₆₄₇. Disulfide bonds pairing in MTSP10 is as follows: C₄₈₈-C₅₀₄, C₅₈₇-C₆₅₃; C₆₁₉-C₆₃₂;
 25 C₆₄₃-C₆₇₃. Cys₅₇₃, which is in the protease domain, binds to Cys₅₉₆ outside the domain, and is unpaired in the single chain form of the protease domain.

MTSP10 includes several domains in addition to a catalytic domain residues 463-692 (SEQ ID No. 23; corresponding to residue 1-230 of SEQ ID No. 6) and a transmembrane domain. These include 2 CUB domains at aa104-
 30

-49-

217 and aa222-335, respectively; three LDLa domains at a aa340-377, aa381-412 and aa415-453 respectively.

Protease domains

MTSP protease domains include the single chain protease domains of
5 MTSP10. Provided are the protease domains or proteins that include a portion of an MTSP that is the protease domain of any MTSP, particularly a MTSP10. The protein can also include other non-MTSP sequences of amino acids, but includes the protease domain or a sufficient portion thereof to exhibit catalytic activity in any *in vitro* assay that assess such protease activity, such as any
10 provided herein. Also provided are two chain activated forms of the full length protease and also two chain forms of the protease domain. Thus, isolated, substantially pure proteases that include the protease domains or catalytically active portions thereof as single chain forms of SPs are provided. The protease domains can be included in a longer protein, and such longer protein is optionally
15 the activated MTSP10 protein, up to and including a full-length, or an MTSP10 zymogen.

In particular, exemplary protease domains include at least a sufficient portion of sequences of amino acids set forth of SEQ ID No. 6 or SEQ ID No. 23 (encoded by nucleotides in SEQ ID No. 5 or SEQ ID No. 22) to retain catalytic
20 activity *in vitro*.

As noted, the protease domains of an MTSP are single-chain polypeptides or two-chain polypeptides, with an N-terminus (such as IV, VV, IL and II) generated at the cleavage site (generally having the consensus sequence R↓VVG, R↓IVG, R↓IVQ, R↓IVNG, R↓ILG, R↓VGL, R↓ILG or a variation
25 thereof; an N-terminus R↓V or R↓I, where the arrow represents the cleavage point) when the zymogen is activated. The protease domain of an exemplary MTSP10, produced is produced by activation cleavage between an R462 and the I at residue 463 of SEQ ID No. 23 (R↓I) includes the sequence R↓IIGT, as set forth in SEQ ID No. 23. A single chain form includes residues 1-230 of SEQ ID
30 No. 6 or catalytically active fragments thereof. Hence any length polypeptide that includes the protease domain (residues 463-692 of SEQ ID No. 23) or catalytically active fragments thereof, is contemplated herein. Two chain forms

RECTIFIED SHEET (RULE 91)

-50-

also are provided and include at least a polypeptide from C₂₉₆ up to and including C₅₇₃ of SEQ ID No. 23 or corresponding residues of an MTSP10 that has at least 60%, 70%, 80%, 90%, or 95% sequence identity there with and/or is:

- a) a polypeptide that includes the sequence of amino acids set forth in SEQ ID No. 6, particularly amino acids 1-230;
- b) a polypeptide that includes the sequence of amino acids set forth in SEQ ID No. 23;
- c) a polypeptide encoded by a sequence of nucleotides that hybridizes under conditions of low, moderate or high stringency to the sequence of nucleotides set forth in SEQ ID No. 5 or to SEQ ID No. 22 or hybridizes along at least 70%, 80%, 90% or 95% of its full-length;
- d) a polypeptide that includes the sequence of amino acids set forth in SEQ ID No. 6 or SEQ ID No. 23 or a catalytically active portion thereof;
- e) a polypeptide that includes a sequence of amino acids having at least about 60%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% sequence identity with the sequence of amino acids set forth in SEQ ID No. 6 or SEQ ID No. 23; and/or
- f) a protease domain of a polypeptide encoded by a splice variant of a sequence of nucleotides that encodes an MTSP10 of any of a)-e).

Also provided are polypeptides that are encoded by nucleic acid molecules that meet criteria specified below as (a)-(e)

Muteins and derivatives

Full-length MTSP10, zymogen and activated forms thereof and MTSP10 protease domains, portions thereof, and muteins and derivatives of such polypeptides are provided. The domains, fragments, derivatives or analogs of an MTSP10 that are functionally active are capable of exhibiting one or more functional activities associated with the MTSP10 polypeptide, such as serine protease activity, immunogenicity and antigenicity, are provided.

Among the derivatives are those based on animal MTSP10s, including, but are not limited to, rodent, such as mouse and rat; fowl, such as chicken; ruminants, such as goats, cows, deer, sheep; ovine, such as pigs; and humans.

-51-

- For example, MTSP10 derivatives can be made by altering their sequences by substitutions, additions or deletions. MTSP10 derivatives include, but are not limited to, those containing, as a primary amino acid sequence, all or part of the amino acid sequence of MTSP10, including altered sequences in which
- 5 functionally equivalent amino acid residues are substituted for residues within the sequence resulting in a silent change. For example, one or more amino acid residues within the sequence can be substituted by another amino acid of a similar polarity which acts as a functional equivalent, resulting in a silent alteration. Substitutes for an amino acid within the sequence can be selected
- 10 from other members of the class to which the amino acid belongs. For example, the nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan and methionine. The polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine. The positively charged (basic) amino acids include arginine,
- 15 lysine and histidine. The negatively charged (acidic) amino acids include aspartic acid and glutamic acid (see, *e.g.*, Table 1). Muteins of the MTSP10 or a domain thereof, such as a protease domain, in which up to about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 85%, 90% or 95% of the amino acids are replaced with another amino acid are provided. Generally such muteins retain at
- 20 least about 1%, 2%, 3%, 5%, 7%, 8%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80% or 90% of the protease activity of the unmutated protein. Those of skill in the art recognize that a polypeptide that retains at least 1% of the activity of the wild-type protease is sufficiently active for use in screening assays or for other applications.
- 25 Muteins of the MTSP10 or a domain thereof, such as a protease domain, in which up to about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% of the amino acids are replaced with another amino acid are provided. Generally such muteins retain at least about 1%, 2%,
- 30 3%, 5%, 7%, 8%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80% or 90% (in increased activity, *i.e.*, 101, 102, 103, 104, 105, 110% or greater) of the protease activity of the unmutated protein.

RECTIFIED SHEET (RULE 91)

-52-

Included among the polypeptides provided herein are the MTSP10 protease domain or a polypeptide with amino acid changes such that the specificity and protease activity remains substantially unchanged or changed (increased or decreased) by a specified percentage, such as 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5%. In particular, a substantially purified mammalian MTSP polypeptide is provided that has a transmembrane (TM) domain and additionally includes two CUB domains, three LDL receptor type a domains and a serine protease catalytic domain is provided.

Also provided is a substantially purified protein containing a sequence of amino acids that has at least 60%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identity to the MTSP10 where the percentage identity is determined using standard algorithms and gap penalties that maximize the percentage identity. The human MTSP10 polypeptide is included, although other mammalian MTSP10 polypeptides are contemplated. The precise percentage of identity can be specified if needed.

Muteins in which one or more of the Cys residues, particularly, a residue that is paired in the activated two chain form, but unpaired in the protease domain alone is/are replaced with any amino acid, typically, although not necessarily, a conservative amino acid residue, such as Ser, are contemplated. Muteins of MTSP10, particularly those in which Cys residues, such as the Cys₅₇₃ in the single chain protease domain, is replaced with another amino acid, such as Ser, Gly or Ala, that does not eliminate the activity, are provided. Also provided are substantially purified MTSP10 polypeptides and functional domains thereof, including catalytically active domains and portions, that have at least about 60%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% sequence identity with a protease domain that includes a sequence of amino acids set forth in SEQ ID No. 6, particularly amino acids 1-230 or catalytically active fragments thereof.

RECTIFIED SHEET (RULE 91)

-53-

Muteins of the protein are also provided in which amino acids are replaced with other amino acids. Among the muteins are those in which the Cys residues, is/are replaced typically with a conservative amino acid residues, such as a serine. Such muteins are also provided herein. Muteins in which 10%,
5 20%, 30%, 35%, 40%, 45%, 50% or more of the amino acids are replaced but the resulting polypeptide retains at least about 1%, 2%, 3%, 5%, 7%, 8%, 10%, 20%, 30%, 35%, 40%, 45%, 50%, 60%, 70%, 80%, 90% or 95% of the catalytic activity as the unmodified form for the same substrate are provided.

Muteins can be made by making conservative amino acid substitutions
10 and also non-conservative amino acid substitutions. For example, amino acid substitutions that desirably alter properties of the proteins can be made. In one embodiment, mutations that prevent degradation of the polypeptide can be made. Many proteases cleave after basic residues, such as R and K; to eliminate such cleavage, the basic residue is replaced with a non-basic residue. Also, non-
15 conservative changes at amino acids outside of the protease domain can be effected without altering protease activity. Non-conservative changes at amino acids that are responsible for activities other than protease activity may be desirable. For example, interaction of the protease with an inhibitor can be blocked while retaining catalytic activity by effecting a non-conservative change
20 at the site interaction of the inhibitor with the protease. Similarly, receptor binding can be altered without altering catalytic activity by effecting a non-conservative or conservative change at a site of interaction of the receptor with the protease.

Antigenic epitopes that contain at least 4, 5, 6, 7, 8, 9, 10, 11, 12, 13,
25 14, 15, 20, 25, 30, 40, 50, and typically 10-15 amino acids of the MTSP10 polypeptide are provided. These antigenic epitopes are used, for example, to raise antibodies. Antibodies specific for each epitope or combinations thereof and for single and two-chain forms are also provided.

-54-

Nucleic acid molecules, vectors and plasmids, cells and expression of MTSP10 polypeptides

Nucleic acid molecules

Due to the degeneracy of nucleotide coding sequences, other nucleic sequences which encode substantially the same amino acid sequence as a MTSP are contemplated. These include but are not limited to nucleic acid molecules that include all or portions of MTSP10-encoding genes that are altered by the substitution of different codons that encode the amino acid residue within the sequence, thus producing a silent change.

Nucleic acids

Also provided herein are nucleic acid molecules that encode MTSP10 polypeptides and the encoded proteins. In particular, nucleic acid molecules encoding MTSP10 from animals, including splice variants thereof are provided. The encoded proteins are also provided. Also provided are functional domains thereof. For each of the nucleic acid molecules provided, the nucleic acid can be DNA or RNA or PNA or other nucleic acid analogs or can include non-natural nucleotide bases. Also provided are isolated nucleic acid molecules that include a sequence of nucleotides complementary to the nucleotide sequence encoding an MTSP.

Also provided are nucleic acid molecules that encode single chain or two chain MTSP proteases that have proteolytic activity in an *in vitro* proteolysis assay and that have at least 60%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% sequence identity with the full-length of a protease domain of an MTSP10 polypeptide, or that hybridize along their full-length or along at least about 70%, 80% or 90% of the full-length nucleic acid to a nucleic acids that encode a protease domain, particularly under conditions of moderate, generally high, stringency. As above, the encoded polypeptides contain the protease as a single chain; activated forms thereof can be produced and are provided.

In one embodiment, a nucleic acid molecule that encodes an MTSP, designated MTSP10 is provided. The nucleic acid molecule includes the open reading frame in the sequence of nucleotides set forth in SEQ ID No. 5 or SEQ ID

-55-

No. 22. Also provided are nucleic acid molecules that hybridize under conditions of at least low stringency, moderate stringency, and generally high stringency to the following sequence of nucleic acids (SEQ ID No. 5) particularly to the open reading frame encompassed by nucleotides that encode a single protease domain thereof, or any domain of MTSP10.

In certain embodiments, the isolated nucleic acid fragment hybridizes to the nucleic acid having the nucleotide sequence set forth in SEQ ID No. 5 under high stringency conditions, and generally contains the sequence of nucleotides set forth as nucleotides 1-690 in SEQ ID No. 5. The protein contains a transmembrane domain (TM) and a serine protease domain and can contain additional domains, including a CUB domain and LDLR domain.

Also provided, are muteins of the nucleic acid molecules that encode polypeptides in which amino acids are replaced with other amino acids. Among the muteins are those in which the Cys residue-encoding codons, is/are replaced with other amino acid residues, such as a codon encoding a serine. Such muteins are also provided herein. Each of such domains is provided herein as are nucleic acid molecules that include sequences of nucleotides encoding such domains. Some MTSPs can additionally include a LDLR domain, a scavenger-receptor cysteine rich (SRCR) domain and other domains.

The isolated nucleic acid fragment is DNA, including genomic or cDNA, or is RNA, or can include other components, such as peptide nucleic acid (PNA) and other nucleotide analogs. The isolated nucleic acid can include additional components, such as heterologous or native promoters, and other transcriptional and translational regulatory sequences, these genes can be linked to other genes, such as reporter genes or other indicator genes or genes that encode indicators.

Also provided are nucleic acid molecules that hybridize to the above-noted sequences of nucleotides encoding MTSP10 at least at low stringency, moderate stringency, and typically at high stringency, and that encode the protease domain and/or the full-length protein or at least 60%, 70%, 80% or 90% of the full-length protease domain or other domains of an MTSP10 or a splice variant or allelic variant thereof. Generally the molecules hybridize under

RECTIFIED SHEET (RULE 91)

-56-

such conditions along their full-length or along at least 70%, 80% or 90% of the full-length for at least one domain and encode at least one domain, such as the protease or extracellular domain, of the polypeptide. In particular, such nucleic acid molecules include any isolated nucleic acid fragment that encodes at least one domain of a membrane serine protease, that (1) contains a sequence of nucleotides that encodes the protease or a domain thereof, and (2) is selected from among:

- (a) a sequence of nucleotides that encodes the protease or a domain thereof that includes a sequence of nucleotides set forth in SEQ ID No. 5, particularly nucleotides 1-690, or SEQ ID No. 22;
- (b) a sequence of nucleotides that encodes such portion or the full-length protease and hybridizes under conditions of high stringency, generally to nucleic acid that is complementary to a mRNA transcript present in a mammalian cell that encodes such protein or fragment thereof;
- (c) a sequence of nucleotides that encodes the protease or a domain thereof that includes a sequence of nucleotides set having at least about 60%, 70%, 80%, 90% or 95% sequence identity with the sequence set forth in SEQ ID No. 5, particularly nucleotides 1-690, or in SEQ ID No. 22;
- (d) a sequence of nucleotides that encodes a transmembrane protease or domain thereof that includes a sequence of amino acids encoded by such portion or the full-length open reading frame;
- (e) a sequence of nucleotides that encodes the protease or a domain thereof that includes a sequence of nucleotides set having at least about 60%, 70%, 80%, 90% or 95% sequence identity with the sequence set forth in SEQ ID No. 5 or SEQ ID No. 22; and
- (f) a sequence of nucleotides that encodes the transmembrane protease that includes a sequence of amino acids encoded by a sequence of nucleotides that encodes such subunit and hybridizes under conditions of low, moderate or high stringency to DNA that is complementary to the mRNA transcript.

RECTIFIED SHEET (RULE 91)

-57-

The isolated nucleic acids can contain least 10 nucleotides, 25 nucleotides, 50 nucleotides, 100 nucleotides, 150 nucleotides, or 200 nucleotides or more contiguous nucleotides of an MTSP10-encoding sequence, or a full-length SP coding sequence. In another embodiment, the nucleic acids are smaller than 35, 200 or 500 nucleotides in length. Nucleic acids that hybridize to or are complementary to an MTSP10-encoding nucleic acid molecule can be single or double-stranded. For example, nucleic acids are provided that include a sequence complementary to (specifically are the inverse complement of) at least 10, 25, 50, 100, or 200 nucleotides or the entire coding region of an MTSP10 encoding nucleic acid, particularly the protease domain thereof. For MTSP10 the full-length protein or a domain or active fragment thereof is also provided.

Probes, primers, antisense oligonucleotides and dsRNA

Also provided are fragments thereof that can be used as probes or primers and that contain at least about 10 nucleotides, 14 nucleotides, generally at least about 16 nucleotides, often at least about 30 nucleotides. The length of the probe or primer is a function of the size of the genome probed; the larger the genome, the longer the probe or primer required for specific hybridization to a single site. Those of skill in the art can select appropriately sized probes and primers. Generally probes and primers as described are single-stranded. Double stranded probes and primers can be used, if they are denatured when used.

Probes and primers derived from the nucleic acid molecules are provided. Such probes and primers contain at least 8, 14, 16, 30, 100 or more contiguous nucleotides with identity to contiguous nucleotides of an MTSP10, and probes of at least 14, 16, 30, 50 or 100 contiguous sequence of nucleotides of SEQ ID No. 5 or SEQ ID No. 22. The probes and primers are optionally labelled with a detectable label, such as a radiolabel or a fluorescent tag, or can be mass differentiated for detection by mass spectrometry or other means.

-58-

Also provided is an isolated nucleic acid molecule that includes the sequence of molecules that is complementary to the nucleotide sequence encoding MTSP10 or the portion thereof. Double-stranded RNA (dsRNA), such as RNAi is also provided.

5 **Plasmids, Vectors and Cells**

Plasmids and vectors containing the nucleic acid molecules are also provided. Cells containing the vectors, including cells that express the encoded proteins are provided. The cell can be a bacterial cell, a yeast cell, a fungal cell, a plant cell, an insect cell or an animal cell. Methods for producing an MTSP or
10 single chain form of the protease domain thereof by, for example, growing the cell under conditions whereby the encoded MTSP is expressed by the cell, and recovering the expressed protein, are provided herein. As noted, for MTSP10, the full-length zymogens and activated proteins and activated (two chain) protease and single chain protease domains are provided. As described herein,
15 the cells are used for expression of the protein, which can be secreted or expressed in the cytoplasm.

As discussed below, the MTSP10 polypeptide, and catalytically active portions thereof, can be expressed on the surface of a cell. In addition, all or portions thereof can be expressed as a secreted protein using the native signal
20 sequence or a heterologous signal. Alternatively, all or portions of the polypeptide can be expressed as inclusion bodies in the cytoplasm and isolated therefrom. The resulting protein can be treated to refold if necessary.

The above discussion provides an overview and some details of the exemplified MTSP10s.

25 **C. Tumor specificity and tissue expression profiles**

MTSPs are of interest because they appear to be expressed and/or activated at different levels in tumor cells from normal cells, or have functional activity that is different in tumor cells from normal cells, such as by an alteration in a substrate for the MTSP, or a cofactor or receptor of the MTSP. MTSP10 is
30 of interest because it is expressed or is active in tumor cells. Hence the MTSPs provided herein can serve as diagnostic markers for certain tumors.

RECTIFIED SHEET (RULE 91)

-59-

Each MTSP has a characteristic tissue expression profile; the MTSPs in particular, although not exclusively expressed or activated in tumors, exhibit characteristic tumor tissue expression or activation profiles. In some instances, MTSPs can have different activity in a tumor cell from a non-tumor cell by virtue of a change in a substrate or cofactor or receptor therefor or other factor that would alter the functional activity of the MTSP. Hence each can serve as a diagnostic marker for particular tumors, by virtue of a level of activity and/or expression or function in a subject (i.e. a mammal, particularly a human) with neoplastic disease, compared to a subject or subjects that do not have the neoplastic disease. In addition, detection of activity (and/or expression) in a particular tissue can be indicative of neoplastic disease. Shed MTSPs in body fluids can be indicative of neoplastic disease. Also, by virtue of the activity and/or expression profiles of each, they can serve as therapeutic targets, such as by administration of modulators of the activity thereof, or, as by administration of a prodrug specifically activated by one of the MTSPs.

Tissue expression profiles

MTSP10

MTSP10 transcript was detected in pancreas, lung and kidney. MTSP10 transcript was also detected in small intestine Marathon-Ready cDNA (Clontech). The MTSP10 transcript was detected in breast carcinoma (GI-101), lung carcinoma (LX-1 and GI-117), ovarian carcinoma (GI-102), and pancreatic adenocarcinoma (GI-103). The MTSP10 transcript was weakly detected in prostatic adenocarcinoma (PC3). The MTSP10 transcript was also detected in the CWR22R prostate tumor grown on nude mice. No apparent signal was detected in two forms of colon adenocarcinomas (GI-112 and CX-1).

D. Identification and isolation of MTSP10 polypeptide genes

The MTSP polypeptides and/or domains thereof, can be obtained by methods well known in the art for protein purification and recombinant protein expression. Any method known to those of skill in the art for identification of nucleic acids that encode desired genes can be used. Any method available in the art can be used to obtain a full-length (i.e., encompassing the entire coding region) cDNA or genomic DNA clone encoding an MTSP polypeptide. For

-60-

example, the polymerase chain reaction (PCR) can be used to amplify a sequence that is expressed in normal and tumor cells or tissues, *e.g.*, nucleic acids encoding an MTSP10 polypeptide (SEQ. Nos: 5 and 17), in a genomic or cDNA library. Oligonucleotide primers that hybridize to sequences at the 3' and 5' termini of the identified sequences can be used as primers to amplify by PCR sequences from a nucleic acid sample (RNA or DNA), generally a cDNA library, from an appropriate source (*e.g.*, tumor or cancer tissue).

PCR can be carried out, *e.g.*, by use of a Perkin-Elmer Cetus thermal cycler and Taq polymerase (Gene Amp[®]). The DNA being amplified can include mRNA or cDNA or genomic DNA from any eukaryotic species. One can choose to synthesize several different degenerate primers, for use in the PCR reactions. It is also possible to vary the stringency of hybridization conditions used in priming the PCR reactions, to amplify nucleic acid homologs (*e.g.*, to obtain *MTSP polypeptide* sequences from species other than humans or to obtain human sequences with homology to MTSP10 polypeptide) by allowing for greater or lesser degrees of nucleotide sequence similarity between the known nucleotide sequence and the nucleic acid homolog being isolated. For cross-species hybridization, low stringency to moderate stringency conditions are used. For same species hybridization, moderately stringent to highly stringent conditions are used. The conditions can be empirically determined.

After successful amplification of the nucleic acid containing all or a portion of the identified MTSP polypeptide sequence or of a nucleic acid encoding all or a portion of an MTSP polypeptide homolog, that segment can be molecularly cloned and sequenced, and used as a probe to isolate a complete cDNA or genomic clone. This, in turn, permits the determination of the gene's complete nucleotide sequence, the analysis of its expression, and the production of its protein product for functional analysis. Once the nucleotide sequence is determined, an open reading frame encoding the MTSP polypeptide gene protein product can be determined by any method well known in the art for determining open reading frames, for example, using publicly available computer programs for nucleotide sequence analysis. Once an open reading frame is defined, it is routine to determine the amino acid sequence of the protein encoded by the

-61-

open reading frame. In this way, the nucleotide sequences of the entire MTSP polypeptide genes as well as the amino acid sequences of MTSP polypeptide proteins and analogs can be identified.

Any eukaryotic cell potentially can serve as the nucleic acid source for the molecular cloning of the MTSP polypeptide gene. The nucleic acids can be isolated from vertebrate, mammalian, human, porcine, bovine, feline, avian, equine, canine, as well as additional primate sources, insects, plants and other organisms. The DNA can be obtained by standard procedures known in the art from cloned DNA (*e.g.*, a DNA "library"), by chemical synthesis, by cDNA cloning, or by the cloning of genomic DNA, or fragments thereof, purified from the desired cell (see, *e.g.*, Sambrook *et al.* (2001) *Molecular Cloning, A Laboratory Manual*, 3d Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York; Glover, D.M. (ed.), 1985, *DNA Cloning: A Practical Approach*, MRL Press, Ltd., Oxford, U.K. Vol. I, II). Clones derived from genomic DNA can contain regulatory and intron DNA regions in addition to coding regions; clones derived from cDNA will contain only exon sequences. For any source, the gene is cloned into a suitable vector for propagation thereof.

In the molecular cloning of the gene from genomic DNA, DNA fragments are generated, some of which will encode the desired gene.

The DNA can be cleaved at specific sites using various restriction enzymes. Alternatively, one can use DNase in the presence of manganese to fragment the DNA, or the DNA can be physically sheared, for example, by sonication. The linear DNA fragments then can be separated according to size by standard techniques, including but not limited to, agarose and polyacrylamide gel electrophoresis and column chromatography.

Once the DNA fragments are generated, identification of the specific DNA fragment containing the desired gene can be accomplished in a number of ways. For example, a portion of the MTSP polypeptide (of any species) gene (*e.g.*, a PCR amplification product obtained as described above or an oligonucleotide having a sequence of a portion of the known nucleotide sequence) or its specific RNA, or a fragment thereof can be purified and labeled, and the generated DNA fragments can be screened by nucleic acid hybridization to the labeled probe

RECTIFIED SHEET (RULE 91)

-62-

(Benton and Davis, *Science* 196:180 (1977); Grunstein and Hogness, *Proc. Natl. Acad. Sci. U.S.A.* 72:3961 (1975)). Those DNA fragments with substantial homology to the probe will hybridize. It is also possible to identify the appropriate fragment by restriction enzyme digestion(s) and comparison of
5 fragment sizes with those expected according to a known restriction map if such is available or by DNA sequence analysis and comparison to the known nucleotide sequence of MTSP polypeptide. Further selection can be carried out on the basis of the properties of the gene. Alternatively, the presence of the gene can be detected by assays based on the physical, chemical, or
10 immunological properties of its expressed product. For example, cDNA clones, or DNA clones which hybrid-select the proper mRNA, can be selected which produce a protein that, *e.g.*, has similar or identical electrophoretic migration, isoelectric focusing behavior, proteolytic digestion maps, antigenic properties, serine protease activity. If an anti-MTSP polypeptide antibody is available, the
15 protein can be identified by binding of labeled antibody to the putatively MTSP polypeptide synthesizing clones, in an ELISA (enzyme-linked immunosorbent assay)-type procedure.

Alternatives to isolating the MTSP10 polypeptide genomic DNA include, but are not limited to, chemically synthesizing the gene sequence from a known
20 sequence or making cDNA to the mRNA that encodes the MTSP polypeptide. For example, RNA for cDNA cloning of the MTSP polypeptide gene can be isolated from cells expressing the protein. The identified and isolated nucleic acids then can be inserted into an appropriate cloning vector. A large number of vector-host systems known in the art can be used. Possible vectors include, but
25 are not limited to, plasmids or modified viruses, but the vector system must be compatible with the host cell used. Such vectors include, but are not limited to, bacteriophages such as lambda derivatives, or plasmids such as pBR322 or pUC plasmid derivatives or the Bluescript vector (Stratagene, La Jolla, CA). The insertion into a cloning vector can, for example, be accomplished by ligating the
30 DNA fragment into a cloning vector which has complementary cohesive termini. If the complementary restriction sites used to fragment the DNA are not present in the cloning vector, the ends of the DNA molecules can be enzymatically

RECTIFIED SHEET (RULE 91)

-63-

modified. Alternatively, any site desired can be produced by ligating nucleotide sequences (linkers) onto the DNA termini; these ligated linkers can include specific chemically synthesized oligonucleotides encoding restriction endonuclease recognition sequences. In an alternative method, the cleaved
5 vector and MTSP polypeptide gene can be modified by homopolymeric tailing. Recombinant molecules can be introduced into host cells via transformation, transfection, infection, electroporation, calcium precipitation and other methods, so that many copies of the gene sequence are generated.

In specific embodiments, transformation of host cells with recombinant
10 DNA molecules that incorporate the isolated MTSP polypeptide gene, cDNA, or synthesized DNA sequence enables generation of multiple copies of the gene. Thus, the gene can be obtained in large quantities by growing transformants, isolating the recombinant DNA molecules from the transformants and, when necessary, retrieving the inserted gene from the isolated recombinant DNA.

15 **E. Vectors, plasmids and cells that contain nucleic acids encoding an MTSP polypeptide or protease domain thereof and expression of MTSP polypeptides**

Vectors and cells

For recombinant expression of one or more of the MTSP polypeptides, the
20 nucleic acid containing all or a portion of the nucleotide sequence encoding the MTSP polypeptide can be inserted into an appropriate expression vector, *i.e.*, a vector that contains the necessary elements for the transcription and translation of the inserted protein coding sequence. The necessary transcriptional and translational signals can also be supplied by the native promoter for MTSP
25 genes, and/or their flanking regions.

Also provided are vectors that contain nucleic acid encoding the MTSPs. Cells containing the vectors are also provided. The cells include eukaryotic and prokaryotic cells, and the vectors are any suitable for use therein.

Prokaryotic and eukaryotic cells, including endothelial cells, containing the
30 vectors are provided. Such cells include bacterial cells, yeast cells, fungal cells, plant cells, insect cells and animal cells. The cells are used to produce an MTSP polypeptide or protease domain thereof by (a) growing the above-described cells under conditions whereby the encoded MTSP polypeptide or protease domain of

-64-

the MTSP polypeptide is expressed by the cell, and then (b) recovering the expressed protease domain protein. In the exemplified embodiments, the protease domain is secreted into the medium.

In one embodiment, vectors that include a sequence of nucleotides that
5 encode a polypeptide that has protease activity and contains all or a portion of only the protease domain, or multiple copies thereof, of an SP protein are provided. Also provided are vectors that include a sequence of nucleotides that encodes the protease domain and additional portions of an SP protein up to and including a full length SP protein, as well as multiple copies thereof. The vectors
10 can be selected for expression of the SP protein or protease domain thereof in the cell or such that the SP protein is expressed as a secreted protein. Alternatively, the vectors can include signals necessary for secretion of encoded proteins. When the protease domain is expressed the nucleic acid is linked to nucleic acid encoding a secretion signal, such as the *Saccharomyces cerevisiae* α
15 mating factor signal sequence or a portion thereof, or the native signal sequence.

A variety of host-vector systems can be used to express the protein coding sequence. These include but are not limited to mammalian cell systems infected with virus (*e.g.* vaccinia virus, adenovirus, etc.); insect cell systems infected with virus (*e.g.* baculovirus); microorganisms such as yeast containing
20 yeast vectors; or bacteria transformed with bacteriophage, DNA, plasmid DNA, or cosmid DNA. The expression elements of vectors vary in their strengths and specificities. Depending on the host-vector system used, any one of a number of suitable transcription and translation elements can be used.

Any methods known to those of skill in the art for the insertion of nucleic
25 acid fragments into a vector can be used to construct expression vectors containing a chimeric gene containing appropriate transcriptional/translational control signals and protein coding sequences. These methods can include *in vitro* recombinant DNA and synthetic techniques and *in vivo* recombinants (genetic recombination). Expression of nucleic acid sequences encoding MTSP
30 polypeptide, or domains, derivatives, fragments or homologs thereof, can be regulated by a second nucleic acid sequence so that the genes or fragments thereof are expressed in a host transformed with the recombinant DNA

molecule(s). For example, expression of the proteins can be controlled by any promoter/enhancer known in the art. In a specific embodiment, the promoter is not native to the genes for MTSP polypeptide. Promoters which can be used include but are not limited to the SV40 early promoter (Bernoist and Chambon, *Nature* 290:304-310 (1981)), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto et al., *Cell* 22:787-797 (1980)), the herpes thymidine kinase promoter (Wagner et al., *Proc. Natl. Acad. Sci. USA* 78:1441-1445 (1981)), the regulatory sequences of the metallothionein gene (Brinster et al., *Nature* 296:39-42 (1982)); prokaryotic expression vectors such as the β -lactamase promoter (Villa-Komaroff et al., *Proc. Natl. Acad. Sci. USA* 75:3727-3731 (1978)) or the *tac* promoter (DeBoer et al., *Proc. Natl. Acad. Sci. USA* 80:21-25 (1983)); see also "Useful Proteins from Recombinant Bacteria": in *Scientific American* 242:79-94 (1980)); plant expression vectors containing the nopaline synthetase promoter (Herrar-Estrella et al., *Nature* 303:209-213 (1984)) or the cauliflower mosaic virus 35S RNA promoter (Gardner et al., *Nucleic Acids Res.* 9:2871 (1981)), and the promoter of the photosynthetic enzyme ribulose biphosphate carboxylase (Herrera-Estrella et al., *Nature* 310:115-120 (1984)); promoter elements from yeast and other fungi such as the Gal4 promoter, the alcohol dehydrogenase promoter, the phosphoglycerol kinase promoter, the alkaline phosphatase promoter, and the following animal transcriptional control regions that exhibit tissue specificity and have been used in transgenic animals: elastase I gene control region which is active in pancreatic acinar cells (Swift et al., *Cell* 38:639-646 (1984); Ornitz et al., *Cold Spring Harbor Symp. Quant. Biol.* 50:399-409 (1986); MacDonald, *Hepatology* 7:425-515 (1987)); insulin gene control region which is active in pancreatic beta cells (Hanahan et al., *Nature* 315:115-122 (1985)), immunoglobulin gene control region which is active in lymphoid cells (Grosschedl et al., *Cell* 38:647-658 (1984); Adams et al., *Nature* 318:533-538 (1985); Alexander et al., *Mol. Cell Biol.* 7:1436-1444 (1987)), mouse mammary tumor virus control region which is active in testicular, breast, lymphoid and mast cells (Leder et al., *Cell* 45:485-495 (1986)), albumin gene control region which is active in liver (Pinckert et al., *Genes and Devel.* 1:268-276 (1987)), alpha-fetoprotein gene control region which is active in liver

-66-

- (Krumlauf et al., *Mol. Cell. Biol.* 5:1639-1648 (1985); Hammer et al., *Science* 235:53-58 (1987)), alpha-1 antitrypsin gene control region which is active in liver (Kelsey et al., *Genes and Devel.* 1:161-171 (1987)), beta globin gene control region which is active in myeloid cells (Mogam et al., *Nature* 315:338-340 (1985); Kollias et al., *Cell* 46:89-94 (1986)), myelin basic protein gene control region which is active in oligodendrocyte cells of the brain (Readhead et al., *Cell* 48:703-712 (1987)), myosin light chain-2 gene control region which is active in skeletal muscle (Sani, *Nature* 314:283-286 (1985)), and gonadotrophic releasing hormone gene control region which is active in gonadotrophs of the
- 10 hypothalamus (Mason et al., *Science* 234:1372-1378 (1986)).

- In a specific embodiment, a vector is used that contains a promoter operably linked to nucleic acids encoding an MTSP polypeptide, or a domain, fragment, derivative or homolog, thereof, one or more origins of replication, and optionally, one or more selectable markers (*e.g.*, an antibiotic resistance gene).
- 15 Expression vectors containing the coding sequences, or portions thereof, of an MTSP polypeptide, is made, for example, by subcloning the coding portions into the EcoRI restriction site of each of the three pGEX vectors (glutathione S-transferase expression vectors (Smith and Johnson, *Gene* 7:31-40 (1988)). This allows for the expression of products in the correct reading frame. Exemplary
- 20 vectors and systems for expression of the protease domains of the MTSP polypeptides include the well-known *Pichia* vectors (available, for example, from Invitrogen, San Diego, CA), particularly those designed for secretion of the encoded proteins. The protein can also be expressed cytoplasmically, such as in the inclusion bodies. One exemplary vector is described in the EXAMPLES.
- 25 Plasmids for transformation of *E. coli* cells, include, for example, the pET expression vectors (see, U.S patent 4,952,496; available from NOVAGEN, Madison, WI; see, also literature published by Novagen describing the system). Such plasmids include pET 11a, which contains the T7lac promoter, T7 terminator, the inducible *E. coli* lac operator, and the lac repressor gene; pET
- 30 12a-c, which contains the T7 promoter, T7 terminator, and the *E. coli* ompT secretion signal; and pET 15b and pET19b (NOVAGEN, Madison, WI), which contain a His-TagTM leader sequence for use in purification with a His column and

-67-

a thrombin cleavage site that permits cleavage following purification over the column; the T7-lac promoter region and the T7 terminator.

The vectors are introduced into host cells, such as *Pichia* cells and bacterial cells, such as *E. coli*, and the proteins expressed therein. Exemplary
5 *Pichia* strains, include, for example, GS115. Exemplary bacterial hosts contain chromosomal copies of DNA encoding T7 RNA polymerase operably linked to an inducible promoter, such as the lacUV promoter (see, U.S. Patent No. 4,952,496). Such hosts include, but are not limited to, the lysogenic *E. coli* strain BL21(DE3).

10 **Expression and production of proteins**

The MTSP domains, derivatives and analogs can be produced by various methods known in the art. For example, once a recombinant cell expressing an MTSP polypeptide, or a domain, fragment or derivative thereof, is identified, the individual gene product can be isolated and analyzed. This is achieved by
15 assays based on the physical and/or functional properties of the protein, including, but not limited to, radioactive labeling of the product followed by analysis by gel electrophoresis, immunoassay, cross-linking to marker-labeled product, and assays of proteolytic activity.

The MTSP polypeptides can be isolated and purified by standard methods
20 known in the art (either from natural sources or recombinant host cells expressing the complexes or proteins), including but not restricted to column chromatography (*e.g.*, ion exchange, affinity, gel exclusion, reversed-phase high pressure and fast protein liquid), differential centrifugation, differential solubility, or by any other standard technique used for the purification of proteins.
25 Functional properties can be evaluated using any suitable assay known in the art.

Alternatively, once an MTSP polypeptide or its domain or derivative is identified, the amino acid sequence of the protein can be deduced from the nucleotide sequence of the gene which encodes it. As a result, the protein or its
30 domain or derivative can be synthesized by standard chemical methods known in the art (*e.g.* see Hunkapiller et al, *Nature* 310:105-111 (1984)).

-68-

Manipulations of MTSP polypeptide sequences can be made at the protein level. Also contemplated herein are MTSP polypeptide proteins, domains thereof, derivatives or analogs or fragments thereof, which are differentially modified during or after translation, *e.g.*, by glycosylation, acetylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to an antibody molecule or other cellular ligand. Any of numerous chemical modifications can be carried out by known techniques, including but not limited to specific chemical cleavage by cyanogen bromide, trypsin, chymotrypsin, papain, V8 protease, NaBH₄, acetylation, formylation, oxidation, reduction, metabolic synthesis in the presence of tunicamycin and other such agents.

In addition, domains, analogs and derivatives of an MTSP polypeptide can be chemically synthesized. For example, a peptide corresponding to a portion of an MTSP polypeptide, which includes the desired domain or which mediates the desired activity *in vitro* can be synthesized by use of a peptide synthesizer. Furthermore, if desired, nonclassical amino acids or chemical amino acid analogs can be introduced as a substitution or addition into the MTSP polypeptide sequence. Non-classical amino acids include but are not limited to the D-isomers of the common amino acids, α -amino isobutyric acid, 4-aminobutyric acid, Abu, 2-aminobutyric acid, ϵ -Abu, ϵ -Ahx, 6-amino hexanoic acid, Aib, 2-amino isobutyric acid, 3-amino propionic acid, ornithine, norleucine, norvaline, hydroxyproline, sarcosine, citrulline, cysteic acid, t-butylglycine, t-butylalanine, phenylglycine, cyclohexylalanine, β -alanine, fluoro-amino acids, designer amino acids such as β -methyl amino acids, Ca-methyl amino acids, Na-methyl amino acids, and amino acid analogs in general. Furthermore, the amino acid can be D (dextrorotary) or L (levorotary).

In cases where natural products are suspected of being mutant or are isolated from new species, the amino acid sequence of the MTSP polypeptide isolated from the natural source, as well as those expressed *in vitro*, or from synthesized expression vectors *in vivo* or *in vitro*, can be determined from analysis of the DNA sequence, or alternatively, by direct sequencing of the

-69-

isolated protein. Such analysis can be performed by manual sequencing or through use of an automated amino acid sequenator.

Modifications

- A variety of modifications of the MTSP polypeptides and domains are contemplated herein. An MTSP-encoding nucleic acid molecule can be modified by any of numerous strategies known in the art (Sambrook *et al.* (1990), *Molecular Cloning, A Laboratory Manual*, 2d ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, New York). The sequences can be cleaved at appropriate sites with restriction endonuclease(s), followed by further enzymatic modification if desired, isolated, and ligated *in vitro*. In the production of the gene encoding a domain, derivative or analog of MTSP, care should be taken to ensure that the modified gene retains the original translational reading frame, uninterrupted by translational stop signals, in the gene region where the desired activity is encoded.
- Additionally, the MTSP-encoding nucleic acid molecules can be mutated *in vitro* or *in vivo*, to create and/or destroy translation, initiation, and/or termination sequences, or to create variations in coding regions and/or form new restriction endonuclease sites or destroy pre-existing ones, to facilitate further *in vitro* modification. Also, as described herein muteins with primary sequence alterations, such as replacements of Cys residues and elimination or addition of glycosylation sites are contemplated; the MTSP10 that includes the sequence of amino acids set forth in SEQ ID No. 6 has potential glycosylation sites at N₆₇, N₂₇₂, N₃₃₆, N₃₈₃, N₄₀₉ and N₄₁₈ (SEQ ID No. 23). Mutations can be effected by any technique for mutagenesis known in the art, including, but not limited to, chemical mutagenesis and *in vitro* site-directed mutagenesis (Hutchinson et al., *J. Biol. Chem.* 253:6551-6558 (1978)), use of TAB[®] linkers (Pharmacia). In one embodiment, for example, an MTSP polypeptide or domain thereof is modified to include a fluorescent label. In other specific embodiments, the MTSP polypeptide is modified such that heterobifunctional reagents can be used to crosslink the members of a complex.

In addition, domains, analogs and derivatives of an MTSP can be chemically synthesized. For example, a peptide corresponding to a portion of an

-70-

MTSP, which includes the desired domain or which mediates the desired activity *in vitro* can be synthesized by use of a peptide synthesizer. Furthermore, if desired, nonclassical amino acids or chemical amino acid analogs can be introduced as a substitution or addition into the MTSP sequence. Non-classical amino acids include but are not limited to the D-isomers of the common amino acids, α -amino isobutyric acid, 4-aminobutyric acid, Abu, 2-aminobutyric acid, ϵ -Abu, e-Ahx, 6-amino hexanoic acid, Aib, 2-amino isobutyric acid, 3-amino propionic acid, ornithine, norleucine, norvaline, hydroxyproline, sarcosine, citrulline, cysteic acid, t-butylglycine, t-butylalanine, phenylglycine, cyclohexylalanine, β -alanine, fluoro-amino acids, designer amino acids such as β -methyl amino acids, Ca-methyl amino acids, Na-methyl amino acids, and amino acid analogs in general. Furthermore, the amino acid can be D (dextrorotary) or L (levorotary).

F. SCREENING METHODS

The single chain protease domains, as shown herein, can be used in a variety of methods to identify compounds that modulate the activity thereof. For SPs that exhibit higher activity or expression in tumor cells, compounds that inhibit the proteolytic activity are of particular interest. For any SPs that are active at lower levels in tumor cells, compounds or agents that enhance the activity are potentially of interest. In all instances the identified compounds include agents that are candidate cancer treatments.

Several types of assays are exemplified and described herein. It is understood that the protease domains can be used in other assays. It is shown here, however, that the single chain protease domains exhibit catalytic activity. As such they are ideal for *in vitro* screening assays. They can also be used in binding assays.

The MTSP10 full length zymogens, activated enzymes, single and two chain protease domains are contemplated for use in any screening assay known to those of skill in the art, including those provided herein. Hence the following description, if directed to proteolytic assays is intended to apply to use of a single chain protease domain or a catalytically active portion thereof of any SP, including an MTSP10. Other assays, such as binding assays are provided

-71-

herein, particularly for use with an MTSP10, including any variants, such as splice variants thereof.

1. Catalytic Assays for identification of agents that modulate the protease activity of an SP protein

5 Methods for identifying a modulator of the catalytic activity of an SP, particularly a single chain protease domain or catalytically active portion thereof, are provided herein. The methods can be practiced by: contacting the MTSP10, a full-length zymogen or activated form, and particularly a single-chain domain thereof, with a substrate of the MTSP10 in the presence of a test substance,
10 and detecting the proteolysis of the substrate, whereby the activity of the MTSP10 is assessed, and comparing the activity to a control. For example, a control can be the activity of the MTSP10 assessed by contacting an MTSP10, including a full-length zymogen or activated form, and particularly a single-chain domain thereof, particularly a single-chain domain thereof, with a substrate of
15 the MTSP10, and detecting the proteolysis of the substrate, whereby the activity of the MTSP10 is assessed. The results in the presence and absence of the test compounds are compared. A difference in the activity indicates that the test substance modulates the activity of the MTSP10. Activators of MTSP10 activation cleavage are also contemplated; such assays are discussed below.

20 In one embodiment a plurality of the test substances are screened simultaneously in the above screening method. In another embodiment, the MTSP10 is isolated from a target cell as a means for then identifying agents that are potentially specific for the target cell.

 In another embodiment, a test substance is a therapeutic compound, and
25 whereby a difference of the MTSP10 activity measured in the presence and in the absence of the test substance indicates that the target cell responds to the therapeutic compound.

 One method includes the steps of (a) contacting the MTSP10 polypeptide or protease domain thereof with one or a plurality of test compounds under
30 conditions conducive to interaction between the ligand and the compounds; and (b) identifying one or more compounds in the plurality that specifically binds to the ligand.

-72-

Another method provided herein includes the steps of a) contacting an MTSP10 polypeptide or protease domain thereof with a substrate of the MTSP10 polypeptide, and detecting the proteolysis of the substrate, whereby the activity of the MTSP10 polypeptide is assessed; b) contacting the MTSP10 polypeptide with a substrate of the MTSP10 polypeptide in the presence of a test substance, and detecting the proteolysis of the substrate, whereby the activity of the MTSP10 polypeptide is assessed; and c) comparing the activity of the MTSP10 polypeptide assessed in steps a) and b), whereby the activity measured in step a) differs from the activity measured in step b) indicates that the test substance modulates the activity of the MTSP10 polypeptide.

In another embodiment, a plurality of the test substances are screened simultaneously. In comparing the activity of an MTSP10 polypeptide in the presence and absence of a test substance to assess whether the test substance is a modulator of the MTSP10 polypeptide, it is unnecessary to assay the activity in parallel, although such parallel measurement is typical. It is possible to measure the activity of the MTSP10 polypeptide at one time point and compare the measured activity to a historical value of the activity of the MTSP10 polypeptide.

For instance, one can measure the activity of the MTSP10 polypeptide in the presence of a test substance and compare with historical value of the activity of the MTSP10 polypeptide measured previously in the absence of the test substance, and *vice versa*. This can be accomplished, for example, by providing the activity of the MTSP10 polypeptide on an insert or pamphlet provided with a kit for conducting the assay.

Methods for selecting substrates for a particular SP are described in the EXAMPLES, and particular proteolytic assays are exemplified.

Combinations and kits containing the combinations optionally including instructions for performing the assays are provided. The combinations include an MTSP10 polypeptide and a substrate of the MTSP10 polypeptide to be assayed; and, optionally reagents for detecting proteolysis of the substrate. The substrates, which can be chromogenic or fluorogenic molecules, including proteins, subject to proteolysis by a particular MTSP10 polypeptide, can be

RECTIFIED SHEET (RULE 91)

-73-

identified empirically by testing the ability of the MTSP10 polypeptide to cleave the test substrate. Substrates that are cleaved most effectively (*i.e.*, at the lowest concentrations and/or fastest rate or under desirable conditions), are identified.

- 5 Additionally provided herein is a kit containing the above-described combination. The kit optionally includes instructions for identifying a modulator of the activity of an MTSP10 polypeptide. Any MTSP10 polypeptide is contemplated as target for identifying modulators of the activity thereof.

2. Binding assays

- 10 Also provided herein are methods for identification and isolation of agents, particularly compounds that bind to MTSP10s. The assays are designed to identify agents that bind to the zymogen form, the single chain isolated protease domain (or a protein, other than an MTSP10 polypeptide, that contains the protease domain of an MTSP10 polypeptide), and to the activated form,
- 15 including the activated form derived from the full length zymogen or from an extended protease domain. The identified compounds are candidates or leads for identification of compounds for treatments of tumors and other disorders and diseases involving aberrant angiogenesis. The MTSP10 polypeptides used in the methods include any MTSP10 polypeptide as defined herein, including the
- 20 MTSP10 single chain protease domain or proteolytically active portion thereof.

- A variety of methods are provided herein. These methods can be performed in solution or in solid phase reactions in which the MTSP10 polypeptide(s) or protease domain(s) thereof are linked, either directly or indirectly via a linker, to a solid support. Screening assays are described in the
- 25 Examples, and these assays have been used to identify candidate compounds. For purposes herein, all binding assays described above are provided for MTSP10.

- Methods for identifying an agent, such as a compound, that specifically binds to an MTSP10 single and/or two chain protease domain, a zymogen and/or
- 30 full-length activated MTSP10 or two chain protease domain thereof are provided herein. The method can be practiced by (a) contacting the MTSP10 with one or a plurality of test agents under conditions conducive to binding between the

-74-

MTSP10 and an agent; and (b) identifying one or more agents within the plurality that specifically binds to the MTSP10.

For example, in practicing such methods the MTSP10 polypeptide is mixed with a potential binding partner or an extract or fraction of a cell under

- 5 conditions that allow the association of potential binding partners with the polypeptide. After mixing, peptides, polypeptides, proteins or other molecules that have become associated with an MTSP10 are separated from the mixture. The binding partner that bound to the MTSP10 can then be removed and further analyzed. To identify and isolate a binding partner, the entire protein, for
10 instance the entire polypeptide of SEQ ID No. 6 can be used. Alternatively, a fragment of the protein can be used.

- A variety of methods can be used to obtain cell extracts or body fluids, such as blood, serum, urine, sweat, synovial fluid, CSF and other such fluids. For example, cells can be disrupted using either physical or chemical disruption
15 methods. Examples of physical disruption methods include, but are not limited to, sonication and mechanical shearing. Examples of chemical lysis methods include, but are not limited to, detergent lysis and enzyme lysis. A skilled artisan can readily adapt methods for preparing cellular extracts in order to obtain extracts for use in the present methods.

- 20 Once an extract of a cell is prepared, the extract is mixed with the MTSP10 under conditions in which association of the protein with the binding partner can occur. A variety of conditions can be used, including conditions that resemble conditions found in the cytoplasm of a human cell or in a body fluid, such as blood. Features, such as osmolarity, pH, temperature, and the
25 concentration of cellular extract used, can be varied to optimize the association of the protein with the binding partner. Similarly, methods for isolation of molecules of interest from body fluids are known.

- After mixing under appropriate conditions, the bound complex is separated from the mixture. A variety of techniques can be used to separate the
30 mixture. For example, antibodies specific to an MTSP10 can be used to immunoprecipitate the binding partner complex. Alternatively, standard chemical

RECTIFIED SHEET (RULE 91)

-75-

separation techniques such as chromatography and density/sediment centrifugation can be used.

After removing the non-associated cellular constituents in the extract, the binding partner can be dissociated from the complex using conventional
5 methods. For example, dissociation can be accomplished by altering the salt concentration or pH of the mixture.

To aid in separating associated binding partner pairs from the mixed extract, the MTSP10 can be immobilized on a solid support. For example, the protein can be attached to a nitrocellulose matrix or acrylic beads. Attachment
10 of the protein or a fragment thereof to a solid support aids in separating peptide/binding partner pairs from other constituents found in the extract. The identified binding partners can be either a single protein or a complex made up of two or more proteins.

Alternatively, the nucleic acid molecules encoding the single chain
15 proteases can be used in a yeast two-hybrid system. The yeast two-hybrid system has been used to identify other protein partner pairs and can readily be adapted to employ the nucleic acid molecules herein described.

Another *in vitro* binding assay, particularly for an MTSP10, uses a mixture of a polypeptide that contains at least the catalytic domain of one of
20 these proteins and one or more candidate binding targets or substrates. After incubating the mixture under appropriate conditions, the ability of the MTSP10 or a polypeptide fragment thereof containing the catalytic domain to bind to or interact with the candidate substrate is assessed. For cell-free binding assays, one of the components includes or is coupled to a detectable label. The label
25 can provide for direct detection, such as radioactivity, luminescence, optical or electron density, *etc.*, or indirect detection such as an epitope tag, an enzyme, *etc.* A variety of methods can be employed to detect the label depending on the nature of the label and other assay components. For example, the label can be detected bound to the solid substrate or a portion of the bound complex
30 containing the label can be separated from the solid substrate, and the label thereafter detected.

3. Detection of signal transduction

MTSP10, which is a transmembrane protein, can be involved directly or indirectly in signal transduction directly as a cell surface receptor or indirectly by activating proteins, such as pro-growth factors that can initiate signal

5 transduction.

In addition, secretion of MTSP10, such as the extracellular domain of MTSP10, can be involved in signal transduction either directly by binding to or interacting with a cell surface receptor or indirectly by activating proteins, such as pro-growth factors that can initiate signal transduction. Assays for assessing

10 signal transduction are well known to those of skill in the art, and can be adapted for use with the MTSP10 polypeptide.

Assays for identifying agents that affect or alter signal transduction mediated directly or indirectly, such as via activation of a pro-growth factor, by an MTSP10, particularly the full length or a sufficient portion to anchor the

15 extracellular domain or a functional portion thereof of an MTSP10 on the surface of a cell are provided. Such assays, include, for example, transcription based assays in which modulation of a transduced signal is assessed by detecting an effect on an expression from a reporter gene (see, *e.g.*, U.S. Patent No. 5,436,128).

20 4. Methods for Identifying Agents that Modulate the Expression a Nucleic Acid Encoding an MTSP10

Another embodiment provides methods for identifying agents that modulate the expression of a nucleic acid encoding an MTSP10. Such assays use any available means of monitoring for changes in the expression level of the

25 nucleic acids encoding an MTSP10.

In one assay format, cell lines that contain reporter gene fusions between the open reading frame of MTSP10 or a domain thereof, particularly the protease domain and any assayable fusion partner can be prepared. Numerous assayable fusion partners are known and readily available including the firefly luciferase

30 gene and the gene encoding chloramphenicol acetyltransferase (Alam *et al.*, *Anal. Biochem.* 188: 245-54 (1990)). Cell lines containing the reporter gene fusions are then exposed to the agent to be tested under appropriate conditions

-77-

and time. Differential expression of the reporter gene between samples exposed to the agent and control samples identifies agents which modulate the expression of a nucleic acid encoding an MTSP10.

Additional assay formats can be used to monitor the ability of the agent
5 to modulate the expression of a nucleic acid encoding an MTSP10. For instance, mRNA expression can be monitored directly by hybridization to the nucleic acids. Cell lines are exposed to the agent to be tested under appropriate conditions and time and total RNA or mRNA is isolated by standard procedures (see, *e.g.*, Sambrook *et al.* (1989) MOLECULAR CLONING: A LABORATORY
10 MANUAL, 2nd Ed. Cold Spring Harbor Laboratory Press). Probes to detect differences in RNA expression levels between cells exposed to the agent and control cells can be prepared from the nucleic acids. It is typical, but not necessary, to design probes which hybridize only with target nucleic acids under conditions of high stringency. Only highly complementary nucleic acid hybrids
15 form under conditions of high stringency. Accordingly, the stringency of the assay conditions determines the amount of complementarity which should exist between two nucleic acid strands in order to form a hybrid. Stringency should be chosen to maximize the difference in stability between the probe:target hybrid and potential probe:non-target hybrids.

20 Probes can be designed from the nucleic acids through methods known in the art. For instance, the G+C content of the probe and the probe length can affect probe binding to its target sequence. Methods to optimize probe specificity are commonly available (see, *e.g.*, Sambrook *et al.* (1989) MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed. Cold Spring
25 Harbor Laboratory Press); and Ausubel *et al.* (1995) CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, Greene Publishing Co., NY).

Hybridization conditions are modified using known methods (see, *e.g.*, Sambrook *et al.* (1989) MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed. Cold Spring Harbor Laboratory Press); and Ausubel *et al.* (1995)
30 CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, Greene Publishing Co., NY), as required for each probe. Hybridization of total cellular RNA or RNA enriched for polyA RNA can be accomplished in any available format. For instance, total

-78-

cellular RNA or RNA enriched for polyA RNA can be affixed to a solid support, and the solid support exposed to at least one probe comprising at least one, or part of one of the nucleic acid molecules under conditions in which the probe specifically hybridizes. Alternatively, nucleic acid fragments comprising at least one, or part of one of the sequences can be affixed to a solid support, such as a porous glass wafer. The glass wafer can then be exposed to total cellular RNA or polyA RNA from a sample under conditions in which the affixed sequences specifically hybridize. Such glass wafers and hybridization methods are widely available, for example, those disclosed by Beattie (WO 95/11755). By examining for the ability of a given probe to specifically hybridize to an RNA sample from an untreated cell population and from a cell population exposed to the agent, agents which up or down regulate the expression of a nucleic acid encoding the MTSP10 polypeptide, are identified.

In one format, the relative amounts of a protein between a cell population that has been exposed to the agent to be tested compared to an un-exposed control cell population can be assayed (*e.g.*, a prostate cancer cell line, a lung cancer cell line, a colon cancer cell line or a breast cancer cell line). In this format, probes, such as specific antibodies, are used to monitor the differential expression or level of activity of the protein in the different cell populations or body fluids. Cell lines or populations or body fluids are exposed to the agent to be tested under appropriate conditions and time. Cellular lysates or body fluids can be prepared from the exposed cell line or population and a control, unexposed cell line or population or unexposed body fluid. The cellular lysates or body fluids are then analyzed with the probe.

For example, N- and C- terminal fragments of the MTSP10 can be expressed in bacteria and used to search for proteins which bind to these fragments. Fusion proteins, such as His-tag or GST fusion to the N- or C-terminal regions of the MTSP10 can be prepared for use as a substrate. These fusion proteins can be coupled to, for example, Glutathione-Sepharose beads and then probed with cell lysates or body fluids. Prior to lysis, the cells or body fluids can be treated with a candidate agent which can modulate an MTSP10 or proteins that interact with domains thereon. Lysate proteins binding to the

-79-

fusion proteins can be resolved by SDS-PAGE, isolated and identified by protein sequencing or mass spectroscopy, as is known in the art.

Antibody probes are prepared by immunizing suitable mammalian hosts in appropriate immunization protocols using the peptides, polypeptides or proteins if they are of sufficient length (*e.g.*, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 20, 25, 30, 35, 40 or more consecutive amino acids the MTSP10 polypeptide or if required to enhance immunogenicity, conjugated to suitable carriers. Methods for preparing immunogenic conjugates with carriers, such as bovine serum albumin (BSA), keyhole limpet hemocyanin (KLH), or other carrier proteins are well known in the art. In some circumstances, direct conjugation using, for example, carbodiimide reagents can be effective; in other instances linking reagents such as those supplied by Pierce Chemical Co., Rockford, IL, can be desirable to provide accessibility to the hapten. Hapten peptides can be extended at either the amino or carboxy terminus with a Cys residue or interspersed with cysteine residues, for example, to facilitate linking to a carrier. Administration of the immunogens is conducted generally by injection over a suitable time period and with use of suitable adjuvants, as is generally understood in the art. During the immunization schedule, titers of antibodies are taken to determine adequacy of antibody formation.

Anti-peptide antibodies can be generated using synthetic peptides corresponding to, for example, the carboxy terminal amino acids of the MTSP10. Synthetic peptides can be as small as 1-3 amino acids in length, generally at least 4 or more amino acid residues long. The peptides can be coupled to KLH using standard methods and can be immunized into animals, such as rabbits or ungulates. Polyclonal antibodies can then be purified, for example using Actigel beads containing the covalently bound peptide.

While the polyclonal antisera produced in this way can be satisfactory for some applications, for pharmaceutical compositions, use of monoclonal preparations are generally used. Immortalized cell lines which secrete the desired monoclonal antibodies can be prepared using the standard method of Kohler *et al.*, (*Nature* 256: 495-7 (1975)) or modifications which effect immortalization of lymphocytes or spleen cells, as is generally known. The

-80-

immortalized cell lines secreting the desired antibodies are screened by immunoassay in which the antigen is the peptide hapten, polypeptide or protein. When the appropriate immortalized cell culture secreting the desired antibody is identified, the cells can be cultured either *in vitro* or by production *in vivo* via
5 ascites fluid. Of particular interest, are monoclonal antibodies that recognize the catalytic domain or activation cleavage site (region) of an MTSP10.

Additionally, the zymogen or two-chain form of the MTSP10 can be used to make monoclonal antibodies that recognize conformation epitopes. The desired monoclonal antibodies are then recovered from the culture supernatant
10 or from the ascites supernatant. Fragments of the monoclonals or the polyclonal antisera which contain the immunologically significant portion can be used as antagonists, as well as the intact antibodies. Immunologically reactive fragments, such as the Fab, Fab', or F(ab')₂ fragments are often used, especially in a therapeutic context, as these fragments are generally less immunogenic than
15 the whole immunoglobulin.

The antibodies or fragments can also be produced. Regions that bind specifically to the desired regions of receptor also can be produced in the context of chimeras with multiple species origin.

Agents that are assayed in the above method can be randomly selected
20 or rationally selected or designed.

The agents can be, as examples, peptides, small molecules, and carbohydrates. A skilled artisan can readily recognize that there is no limit as to the structural nature of the agents.

The peptide agents can be prepared using standard solid phase (or
25 solution phase) peptide synthesis methods, as is known in the art. In addition, the DNA encoding these peptides can be synthesized using commercially available oligonucleotide synthesis instrumentation and produced recombinantly using standard recombinant production systems. The production using solid phase peptide synthesis is necessitated if non-gene-encoded amino acids are to
30 be included.

RECTIFIED SHEET (RULE 91)

-81-

G. Assay formats and selection of test substances that modulate at least one activity of an MTSP10 polypeptide

Methods for identifying agents that modulate at least one activity of an MTSP10 are provided. The methods include phage display and other methods
5 for assessing alterations in the activity of an MTSP10. Such methods or assays can use any means of monitoring or detecting the desired activity. A variety of formats and detection protocols are known for performing screening assays. Any such formats and protocols can be adapted for identifying modulators of MTSP10 polypeptide activities. The following includes a discussion of exemplary
10 protocols.

1. High throughput screening assays

Although the above-described assay can be conducted where a single MTSP10 polypeptide is screened, and/or a single test substance is screened in one assay, the assay typically is conducted in a high throughput screening
15 mode, *i.e.*, a plurality of the SP proteins are screened against and/or a plurality of the test substances are screened simultaneously (*See generally, High Throughput Screening: The Discovery of Bioactive Substances* (Devlin, Ed.) Marcel Dekker, 1997; Sittampalam et al., *Curr. Opin. Chem. Biol.*, 1:384-91 (1997); and Silverman et al., *Curr. Opin. Chem. Biol.*, 2:397-403 (1998)). For
20 example, the assay can be conducted in a multi-well (*e.g.*, 24-, 48-, 96-, 384-, 1536-well or higher density), chip or array format.

High-throughput screening (HTS) is the process of testing a large number of diverse chemical structures against disease targets to identify "hits" (Sittampalam et al., *Curr. Opin. Chem. Biol.*, 1:384-91 (1997)). Current state-of-
25 the-art HTS operations are highly automated and computerized to handle sample preparation, assay procedures and the subsequent processing of large volumes of data.

Detection technologies employed in high-throughput screens depend on the type of biochemical pathway being investigated (Sittampalam et al., *Curr. Opin. Chem. Biol.*, 1:384-91 (1997)). These methods include, radiochemical
30 methods, such as the scintillation proximity assays (SPA), which can be adapted to a variety of enzyme assays (Lerner et al., *J. Biomol. Screening*, 1:135-143

(1996); Baker et al., *Anal. Biochem.*, 239:20-24 (1996); Baum et al., *Anal. Biochem.*, 237:129-134 (1996); and Sullivan et al., *J. Biomol. Screening* 2:19-23 (1997)) and protein-protein interaction assays (Braunwalder et al., *J. Biomol. Screening* 1:23-26 (1996); Sonatore et al., *Anal. Biochem.* 240:289-297 (1996); and Chen et al., *J. Biol. Chem.* 271:25308-25315 (1996)), and non-isotopic detection methods, including but are not limited to, colorimetric and luminescence detection methods, resonance energy transfer (RET) methods, time-resolved fluorescence (HTRF) methods, cell-based fluorescence assays, such as fluorescence resonance energy transfer (FRET) procedures (see, e.g., Gonzalez et al., *Biophys. J.*, 69:1272-1280 (1995)), fluorescence polarization or anisotropy methods (see, e.g., Jameson et al., *Methods Enzymol.* 246:283-300 (1995); Jolley, *J. Biomol. Screening* 1:33-38 (1996); Lynch et al., *Anal. Biochem.* 247:77-82 (1997)), fluorescence correlation spectroscopy (FCS) and other such methods.

2. Test Substances

Test compounds, including small molecules, antibodies, proteins, nucleic acids, peptides, natural products, extracts containing natural products and libraries and collections thereof, can be screened in the above-described assays and assays described below to identify compounds that modulate the activity of an MTSP10 polypeptide. Rational drug design methodologies that rely on computational chemistry can be used to screen and identify candidate compounds.

The compounds identified by the screening methods include inhibitors, including antagonists, and can be agonists. Compounds for screening include any compounds and collections of compounds available, known or that can be prepared.

a. Selection of Compounds

Compounds can be selected for their potency and selectivity of inhibition of serine proteases, especially an MTSP10 polypeptide. As described herein, and as generally known, a target serine protease and its substrate are combined under assay conditions permitting reaction of the protease with its substrate. The assay is performed in the absence of test compound, and in the presence of

-83-

increasing concentrations of the test compound. The concentration of test compound at which 50% of the serine protease activity is inhibited by the test compound is the IC_{50} value (Inhibitory Concentration) or EC_{50} (Effective Concentration) value for that compound. Within a series or group of test compounds, those having lower IC_{50} or EC_{50} values are considered more potent inhibitors of the serine protease than those compounds having higher IC_{50} or EC_{50} values. The IC_{50} measurement is often used for more simplistic assays, whereas the EC_{50} is often used for more complicated assays, such as those employing cells.

Typically candidate compounds have an IC_{50} value of 100 nM or less as measured in an *in vitro* assay for inhibition of MTSP10 polypeptide activity. The test compounds also are evaluated for selectivity toward a serine protease. As described herein, and as generally known, a test compound is assayed for its potency toward a panel of serine proteases and other enzymes and an IC_{50} value or EC_{50} value is determined for each test compound in each assay system. A compound that demonstrates a low IC_{50} value or EC_{50} value for the target enzyme, *e.g.*, MTSP10 polypeptide, and a higher IC_{50} value or EC_{50} value for other enzymes within the test panel (*e.g.*, urokinase tissue plasminogen activator, thrombin, Factor Xa), is considered to be selective toward the target enzyme. Generally, a compound is deemed selective if its IC_{50} value or EC_{50} value in the target enzyme assay is at least one order of magnitude less than the next smallest IC_{50} value or EC_{50} value measured in the selectivity panel of enzymes.

Compounds are also evaluated for their activity *in vivo*. The type of assay chosen for evaluation of test compounds depends on the pathological condition to be treated or prevented by use of the compound, as well as the route of administration to be evaluated for the test compound.

For instance, to evaluate the activity of a compound to reduce tumor growth through inhibition of MTSP10 polypeptide, the procedures described by Jankun et al., *Canc. Res.* 57:559-563 (1997) to evaluate PAI-1 can be employed. Briefly, the ATCC cell lines DU145 and LnCaP are injected into SCID mice. After tumors are established, the mice are given test compound according

-84-

to a dosing regime determined from the compound's *in vitro* characteristics. The Jankun *et al.* compound was administered in water. Tumor volume measurements are taken twice a week for about five weeks. A compound is deemed active if an animal to which the compound was administered exhibited
5 decreased tumor volume, as compared to animals receiving appropriate control compounds.

Another *in vivo* experimental model designed to evaluate the effect of p-aminobenzamidine, a swine protease inhibitor, on reducing tumor volume is described by Billström *et al.*, *Int. J. Cancer* 61:542-547 (1995).

10 To evaluate the ability of a compound to reduce the occurrence of, or inhibit, metastasis, the procedures described by Kobayashi *et al.* *Int. J. Canc.* 57:727-733d (1994) can be employed. Briefly, a murine xenograft selected for high lung colonization potential is injected into C57B1/6 mice i.v. (experimental metastasis) or s.c. into the abdominal wall (spontaneous metastasis). Various
15 concentrations of the compound to be tested can be admixed with the tumor cells in Matrigel prior to injection. Daily i.p. injections of the test compound are made either on days 1-6 or days 7-13 after tumor inoculation. The animals are sacrificed about three or four weeks after tumor inoculation, and the lung tumor colonies are counted. Evaluation of the resulting data permits a determination as
20 to efficacy of the test compound, optimal dosing and route of administration.

The activity of the tested compounds toward decreasing tumor volume and metastasis can be evaluated in model described in Rabbani *et al.*, *Int. J. Cancer* 63:840-845 (1995) to evaluate their inhibitor. There, Mat LyLu tumor cells were injected into the flank of Copenhagen rats. The animals were
25 implanted with osmotic minipumps to continuously administer various doses of test compound for up to three weeks. The tumor mass and volume of experimental and control animals were evaluated during the experiment, as were metastatic growths. Evaluation of the resulting data permits a determination as to efficacy of the test compound, optimal dosing, and route of administration.
30 Some of these authors described a related protocol in Xing *et al.*, *Canc. Res.* 57:3585-3593 (1997).

To evaluate the anti-angiogenesis activity of a compound, a rabbit cornea neovascularization model can be employed (see, *e.g.*, Avery *et al.* (1990) *Arch. Ophthalmol.*, 108:1474-147). Avery *et al.* describes anesthetizing New Zealand albino rabbits and then making a central corneal incision and forming a radial
5 corneal pocket. A slow release prostaglandin pellet was placed in the pocket to induce neovascularization. Test compound was administered i.p. for five days, at which time the animals were sacrificed. The effect of the test compound is evaluated by review of periodic photographs taken of the limbus, which can be used to calculate the area of neovascular response and, therefore, limbal
10 neovascularization. A decreased area of neovascularization as compared with appropriate controls indicates the test compound was effective at decreasing or inhibiting neovascularization.

An angiogenesis model used to evaluate the effect of a test compound in preventing angiogenesis is described by Min *et al.* *Canc. Res.* 56:2428-2433
15 (1996). C57BL6 mice receive subcutaneous injections of a Matrigel mixture containing bFGF, as the angiogenesis-inducing agent, with and without the test compound. After five days, the animals are sacrificed and the Matrigel plugs, in which neovascularization can be visualized, are photographed. An experimental animal receiving Matrigel and an effective dose of test compound exhibits less
20 vascularization than a control animal or an experimental animal receiving a less- or non-effective dose of compound.

An *in vivo* system designed to test compounds for their ability to limit the spread of primary tumors is described by Crowley *et al.*, *Proc. Natl. Acad. Sci.* 90:5021-5025 (1993). Nude mice are injected with tumor cells (PC3)
25 engineered to express CAT (chloramphenicol acetyltransferase). Compounds to be tested for their ability to decrease tumor size and/or metastases are administered to the animals, and subsequent measurements of tumor size and/or metastatic growths are made. In addition, the level of CAT detected in various organs provides an indication of the ability of the test compound to inhibit
30 metastasis; detection of less CAT in tissues of a treated animal versus a control animal indicates less CAT-expressing cells migrated to that tissue.

In vivo experimental models designed to evaluate the inhibitory potential of a test serine protease inhibitors, using a tumor cell line F3II known to be highly invasive (see, e.g., Alonso *et al.* (1996) *Breast Canc. Res. Treat.* 40:209-223) are provided. Alonso describes *in vivo* studies for toxicity determination, tumor growth, invasiveness, spontaneous metastasis, experimental lung metastasis, and an angiogenesis assay.

The CAM model (chick embryo chorioallantoic membrane model; Ossowski (1988) *J. Cell Biol.* 107:2437-2445), provides another method for evaluating the inhibitory activity of a test compound. In the CAM model, tumor cells invade through the chorioallantoic membrane containing CAM (with tumor cells in the presence of several serine protease inhibitors results in less or no invasion of the tumor cells through the membrane). Thus, the CAM assay is performed with CAM and tumor cells in the presence and absence of various concentrations of test compound. The invasiveness of tumor cells is measured under such conditions to provide an indication of the compound's inhibitory activity. A compound having inhibitory activity correlates with less tumor invasion.

The CAM model is also used in a standard assay of angiogenesis (*i.e.*, effect on formation of new blood vessels (Brooks *et al. Methods in Molecular Biology* 129:257-269 (1999)). According to this model, a filter disc containing an angiogenesis inducer, such as basic fibroblast growth factor (bFGF) is placed onto the CAM. Diffusion of the cytokine into the CAM induces local angiogenesis, which can be measured in several ways such as by counting the number of blood vessel branch points within the CAM directly below the filter disc. The ability of identified compounds to inhibit cytokine-induced angiogenesis can be tested using this model. A test compound can either be added to the filter disc that contains the angiogenesis inducer, be placed directly on the membrane or be administered systemically. The extent of new blood vessel formation in the presence and/or absence of test compound can be compared using this model. The formation of fewer new blood vessels in the presence of a test compound would be indicative of anti-angiogenesis activity.

-87-

Demonstration of anti-angiogenesis activity for inhibitors of an MTSP10 polypeptide indicates a role in angiogenesis for that SP protein.

b. Known serine protease inhibitors

Compounds for screening can be serine protease inhibitors, which can be
5 tested for their ability to inhibit the activity of an MTSP10.
Exemplary, serine protease inhibitors for use in the screening assays, include, but are not limited to: Serine Protease Inhibitor 3 (SPI-3) (Chen, *et al. Citokine*, 11:856-862 (1999)); Aprotinin (Iijima, R., *et al., J. Biochem. (Tokyo)* 126:912-916 (1999)); Kazal-type serine protease inhibitor-like proteins (Niimi, *et al. Eur.*
10 *J. Biochem.*, 266:282-292 (1999)); Kunitz-type serine protease inhibitor (Ravichandran, S., *et al., Acta Crystallogr. D. Biol. Crystallogr.*, 55:1814-1821 (1999)); Tissue factor pathway inhibitor-2/Matrix-associated serine protease inhibitor (TFPI-2/MSPI), (Liu, Y. *et al. Arch. Biochem. Biophys.* 370:112-8 (1999)); Bukunin (Cui, C.Y. *et al. J. Invest. Dermatol.* 113:182-8 (1999));
15 Nafmostat mesilate (Ryo, R. *et al. Vox Sang.* 76:241-6 (1999)); TPCK (Huang *et al. Oncogene* 18:3431-3439 (1999)); A synthetic cotton-bound serine protease inhibitor (Edwards (1999) *et al. Wound Repair Regen.* 7:106-18); FUT-175 (Sawada (1999) *et al. Stroke* 30:644-50); Combination of serine protease inhibitor FUT-0175 and thromboxane synthetase inhibitor OKY-046 (Kaminogo *et*
20 *al.* (1998) *Neurol. Med. Chir. (Tokyo)* 38:704-8; discussion 708-9); the rat serine protease inhibitor 2.1 gene (LeCam, A., *et al., Biochem. Biophys. Res. Commun.*, 253:311-4 (1998)); A new intracellular serine protease inhibitor expressed in the rat pituitary gland complexes with granzyme B (Hill *et al. FEBS Lett.* 440:361-4 (1998)); 3,4-Dichloroisocoumarin (Hammed *et al. Proc. Soc.*
25 *Exp. Biol. Med.*, 219:132-7 (1998)); LEX032 (Bains *et al. Eur. J. Pharmacol.* 356:67-72 (1998)); N-tosyl-L-phenylalanine chloromethyl ketone (Dryjanski *et al. Biochemistry* 37:14151-6 (1998)); Mouse gene for the serine protease inhibitor neuroserpin (P112) (Berger *et al. Gene*, 214:25-33 (1998)); Rat serine protease inhibitor 2.3 gene (Paul *et al. Eur. J. Biochem.* 254:538-46 (1998)); Ecotin
30 (Yang *et al. J. Mol. Biol.* 279:945-57 (1998)); A 14 kDa plant-related serine protease inhibitor (Roch *et al. Dev. Comp. Immunol.* 22(1):1-12 (1998)); Matrix-associated serine protease inhibitor TFPI-2/33 kDa MSPI (Rao *et al. Int. J.*

RECTIFIED SHEET (RULE 91)

- Cancer* 76:749-56 (1998)); ONO-3403 (Hiwasa *et al. Cancer Lett.* 126:221-5 (1998)); Bdelastasin (Moser *et al. Eur. J. Biochem.* 253:212-20 (1998)); Bikunin (Xu *et al. J. Mol. Biol.* 276:955-66 (1998)); Nafamostat mesilate (Mellgren *et al. Thromb. Haemost.* 79:342-7 (1998)); The growth hormone dependent serine
- 5 protease inhibitor, Spi 2.1 (Maake *et al. Endocrinology* 138:5630-6 (1997)); Growth factor activator inhibitor type 2, a Kunitz-type serine protease inhibitor (Kawaguchi *et al. J. Biol. Chem.*, 272:27558-64 (1997)); Heat-stable serine protease inhibitor protein from ovaries of the desert locust, *Schistocerca gregaria* (Hamdaoui *et al. Biochem. Biophys. Res. Commun.* 238:357-60 (1997)); Human
- 10 placental Hepatocyte growth factor activator inhibitor, a Kunitz-type serine protease inhibitor (Shimomura *et al. J. Biol. Chem.* 272:6370-6 (1997)); FUT-187, oral serine protease inhibitor (Shiozaki *et al. Gan To Kagaku Ryoho*, 23(14):1971-9 (1996)); Extracellular matrix-associated serine protease inhibitors (Mr 33,000, 31,000, and 27,000 (Rao, C.N., *et al., Arch. Biochem. Biophys.*,
- 15 335:82-92 (1996)); An irreversible isocoumarin serine protease inhibitor (Palencia, D.D., *et al., Biol. Reprod.*, 55:536-42 (1996)); 4-(2-aminoethyl)-benzenesulfonyl fluoride (AEBSF) (Nakabo *et al. J. Leukoc. Biol.* 60:328-36 (1996)); Neuroserpin (Osterwalder, T., *et al., EMBO J.* 15:2944-53 (1996)); Human serine protease inhibitor alpha-1-antitrypsin (Forney *et al. J. Parasitol.*
- 20 82:496-502 (1996)); Rat serine protease inhibitor 2.3 (Simar-Blanchet, A.E., *et al., Eur. J. Biochem.*, 236:638-48 (1996)); Gebaxate mesilate (parodi, F., *et al., J. Cardiothorac. Vasc. Anesth.* 10:235-7 (1996)); Recombinant serine protease inhibitor, CPTI II (Stankiewicz, M., *et al., Acta Biochim. Pol.*, 43(3):525-9 (1996)); A cysteine-rich serine protease inhibitor (Guamerin II) (Kim, D.R., *et al., J. Enzym. Inhib.*, 10:81-91 (1996)); Diisopropylfluorophosphate (Lundqvist, H., *et al., Inflamm. Res.*, 44(12):510-7 (1995)); Nexin 1 (Yu, D.W., *et al., J. Cell Sci.*, 108(Pt 12):3867-74 (1995)); LEX032 (Scalia, R., *et al., Shock*, 4(4):251-6 (1995)); Protease nexin I (Houenou, L.J., *et al., Proc. Natl. Acad. Sci. U.S.A.*, 92(3):895-9 (1995)); Chymase-directed serine protease inhibitor (Woodard S.L.,
- 30 *et al., J. Immunol.*, 153(11):5016-25 (1994)); N-alpha-tosyl-L-lysyl-chloromethyl ketone (TLCK) (Bourinbaier, A.S., *et al., Cell Immunol.*, 155(1):230-6 (1994)); Smp156 (Ghendler, Y., *et al., Exp. Parasitol.*, 78(2):121-31 (1994)); *Schistosoma*

- haematobium serine protease (Blanton, R.E., et al., *Mol. Biochem. Parasitol.*, 63(1):1-11 (1994)); Spi-1 (Warren, W.C., et al., *Mol. Cell Endocrinol.*, 98(1):27-32 (1993)); TAME (Jessop, J.J., et al., *Inflammation*, 17(5):613-31 (1993)); Antithrombin III (Kalaria, R.N., et al., *Am. J. Pathol.*, 143(3):886-93 (1993));
- 5 FOY-305 (Ohkoshi, M., et al., *Anticancer Res.*, 13(4):963-6 (1993)); Camostat mesilate (Senda, S., et al., *Intern. Med.*, 32(4):350-4 (1993)); Pigment epithelium-derived factor (Steele, F.R., et al., *Proc. Natl. Acad. Sci. U.S.A.*, 90(4):1526-30 (1993)); Antistasin (Holstein, T.W., et al., *FEBS Lett.*, 309(3):288-92 (1992)); The vaccinia virus K2L gene encodes a serine protease
- 10 inhibitor (Zhou, J., et al., *Virology*, 189(2):678-86 (1992)); Bowman-Birk serine-protease inhibitor (Werner, M.H., et al., *J. Mol. Biol.*, 225(3):873-89 (1992); FUT-175 (Yanamoto, H., et al., *Neurosurgery*, 30(3):358-63 (1992)); FUT-175; (Yanamoto, H., et al., *Neurosurgery*, 30(3):351-6, discussion 356-7 (1992)); PAI-I (Yreadwell, B.V., et al., *J. Orthop. Res.*, 9(3):309-16 (1991)); 3,4-
- 15 Dichloroisocoumarin (Rusbridge, N.M., et al., *FEBS Lett.*, 268(1):133-6 (1990)); Alpha 1-antichymotrypsin (Lindmark, B.E., et al., *Am. Rev. Respir. Des.*, 141(4 Pt 1):884-8 (1990)); P-toluenesulfonyl-L-arginine methyl ester (TAME) (Scuderi, P., *J. Immunol.*, 143(1):168-73 (1989)); Alpha 1-antichymotrypsin (Abraham, C.R., et al., *Cell*, 52(4):487-501 (1988)); Contrapsin (Modha, J., et al.,
- 20 *Parasitology*, 96 (Pt 1):99-109 (1988)); Alpha 2-antiplasmin (Holmes, W.E., et al., *J. Biol. Chem.*, 262(4):1659-64 (1987)); 3,4-dichloroisocoumarin (Harper, J.W., et al., *Biochemistry*, 24(8):1831-41 (1985)); Diisopropylfluorophosphate (Tsutsui, K., et al., *Biochem. Biophys. Res. Commun.*, 123(1):271-7 (1984)); Gabexate mesilate (Hesse, B., et al., *Pharmacol. Res. Commun.*, 16(7):637-45
- 25 (1984)); Phenyl methyl sulfonyl fluoride (Dufer, J., et al., *Scand. J. Haematol.*, 32(1):25-32 (1984)); Protease inhibitor CI-2 (McPhalen, C.A., et al., *J. Mol. Biol.*, 168(2):445-7 (1983)); Phenylmethylsulfonyl fluoride (Sekar V., et al., *Biochem. Biophys. Res. Commun.*, 89(2):474-8 (1979)); PGE1 (Feinstein, M.D., et al., *Prostaglandine*, 14(6):1075-93 (1977)).

-90-

c. Combinatorial libraries and other libraries

The source of compounds for the screening assays, can be libraries, including, but are not limited to, combinatorial libraries. Methods for synthesizing combinatorial libraries and characteristics of such combinatorial
5 libraries are known in the art (*See generally, Combinatorial Libraries: Synthesis, Screening and Application Potential* (Cortese Ed.) Walter de Gruyter, Inc., 1995; Tietze and Lieb, *Curr. Opin. Chem. Biol.*, 2(3):363-71 (1998); Lam, *Anticancer Drug Des.*, 12(3):145-67 (1997); Blaney and Martin, *Curr. Opin. Chem. Biol.*, 1(1):54-9 (1997); and Schultz and Schultz, *Biotechnol. Prog.*, 12(6):729-43
10 (1996)).

Methods and strategies for generating diverse libraries, primarily peptide- and nucleotide-based oligomer libraries, have been developed using molecular biology methods and/or simultaneous chemical synthesis methodologies (*see, e.g.,* Dower et al., *Annu. Rep. Med. Chem.*, 26:271-280 (1991); Fodor et al.,
15 *Science*, 251:767-773 (1991); Jung et al., *Angew. Chem. Int. Ed. Engl.*, 31:367-383 (1992); Zuckerman et al., *Proc. Natl. Acad. Sci. USA*, 89:4505-4509 (1992); Scott et al., *Science*, 249:386-390 (1990); Devlin et al., *Science*, 249:404-406 (1990); Cwirla et al., *Proc. Natl. Acad. Sci. USA*, 87:6378-6382 (1990); and Gallop et al., *J. Medicinal Chemistry*, 37:1233-1251 (1994)). The
20 resulting combinatorial libraries potentially contain millions of compounds that can be screened to identify compounds that exhibit a selected activity.

The libraries fall into roughly three categories: fusion-protein-displayed peptide libraries in which random peptides or proteins are presented on the surface of phage particles or proteins expressed from plasmids; support-bound
25 synthetic chemical libraries in which individual compounds or mixtures of compounds are presented on insoluble matrices, such as resin beads (*see, e.g.,* Lam et al., *Nature*, 354:82-84 (1991)) and cotton supports (*see, e.g.,* Eichler et al., *Biochemistry* 32:11035-11041 (1993)); and methods in which the compounds are used in solution (*see, e.g.,* Houghten et al., *Nature*, 354:84-86
30 (1991); Houghten et al., *BioTechniques*, 31:412-421 (1992); and Scott et al., *Curr. Opin. Biotechnol.*, 5:40-48 (1994)). There are numerous examples of synthetic peptide and oligonucleotide combinatorial libraries and there are many

RECTIFIED SHEET (RULE 91)

-91-

methods for producing libraries that contain non-peptidic small organic molecules. Such libraries can be based on a basis set of monomers that are combined to form mixtures of diverse organic molecules or that can be combined to form a library based upon a selected pharmacophore monomer.

- 5 Either a random or a deterministic combinatorial library can be screened by the presently disclosed and/or claimed screening methods. In either of these two libraries, each unit of the library is isolated and/or immobilized on a solid support. In the deterministic library, one knows *a priori* a particular unit's location on each solid support. In a random library, the location of a particular
- 10 unit is not known *a priori* although each site still contains a single unique unit. Many methods for preparing libraries are known to those of skill in this art (see, *e.g.*, Geysen et al., *Proc. Natl. Acad. Sci. USA*, 81:3998-4002 (1984), Houghten et al., *Proc. Natl. Acad. Sci. USA*, 81:5131-5135 (1985)). Combinatorial library generated by the any techniques known to those of skill in
- 15 the art are contemplated (see, *e.g.*, Table 1 of Schultz and Schultz, *Biotechnol. Prog.*, 12(6):729-43 (1996)) for screening; Bartel et al., *Science*, 261:1411-1418 (1993); Baumbach et al. *BioPharm*, (Can):24-35 (1992); Bock et al. *Nature*, 355:564-566 (1992); Borman, S., Combinatorial chemists focus on small molecules molecular recognition, and automation, *Chem. Eng. News*,
- 20 2(12):29 (1996); Boublik, et al., Eukaryotic Virus Display: Engineering the Major Surface Glycoproteins of the Autographa California Nuclear Polyhedrosis Virus (ACNPV) for the Presentation of Foreign Proteins on the Virus Surface, *Bio/Technology*, 13:1079-1084 (1995); Brenner, et al., Encoded Combinatorial Chemistry, *Proc. Natl. Acad. Sci. U.S.A.*, 89:5381-5383 (1992); Caflisch, et al.,
- 25 Computational Combinatorial Chemistry for *De Novo* Ligand Design: Review and Assessment, *Perspect. Drug Discovery Des.*, 3:51-84 (1995); Cheng, et al., Sequence-Selective Peptide Binding with a Peptido-A,B-*trans*-steroidal Receptor Selected from an Encoded Combinatorial Library, *J. Am. Chem. Soc.*, 118:1813-1814 (1996); Chu, et al., Affinity Capillary Electrophoresis to Identify the
- 30 Peptide in A Peptide Library that Binds Most Tightly to Vancomycin, *J. Org. Chem.*, 58:648-652 (1993); Clackson, et al., Making Antibody Fragments Using Phage Display Libraries, *Nature*, 352:624-628 (1991); Combs, et al., Protein

RECTIFIED SHEET (RULE 91)

- Structure-Based Combinatorial Chemistry: Discovery of Non-Peptide Binding Elements to Src SH3 Domain, *J. Am. Chem. Soc.*, 118:287-288 (1996); Cwirla, et al., Peptides On Phage: A Vast Library of Peptides for Identifying Ligands, *Proc. Natl. Acad. Sci. U.S.A.*, 87:6378-6382 (1990); Ecker, et al., Combinatorial
- 5 Drug Discovery: Which Method will Produce the Greatest Value, *Bio/Technology*, 13:351-360 (1995); Ellington, et al., *In Vitro* Selection of RNA Molecules That Bind Specific Ligands, *Nature*, 346:818-822 (1990); Ellman, J.A., Variants of Benzodiazepines, *J. Am. Chem. Soc.*, 114:10997 (1992); Erickson, et al., *The Proteins*; Neurath, H., Hill, R.L., Eds.: Academic: New York,
- 10 1976; pp. 255-257; Felici, et al., *J. Mol. Biol.*, 222:301-310 (1991); Fodor, et al., Light-Directed, Spatially Addressable Parallel Chemical Synthesis, *Science*, 251:767-773 (1991); Francisco, et al., Transport and Anchoring of Beta-Lactamase to the External Surface of *E. Coli.*, *Proc. Natl. Acad. Sci. U.S.A.*, 89:2713-2717 (1992); Georgiou, et al., Practical Applications of Engineering
- 15 Gram-Negative Bacterial Cell Surfaces, *TIBTECH*, 11:6-10 (1993); Geysen, et al., Use of peptide synthesis to probe viral antigens for epitopes to a resolution of a single amino acid, *Proc. Natl. Acad. Sci. U.S.A.*, 81:3998-4002 (1984); Glaser, et al., Antibody Engineering by Codon-Based Mutagenesis in a Filamentous Phage Vector System, *J. Immunol.*, 149:3903-3913 (1992); Gram, et al., In
- 20 vitro selection and affinity maturation of antibodies from a naive combinatorial immunoglobulin library, *Proc. Natl. Acad. Sci.*, 89:3576-3580 (1992); Han, et al., Liquid-Phase Combinatorial Synthesis, *Proc. Natl. Acad. Sci. U.S.A.*, 92:6419-6423 (1995); Hoogenboom, et al., Multi-Subunit Proteins on the Surface of Filamentous Phage: Methodologies for Displaying Antibody (Fab)
- 25 Heavy and Light Chains, *Nucleic Acids Res.*, 19:4133-4137 (1991); Houghten, et al., General Method for the Rapid Solid-Phase Synthesis of Large Numbers of Peptides: Specificity of Antigen-Antibody Interaction at the Level of Individual Amino Acids, *Proc. Natl. Acad. Sci. U.S.A.*, 82:5131-5135 (1985); Houghten, et al., The Use of Synthetic Peptide Combinatorial Libraries for the Determination
- 30 of Peptide Ligands in Radio-Receptor Assays-Opioid-Peptides, *Bioorg. Med. Chem. Lett.*, 3:405-412 (1993); Houghten, et al., Generation and Use of Synthetic Peptide Combinatorial Libraries for Basic Research and Drug Discovery,

- Nature*, 354:84-86 (1991); Huang, et al., Discovery of New Ligand Binding Pathways in Myoglobin by Random Mutagenesis, *Nature Struct. Biol.*, 1:226-229 (1994); Huse, et al., Generation of a Large Combinatorial Library of the Immunoglobulin Repertoire In Phage Lambda, *Science*, 246:1275-1281 (1989);
- 5 Janda, K.D., New Strategies for the Design of Catalytic Antibodies, *Biotechnol. Prog.*, 6:178-181 (1990); Jung, et al., Multiple Peptide Synthesis Methods and Their Applications, *Angew. Chem. Int. Ed. Engl.*, 31:367-486 (1992); Kang, et al., Linkage of Recognition and Replication Functions By Assembling Combinatorial Antibody Fab Libraries Along Phage Surfaces, *Proc. Natl. Acad.*
- 10 *Sci. U.S.A.*, 88:4363-4366 (1991a); Kang, et al., Antibody Redesign by Chain Shuffling from Random Combinatorial Immunoglobulin Libraries, *Proc. Natl. Acad. Sci. U.S.A.*, 88:11120-11123 (1991b); Kay, et al., An M13 Phage Library Displaying Random 38-Amino-Acid-Peptides as a Source of Novel Sequences with Affinity to Selected Targets Genes, *Gene*, 128:59-65 (1993); Lam, et al., A
- 15 new type of synthetic peptide library for identifying ligand-binding activity, *Nature*, 354:82-84 (1991) (published errata appear in *Nature*, 358:434 (1992) and *Nature*, 360:768 (1992); Lebl, et al., One Bead One Structure Combinatorial Libraries, *Biopolymers (Pept. Sci.)*, 37:177-198 (1995); Lerner, et al., Antibodies without Immunization, *Science*, 258:1313-1314 (1992); Li, et al., Minimization
- 20 of a Polypeptide Hormone, *Science*, 270:1657-1660 (1995); Light, et al., Display of Dimeric Bacterial Alkaline Phosphatase on the Major Coat Protein of Filamentous Bacteriophage, *Bioorg. Med. Chem. Lett.*, 3:1073-1079 (1992); Little, et al., Bacterial Surface Presentation of Proteins and Peptides: An Alternative to Phage Technology, *Trends Biotechnol.*, 11:3-5 (1993); Marks, et
- 25 al., By-Passing Immunization. Human Antibodies from V-Gene Libraries Displayed on Phage, *J. Mol. Biol.*, 222:581-597 (1991); Matthews, et al., Substrate Phage: Selection of Protease Substrates by Monovalent Phage Display, *Science*, 260:1113-1117 (1993); McCafferty, et al., Phage Enzymes: Expression and Affinity Chromatography of Functional Alkaline Phosphatase on the Surface
- 30 of Bacteriophage, *Protein Eng.*, 4:955-961 (1991); Menger, et al., Phosphatase Catalysis Developed Via Combinatorial Organic Chemistry, *J. Org. Chem.*, 60:6666-6667 (1995); Nicolaou, et al., *Angew. Chem. Int. Ed. Engl.*, 34:2289-

-94-

- 2291 (1995); Oldenburg, et al., Peptide Ligands for A Sugar-Binding Protein Isolated from a Random Peptide Library, *Proc. Natl. Acad. Sci. U.S.A.*, 89:5393-5397 (1992); Parmley, et al., Antibody-Selectable Filamentous fd Phage Vectors: Affinity Purification of Target Genes, *Genes*, 73:305-318 (1988); Pinilla, et al.,
- 5 Synthetic Peptide Combinatorial Libraries (SPCLS)--Identification of the Antigenic Determinant of Beta-Endorphin Recognized by Monoclonal Antibody-3E7, *Gene*, 128:71-76 (1993); Pinilla, et al., Review of the Utility of Soluble Combinatorial Libraries, *Biopolymers*, 37:221-240 (1995); Pistor, et al., Expression of Viral Hemagglutinin On the Surface of *E. Coli*, *Klin. Wochenschr.*, 66:110-116
- 10 (1989); Pollack, et al., Selective Chemical Catalysis by an Antibody, *Science*, 234:1570-1572 (1986); Rigler, et al., Fluorescence Correlations, Single Molecule Detection and Large Number Screening: Applications in Biotechnology, *J. Biotechnol.*, 41:177-186 (1995); Sarvetnick, et al., Increasing the Chemical Potential of the Germ-Line Antibody Repertoire, *Proc. Natl. Acad. Sci. U.S.A.*,
- 15 90:4008-4011 (1993); Sastry, et al., Cloning of the Immunological Repertoire in *Escherichia Coli* for Generation of Monoclonal Catalytic Antibodies: Construction of a Heavy Chain Variable Region-Specific cDNA Library, *Proc. Natl. Acad. Sci. U.S.A.*, 86:5728-5732 (1989); Scott, et al., Searching for Peptide Ligands with an Epitope Library, *Science*, 249:386-390 (1990); Sears, et al., Engineering
- 20 Enzymes for Bioorganic Synthesis: Peptide Bond Formation, *Biotechnol. Prog.*, 12:423-433 (1996); Simon, et. al., Peptides: A Modular Approach to Drug Discovery, *Proc. Natl. Acad. Sci. U.S.A.*, 89:9367-9371 (1992); Still, et al., Discovery of Sequence-Selective Peptide Binding by Synthetic Receptors Using Encoded Combinatorial Libraries, *Acc. Chem. Res.*, 29:155-163 (1996);
- 25 Thompson, et al., Synthesis and Applications of Small Molecule Libraries, *Chem. Rev.*, 96:555-600 (1996); Tramontano, et al., Catalytic Antibodies, *Science*, 234:1566-1570 (1986); Wrighton, et al., Small Peptides as Potent Mimetics of the Protein Hormone Erythropoietin, *Science*, 273:458-464 (1996); York, et al., Combinatorial mutagenesis of the reactive site region in plasminogen activator
- 30 inhibitor I, *J. Biol. Chem.*, 266:8595-8600 (1991); Zebedee, et al., Human Combinatorial Antibody Libraries to Hepatitis B Surface Antigen, *Proc. Natl. Acad. Sci. U.S.A.*, 89:3175-3179 (1992); Zuckerman, et al., Identification of

RECTIFIED SHEET (RULE 91)

-95-

Highest-Affinity Ligands by Affinity Selection from Equimolar Peptide Mixtures Generated by Robotic Synthesis, *Proc. Natl. Acad. Sci. U.S.A.*, 89:4505-4509 (1992).

For example, peptides that bind to an MTSP10 polypeptide or a protease domain of an SP protein can be identified using phage display libraries. In an exemplary embodiment, this method can include a) contacting phage from a phage library with the MTSP10 polypeptide or a protease domain thereof; (b) isolating phage that bind to the protein; and (c) determining the identity of at least one peptide coded by the isolated phage to identify a peptide that binds to an MTSP10 polypeptide.

H. Modulators of the activity of MTSP10 polypeptides

Provided herein are compounds, identified by screening or produced using the MTSP10 polypeptide or protease domain in other screening methods, that modulate the activity of an MTSP10. These compounds act by directly interacting with the MTSP10 polypeptide or by altering transcription or translation thereof. Such molecules include, but are not limited to, antibodies that specifically react with an MTSP10 polypeptide, particularly with the protease domain thereof, antisense nucleic acids or double-stranded RNA (dsRNA) such as RNAi, that alter expression of the MTSP10 polypeptide, antibodies, peptide mimetics and other such compounds.

1. Antibodies

Antibodies, including polyclonal and monoclonal antibodies, that specifically bind to the MTSP10 polypeptide provided herein, particularly to the single chain protease domains thereof or the activated forms of the full-length or protease domain or the zymogen form, are provided.

Generally, the antibody is a monoclonal antibody, and typically the antibody specifically binds to the protease domain of the MTSP10 polypeptide. In particular embodiments, antibodies to each of the single chain and or two chain form of the protease domain of MTSP10 are provided. Also provided are antibodies that specifically bind to any domain of MTSP10 and to two chain forms thereof.

-96-

The MTSP10 polypeptide and domains, fragments, homologs and derivatives thereof can be used as immunogens to generate antibodies that specifically bind such immunogens. Such antibodies include but are not limited to polyclonal, monoclonal, chimeric, single chain, Fab fragments, and an Fab expression library. In a specific embodiment, antibodies to human MTSP10 polypeptide are produced. In another embodiment, complexes formed from fragments of an MTSP10 polypeptide, that contain the serine protease domain are used as immunogens for antibody production.

Various procedures known in the art can be used for the production of polyclonal antibodies to MTSP10 polypeptide, its domains, derivatives, fragments or analogs. For production of the antibody, various host animals can be immunized by injection with the native MTSP10 polypeptide or a synthetic version, or a derivative of the foregoing, such as a cross-linked MTSP10 polypeptide. Such host animals include but are not limited to rabbits, mice, rats, etc. Various adjuvants can be used to increase the immunological response, depending on the host species, and include but are not limited to Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, dinitrophenol, and potentially useful human adjuvants such as bacille Calmette-Guerin (BCG) and corynebacterium parvum.

For preparation of monoclonal antibodies directed towards an MTSP10 polypeptide or domains, derivatives, fragments or analogs thereof, any technique that provides for the production of antibody molecules by continuous cell lines in culture can be used. Such techniques include but are not restricted to the hybridoma technique originally developed by Kohler and Milstein (*Nature* 256:495-497 (1975)), the trioma technique, the human B-cell hybridoma technique (Kozbor et al., *Immunology Today* 4:72 (1983)), and the EBV hybridoma technique to produce human monoclonal antibodies (Cole et al., in *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96 (1985)). In an additional embodiment, monoclonal antibodies can be produced in germ-free animals utilizing recent technology (PCT/US90/02545). Human antibodies can be used and can be obtained by using human hybridomas (Cote

-97-

et al., *Proc. Natl. Acad. Sci. USA* 80:2026-2030 (1983)), or by transforming human B cells with EBV virus *in vitro* (Cole et al., in *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96 (1985)). Techniques developed for the production of "chimeric antibodies" (Morrison et al., *Proc. Natl. Acad. Sci. USA* 81:6851-6855 (1984); Neuberger et al., *Nature* 312:604-608 (1984); Takeda et al., *Nature* 314:452-454 (1985)) by splicing the genes from a mouse antibody molecule specific for the MTSP10 polypeptide together with genes from a human antibody molecule of appropriate biological activity can be used.

MTSP10-encoding nucleic acid molecules or portions thereof can be used in DNA immunization protocols to produce antibodies that bind to MTSP10 (see, e.g., U.S. Patent No. 5,795,872 and U.S. Patent No. 5,643,578 and U.S. Patent No. 6,337,072).

Techniques described for the production of single chain antibodies (U.S. patent 4,946,778) can be adapted to produce MTSP10 polypeptide-specific single chain antibodies. An additional embodiment uses the techniques described for the construction of Fab expression libraries (Huse et al., *Science* 246:1275-1281 (1989)) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity for MTSP10 polypeptide or domains, derivatives, or analogs thereof. Non-human antibodies can be "humanized" by known methods (see, e.g., U.S. Patent No. 5,225,539).

Antibody fragments that specifically bind to MTSP10 polypeptide or epitopes thereof can be generated by techniques known in the art. For example, such fragments include but are not limited to: the F(ab')₂ fragment, which can be produced by pepsin digestion of the antibody molecule; the Fab' fragments that can be generated by reducing the disulfide bridges of the F(ab')₂ fragment; the Fab fragments that can be generated by treating the antibody molecule with papain and a reducing agent; and Fv fragments.

In the production of antibodies, screening for the desired antibody can be accomplished by techniques known in the art, e.g., ELISA (enzyme-linked immunosorbent assay). To select antibodies specific for a particular domain of the MTSP10 polypeptide one can assay generated hybridomas for a product that binds to the fragment of the MTSP10 polypeptide that contains such a domain.

RECTIFIED SHEET (RULE 91)

-98-

The foregoing antibodies can be used in methods known in the art relating to the localization and/or quantitation of MTSP10 polypeptide proteins, *e.g.*, for imaging these proteins, measuring levels thereof in appropriate physiological samples, in, for example, diagnostic methods. In another
5 embodiment, anti-MTSP10 polypeptide antibodies, or fragments thereof, containing the binding domain are used as therapeutic agents.

2. Peptides, Polypeptides and Peptide Mimetics

Provided herein are methods for identifying molecules that bind to and modulate the activity of SP proteins. Included among molecules that bind to
10 SPs, particularly the single chain protease domain or catalytically active fragments thereof, are peptides, polypeptides and peptide mimetics, including cyclic peptides. Peptide mimetics are molecules or compounds that mimic the necessary molecular conformation of a ligand or polypeptide for specific binding to a target molecule such as an MTSP10 polypeptide. In an exemplary
15 embodiment, the peptides, polypeptides and peptide mimetics bind to the protease domain of the MTSP10 polypeptide. Such peptides and peptide mimetics include those of antibodies that specifically bind to an MTSP10 polypeptide and, typically, bind to the protease domain of an MTSP10 polypeptide. The peptides, polypeptides and peptide mimetics identified by
20 methods provided herein can be agonists or antagonists of MTSP10 polypeptides.

Such peptides, polypeptides and peptide mimetics are useful for diagnosing, treating, preventing, and screening for a disease or disorder associated with MTSP10 polypeptide activity in a mammal. In addition, the
25 peptides and peptide mimetics are useful for identifying, isolating, and purifying molecules or compounds that modulate the activity of an MTSP10 polypeptide, or specifically bind to an MTSP10 polypeptide, generally the protease domain of an MTSP10 polypeptide. Low molecular weight peptides and peptide mimetics can have strong binding properties to a target molecule, *e.g.*, an MTSP10
30 polypeptide or the protease domain of an MTSP10 polypeptide.

Peptides, polypeptides and peptide mimetics that bind to MTSP10 polypeptides as described herein can be administered to mammals, including

-99-

humans, to modulate MTSP10 polypeptide activity. Thus, methods for therapeutic treatment and prevention of neoplastic diseases comprise administering a peptide, polypeptide or peptide mimetic compound in an amount sufficient to modulate such activity are provided. Thus, also provided herein are

5 methods for treating a subject having such a disease or disorder in which a peptide, polypeptide or peptide mimetic compound is administered to the subject in a therapeutically effective dose or amount.

Compositions containing the peptides, polypeptides or peptide mimetics can be administered for prophylactic and/or therapeutic treatments. In

10 therapeutic applications, compositions can be administered to a patient already suffering from a disease, as described above, in an amount sufficient to cure or at least partially arrest the symptoms of the disease and its complications. Amounts effective for this use will depend on the severity of the disease and the weight and general state of the patient and can be empirically determined.

15 In prophylactic applications, compositions containing the peptides, polypeptides and peptide mimetics are administered to a patient susceptible to or otherwise at risk of a particular disease. Such an amount is defined to be a "prophylactically effective dose." In this use, the precise amounts again depend on the patient's state of health and weight. Accordingly, the peptides,

20 polypeptides and peptide mimetics that bind to an MTSP10 polypeptide can be used to prepare pharmaceutical compositions containing, as an active ingredient, at least one of the peptides, polypeptides or peptide mimetics in association with a pharmaceutical carrier or diluent. The compounds can be administered, for example, by oral, pulmonary, parenteral (intramuscular, intraperitoneal,

25 intravenous (IV) or subcutaneous injection), inhalation (via a fine powder formulation), transdermal, nasal, vaginal, rectal, or sublingual routes of administration and can be formulated in dosage forms appropriate for each route of administration (see, e.g., International PCT application Nos. WO 93/25221 and WO 94/17784; and European Patent Application 613,683).

30 Peptides, polypeptides and peptide mimetics that bind to MTSP10 polypeptides are useful *in vitro* as unique tools for understanding the biological role of MTSP10 polypeptides, including the evaluation of the many factors

RECTIFIED SHEET (RULE 91)

-100-

thought to influence, and be influenced by, the production of MTSP10 polypeptide. Such peptides, polypeptides and peptide mimetics are also useful in the development of other compounds that bind to and modulate the activity of an MTSP10 polypeptide, because such compounds provide important
5 information on the relationship between structure and activity that should facilitate such development.

The peptides, polypeptides and peptide mimetics are also useful as competitive binders in assays to screen for new MTSP10 polypeptides or MTSP10 polypeptide agonists. In such assay embodiments, the compounds can
10 be used without modification or can be modified in a variety of ways; for example, by labeling, such as covalently or non-covalently joining a moiety which directly or indirectly provides a detectable signal. In any of these assays, the materials thereto can be labeled either directly or indirectly. Possibilities for direct labeling include label groups such as: radiolabels such as ¹²⁵I enzymes
15 (U.S. Pat. No. 3,645,090) such as peroxidase and alkaline phosphatase, and fluorescent labels (U.S. Pat. No. 3,940,475) capable of monitoring the change in fluorescence intensity, wavelength shift, or fluorescence polarization. Possibilities for indirect labeling include biotinylation of one constituent followed by binding to avidin coupled to one of the above label groups. The compounds
20 can also include spacers or linkers in cases where the compounds are to be attached to a solid support.

Moreover, based on their ability to bind to an MTSP10 polypeptide, the peptides, polypeptides and peptide mimetics can be used as reagents for detecting MTSP10 polypeptides in living cells, fixed cells, in biological fluids, in
25 tissue homogenates and in purified, natural biological materials. For example, by labelling such peptides, polypeptides and peptide mimetics, cells having MTSP10 polypeptides can be identified. In addition, based on their ability to bind an MTSP10 polypeptide, the peptides, polypeptides and peptide mimetics can be used in *in situ* staining, FACS (fluorescence-activated cell sorting),
30 Western blotting, ELISA and other analytical protocols. Based on their ability to bind to an MTSP10 polypeptide, the peptides, polypeptides and peptide mimetics can be used in purification of MTSP10 polypeptides or in purifying cells

-101-

expressing the MTSP10 polypeptides, e.g., a polypeptide encoding the protease domain of an MTSP10 polypeptide.

The peptides, polypeptides and peptide mimetics can also be used as commercial reagents for various medical research and diagnostic uses. The activity of the peptides and peptide mimetics can be evaluated either *in vitro* or *in vivo* in one of the numerous models described in McDonald (1992) *Am. J. of Pediatric Hematology/Oncology*, 14:8-21.

3. Peptide, polypeptide and peptide mimetic therapy

Peptide analogs are commonly used in the pharmaceutical industry as non-peptide drugs with properties analogous to those of the template peptide. These types of non-peptide compounds are termed "peptide mimetics" or "peptidomimetics" (Luthman *et al.*, *A Textbook of Drug Design and Development*, 14:386-406, 2nd Ed., Harwood Academic Publishers (1996); Joachim Grante (1994) *Angew. Chem. Int. Ed. Engl.*, 33:1699-1720; Fauchere (1986) *J. Adv. Drug Res.*, 15:29; Veber and Freidinger (1985) *TINS*, p. 392; and Evans *et al.* (1987) *J. Med. Chem.* 30:1229). Peptide mimetics that are structurally similar to therapeutically useful peptides can be used to produce an equivalent or enhanced therapeutic or prophylactic effect. Preparation of peptidomimetics and structures thereof are known to those of skill in this art.

Systematic substitution of one or more amino acids of a consensus sequence with a D-amino acid of the same type (e.g., D-lysine in place of L-lysine) can be used to generate more stable peptides. In addition, constrained peptides containing a consensus sequence or a substantially identical consensus sequence variation can be generated by methods known in the art (Rizo *et al.* (1992) *An. Rev. Biochem.*, 61:387, incorporated herein by reference); for example, by adding internal cysteine residues capable of forming intramolecular disulfide bridges which cyclize the peptide.

Those skilled in the art appreciate that modifications can be made to the peptides and mimetics without deleteriously effecting the biological or functional activity of the peptide. Further, the skilled artisan would know how to design non-peptide structures in three dimensional terms, that mimic the peptides that bind to a target molecule, e.g., an MTSP10 polypeptide or, generally, the

-102-

protease domain of MTSP10 polypeptides (see, *e.g.*, Eck and Sprang (1989) *J. Biol. Chem.*, 26: 17605-18795).

When used for diagnostic purposes, the peptides and peptide mimetics can be labeled with a detectable label and, accordingly, the peptides and peptide
5 mimetics without such a label can serve as intermediates in the preparation of labeled peptides and peptide mimetics. Detectable labels can be molecules or compounds, which when covalently attached to the peptides and peptide mimetics, permit detection of the peptide and peptide mimetics *in vivo*, for example, in a patient to whom the peptide or peptide mimetic has been
10 administered, or *in vitro*, *e.g.*, in a sample or cells. Suitable detectable labels are well known in the art and include, by way of example, radioisotopes, fluorescent labels (*e.g.*, fluorescein), and the like. The particular detectable label employed is not critical and is selected to be detectable at non-toxic levels. Selection of the such labels is well within the skill of the art.

15 Covalent attachment of a detectable label to the peptide or peptide mimetic is accomplished by conventional methods well known in the art. For example, when the ¹²⁵I radioisotope is employed as the detectable label, covalent attachment of ¹²⁵I to the peptide or the peptide mimetic can be achieved by incorporating the amino acid tyrosine into the peptide or peptide mimetic and
20 then iodinating the peptide (see, *e.g.*, Weaner *et al.* (1994) *Synthesis and Applications of Isotopically Labelled Compounds*, pp. 137-140). If tyrosine is not present in the peptide or peptide mimetic, incorporation of tyrosine to the N or C terminus of the peptide or peptide mimetic can be achieved by well known chemistry. Likewise, ³²P can be incorporated onto the peptide or peptide
25 mimetic as a phosphate moiety through, for example, a hydroxyl group on the peptide or peptide mimetic using conventional chemistry.

Labeling of peptidomimetics usually involves covalent attachment of one or more labels, directly or through a spacer (*e.g.*, an amide group), to non-interfering position(s) on the peptidomimetic that are predicted by
30 quantitative structure-activity data and/or molecular modeling. Such non-interfering positions generally are positions that do not form direct contacts with the macromolecules(s) to which the peptidomimetic binds to produce the

-103-

therapeutic effect. Derivatization (e.g., labeling) of peptidomimetics should not substantially interfere with the desired biological or pharmacological activity of the peptidomimetic.

Peptides, polypeptides and peptide mimetics that can bind to an MTSP10 polypeptide or the protease domain of MTSP10 polypeptides and/or modulate the activity thereof, or exhibit MTSP10 polypeptide activity, can be used for treatment of neoplastic disease. The peptides, polypeptides and peptide mimetics can be delivered, *in vivo* or *ex vivo*, to the cells of a subject in need of treatment. Further, peptides which have MTSP10 polypeptide activity can be delivered, *in vivo* or *ex vivo*, to cells which carry mutant or missing alleles encoding the MTSP10 polypeptide gene. Any of the techniques described herein or known to the skilled artisan can be used for preparation and *in vivo* or *ex vivo* delivery of such peptides, polypeptides and peptide mimetics that are substantially free of other human proteins. For example, the peptides, polypeptides and peptide mimetics can be readily prepared by expression in a microorganism or synthesis *in vitro*.

The peptides or peptide mimetics can be introduced into cells, *in vivo* or *ex vivo*, by microinjection or by use of liposomes, for example. Alternatively, the peptides, polypeptides or peptide mimetics can be taken up by cells, *in vivo* or *ex vivo*, actively or by diffusion. In addition, extracellular application of the peptide, polypeptide or peptide mimetic can be sufficient to effect treatment of a neoplastic disease. Other molecules, such as drugs or organic compounds, that: 1) bind to a MTSP10 polypeptide or protease domain thereof; or 2) have a similar function or activity to an MTSP10 polypeptide or protease domain thereof, can be used in methods for treatment.

4. Rational drug design

The goal of rational drug design is to produce structural analogs of biologically active polypeptides or peptides of interest or of small molecules or peptide mimetics with which they interact (e.g., agonists and antagonists) in order to fashion drugs which are, e.g., more active or stable forms thereof; or which, for example, enhance or interfere with the function of a polypeptide *in vivo* (e.g., an MTSP10 polypeptide). In one approach, one first determines the

-104-

three-dimensional structure of a protein of interest (e.g., an MTSP10 polypeptide or polypeptide having a protease domain) or, for example, of an MTSP10 polypeptide-ligand complex, by X-ray crystallography, by computer modeling or most typically, by a combination of approaches (see, e.g., Erickson *et al.* 1990).

5 Also, useful information regarding the structure of a polypeptide can be gained by modeling based on the structure of homologous proteins. In addition, peptides can be analyzed by an alanine scan. In this technique, an amino acid residue is replaced by Ala, and its effect on the peptide's activity is determined. Each of the amino acid residues of the peptide is analyzed in this manner to
10 determine the important regions of the peptide.

Also, a polypeptide or peptide that binds to an MTSP10 polypeptide or, generally, the protease domain of an MTSP10 polypeptide, can be selected by a functional assay, and then the crystal structure of this polypeptide or peptide can be determined. The polypeptide can be, for example, an antibody specific
15 for an MTSP10 polypeptide or the protein domain of an MTSP10 polypeptide. This approach can yield a pharmacophore upon which subsequent drug design can be based. Further, it is possible to bypass the crystallography altogether by generating anti-idiotypic polypeptides or peptides, (anti-ids) to a functional, pharmacologically active polypeptide or peptide that binds to an MTSP10
20 polypeptide or protease domain of an MTSP10 polypeptide. As a mirror image of a mirror image, the binding site of the anti-ids is expected to be an analog of the original target molecule, e.g., an MTSP10 polypeptide or polypeptide having an MTSP10 polypeptide. The anti-id could then be used to identify and isolate peptides from banks of chemically or biologically produced banks of peptides.
25 Selected peptides would then act as the pharmacophore.

Thus, one can design drugs which have, for example, improved activity or stability or which act as modulators (e.g., inhibitors, agonists or antagonists) of MTSP10 polypeptide activity, and are useful in the methods, particularly the methods for diagnosis, treatment, prevention, and screening of a neoplastic
30 disease. By virtue of the availability of nucleic acid that encodes MTSP10 polypeptides, sufficient amounts of the MTSP10 polypeptide can be made available to perform such analytical studies as X-ray crystallography. In addition,

-105-

the knowledge of the amino acid sequence of an MTSP10 polypeptide or the protease domain thereof, e.g., the protease domain encoded by the amino acid sequence of SEQ ID Nos. 23 and 6, can provide guidance on computer modeling techniques in place of, or in addition to, X-ray crystallography.

5 **Methods of identifying peptides and peptide mimetics that bind to MTSP10 polypeptides**

Peptides having a binding affinity to the MTSP10 polypeptide provided herein (e.g., an MTSP10 polypeptide or a polypeptide having a protease domain of an MTSP10 polypeptide) can be readily identified, for example, by random
10 peptide diversity generating systems coupled with an affinity enrichment process. Specifically, random peptide diversity generating systems include the "peptides on plasmids" system (see, e.g., U.S. Patent Nos. 5,270,170 and 5,338,665); the "peptides on phage" system (see, e.g., U.S. Patent No. 6,121,238 and Cwirla, *et al.* (1990) *Proc. Natl. Acad. Sci. U.S.A.*
15 87:6378-6382); the "polysome system;" the "encoded synthetic library (ESL)" system; and the "very large scale immobilized polymer synthesis" system (see, e.g., U.S. Patent No. 6,121,238; and Dower *et al.* (1991) *An. Rep. Med. Chem.* 26:271-280

For example, using the procedures described above, random peptides can
20 generally be designed to have a defined number of amino acid residues in length (e.g., 12). To generate the collection of oligonucleotides encoding the random peptides, the codon motif (NNK)x, where N is nucleotide A, C, G, or T (equimolar; depending on the methodology employed, other nucleotides can be employed), K is G or T (equimolar), and x is an integer corresponding to the
25 number of amino acids in the peptide (e.g., 12) can be used to specify any one of the 32 possible codons resulting from the NNK motif: 1 for each of 12 amino acids, 2 for each of 5 amino acids, 3 for each of 3 amino acids, and only one of the three stop codons. Thus, the NNK motif encodes all of the amino acids, encodes only one stop codon, and reduces codon bias.

30 The random peptides can be presented, for example, either on the surface of a phage particle, as part of a fusion protein containing either the pIII or the pVIII coat protein of a phage fd derivative (peptides on phage) or as a fusion

-106-

protein with the LacI peptide fusion protein bound to a plasmid (peptides on plasmids). The phage or plasmids, including the DNA encoding the peptides, can be identified and isolated by an affinity enrichment process using immobilized MTSP10 polypeptide having a protease domain. The affinity enrichment
5 process, sometimes called "panning," typically involves multiple rounds of incubating the phage, plasmids, or polysomes with the immobilized MTSP10 polypeptide, collecting the phage, plasmids, or polysomes that bind to the MTSP10 polypeptide (along with the accompanying DNA or mRNA), and producing more of the phage or plasmids (along with the accompanying
10 LacI-peptide fusion protein) collected.

Characteristics of peptides and peptide mimetics

Among the peptides, polypeptides and peptide mimetics for therapeutic application are those of having molecular weights from about 250 to about 8,000 daltons. If such peptides are oligomerized, dimerized and/or derivatized
15 with a hydrophilic polymer (*e.g.*, to increase the affinity and/or activity of the compounds), the molecular weights of such peptides can be substantially greater and can range anywhere from about 500 to about 120,000 daltons, generally from about 8,000 to about 80,000 daltons. Such peptides can contain 9 or more amino acids that are naturally occurring or synthetic (non-naturally
20 occurring) amino acids. One skilled in the art can determine the affinity and molecular weight of the peptides and peptide mimetics suitable for therapeutic and/or diagnostic purposes (*e.g.*, see Dower *et al.*, U.S. Patent No. 6,121,238).

The peptides can be covalently attached to one or more of a variety of hydrophilic polymers. Suitable hydrophilic polymers include, but are not limited
25 to, polyalkylethers as exemplified by polyethylene glycol and polypropylene glycol, polylactic acid, polyglycolic acid, polyoxyalkenes, polyvinylalcohol, polyvinylpyrrolidone, cellulose and cellulose derivatives, dextran and dextran derivatives. When the peptide compounds are derivatized with such polymers, their solubility and circulation half-lives can be increased with little, if any,
30 diminishment in their binding activity. The peptide compounds can be dimerized and each of the dimeric subunits can be covalently attached to a hydrophilic

-107-

polymer. The peptide compounds can be PEGylated, i.e., covalently attached to polyethylene glycol (PEG).

5. Methods of preparing peptides and peptide mimetics

Peptides that bind to MTSP10 polypeptides can be prepared by classical methods known in the art, for example, by using standard solid phase techniques. The standard methods include exclusive solid phase synthesis, partial solid phase synthesis methods, fragment condensation, classical solution synthesis, and even by recombinant DNA technology (see, *e.g.*, Merrifield (1963) *J. Am. Chem. Soc.*, 85:2149, incorporated herein by reference.)

Using the "encoded synthetic library" or "very large scale immobilized polymer synthesis" systems (see, *e.g.*, U.S. Patent No. 5,925,525, and 5,902,723), the minimum size of a peptide with the activity of interest can be determined. In addition all peptides that form the group of peptides that differ from the desired motif (or the minimum size of that motif) in one, two, or more residues can be prepared. This collection of peptides then can be screened for the ability to bind to the target molecule, *e.g.*, MTSP10 polypeptide or, generally, the protease domain of an MTSP10 polypeptide. This immobilized polymer synthesis system or other peptide synthesis methods can also be used to synthesize truncation analogs and deletion analogs and combinations of truncation and deletion analogs of the peptide compounds.

These procedures can also be used to synthesize peptides in which amino acids other than the 20 naturally occurring, genetically encoded amino acids are substituted at one, two, or more positions of the peptide. For instance, naphthylalanine can be substituted for tryptophan, facilitating synthesis. Other synthetic amino acids that can be substituted into the peptides include L-hydroxypropyl, L-3, 4-dihydroxy-phenylalanyl, D amino acids such as L-D-hydroxylysyl and D-D-methylalanyl, L- α -methylalanyl, β amino acids, and isoquinolyl. D amino acids and non-naturally occurring synthetic amino acids can also be incorporated into the peptides (see, *e.g.*, Roberts *et al.* (1983) *Unusual Amino/Acids in Peptide Synthesis*, 5(6):341-449).

The peptides also can be modified by phosphorylation (see, *e.g.*, W. Bannwarth *et al.* (1996) *Biorganic and Medicinal Chemistry Letters*,

-108-

6(17):2141-2146), and other methods for making peptide derivatives (see, *e.g.*, Hruby *et al.* (1990) *Biochem. J.*, 268(2):249-262). Thus, peptide compounds also serve as a basis to prepare peptide mimetics with similar or improved biological activity.

- 5 Those of skill in the art recognize that a variety of techniques are available for constructing peptide mimetics with the same or similar desired biological activity as the corresponding peptide compound but with more favorable activity than the peptide with respect to solubility, stability, and susceptibility to hydrolysis and proteolysis (see, *e.g.*, Morgan *et al.* (1989) *An.*
10 *Rep. Med. Chem.*, 24:243-252). Methods for preparing peptide mimetics modified at the N-terminal amino group, the C-terminal carboxyl group, and/or changing one or more of the amido linkages in the peptide to a non-amido linkage are known to those of skill in the art.

- Amino terminus modifications include, but are not limited to, alkylating,
15 acetylating and adding a carbobenzoyl group, forming a succinimide group (see, *e.g.*, Murray *et al.* (1995) *Burger's Medicinal Chemistry and Drug Discovery*, 5th ed., Vol. 1, Manfred E. Wolf, ed., John Wiley and Sons, Inc.). C-terminal modifications include mimetics wherein the C-terminal carboxyl group is replaced by an ester, an amide or modifications to form a cyclic peptide.

- 20 In addition to N-terminal and C-terminal modifications, the peptide compounds, including peptide mimetics, can advantageously be modified with or covalently coupled to one or more of a variety of hydrophilic polymers. It has been found that when peptide compounds are derivatized with a hydrophilic polymer, their solubility and circulation half-lives can be increased and their
25 immunogenicity is masked, with little, if any, diminishment in their binding activity. Suitable nonproteinaceous polymers include, but are not limited to, polyalkylethers as exemplified by polyethylene glycol and polypropylene glycol, polylactic acid, polyglycolic acid, polyoxyalkenes, polyvinylalcohol, polyvinylpyrrolidone, cellulose and cellulose derivatives, dextran and dextran
30 derivatives. Generally, such hydrophilic polymers have an average molecular weight ranging from about 500 to about 100,000 daltons, including from about 2,000 to about 40,000 daltons and, from about 5,000 to about 20,000 daltons.

The hydrophilic polymers also can have an average molecular weights of about 5,000 daltons, 10,000 daltons and 20,000 daltons.

Methods for derivatizing peptide compounds or for coupling peptides to such polymers have been described (see, *e.g.*, Zallipsky (1995) *Bioconjugate Chem.*, 6:150-165; Monfardini *et al.* (1995) *Bioconjugate Chem.*, 6:62-69; U.S. Pat. No. 4,640,835; U.S. Pat. No. 4,496,689; U.S. Pat. No. 4,301,144; U.S. Pat. No. 4,670,417; U.S. Pat. No. 4,791,192; U.S. Pat. No. 4,179,337 and WO 95/34326, all of which are incorporated by reference in their entirety herein).

Other methods for making peptide derivatives are described, for example, in Hruby *et al.* (1990), *Biochem J.*, 268(2):249-262, which is incorporated herein by reference. Thus, the peptide compounds also serve as structural models for non-peptidic compounds with similar biological activity. Those of skill in the art recognize that a variety of techniques are available for constructing compounds with the same or similar desired biological activity as a particular peptide compound but with more favorable activity with respect to solubility, stability, and susceptibility to hydrolysis and proteolysis (see, *e.g.*, Morgan *et al.* (1989) *An. Rep. Med. Chem.*, 24:243-252, incorporated herein by reference). These techniques include replacing the peptide backbone with a backbone composed of phosphonates, amidates, carbamates, sulfonamides, secondary amines, and N-methylamino acids.

Peptide compounds can exist in a cyclized form with an intramolecular disulfide bond between the thiol groups of the cysteines. Alternatively, an intermolecular disulfide bond between the thiol groups of the cysteines can be produced to yield a dimeric (or higher oligomeric) compound. One or more of the cysteine residues can also be substituted with a homocysteine.

I. Conjugates

A conjugate, containing: a) a single chain protease domain (or proteolytically active portion thereof) of an MTSP10 polypeptide or a full length zymogen, activated form thereof, or two or single chain protease domain thereof; and b) a targeting agent linked to the MTSP10 polypeptide directly or via a linker, wherein the agent facilitates: i) affinity isolation or purification of the conjugate; ii) attachment of the conjugate to a surface; iii) detection of the

-110-

conjugate; or iv) targeted delivery to a selected tissue or cell, is provided herein. The conjugate can be a chemical conjugate or a fusion protein mixture thereof.

The targeting agent can be a protein or peptide fragment, such as a tissue specific or tumor specific monoclonal antibody or growth factor or
5 fragment thereof linked either directly or via a linker to an MTSP10 polypeptide or a protease domain thereof. The targeting agent can also be a protein or peptide fragment that contains a protein binding sequence, a nucleic acid binding sequence, a lipid binding sequence, a polysaccharide binding sequence, or a metal binding sequence, or a linker for attachment to a solid support. In a
10 particular embodiment, the conjugate contains a) the MTSP10 or portion thereof, as described herein; and b) a targeting agent linked to the MTSP10 polypeptide directly or via a linker.

Conjugates, such as fusion proteins and chemical conjugates, of the MTSP10 polypeptide with a protein or peptide fragment (or plurality thereof) that
15 functions, for example, to facilitate affinity isolation or purification of the MTSP10 polypeptide domain, attachment of the MTSP10 polypeptide domain to a surface, or detection of the MTSP10 polypeptide domain are provided. The conjugates can be produced by chemical conjugation, such as via thiol linkages, and can be produced by recombinant means as fusion proteins. In the fusion
20 protein, the peptide or fragment thereof is linked to either the N-terminus or C-terminus of the MTSP10 polypeptide domain. In chemical conjugates the peptide or fragment thereof can be linked anywhere that conjugation can be effected, and there can be a plurality of such peptides or fragments linked to a single MTSP10 polypeptide domain or to a plurality thereof.

25 The targeting agent is for *in vitro* or *in vivo* delivery to a cell or tissue, and includes agents such as cell or tissue-specific antibodies, growth factors and other factors (including compounds) that bind to moieties expressed on specific cells; and other cell or tissue specific agents that promote directed delivery of a linked protein. The targeting agent can be one that specifically delivers the

-111-

MTSP10 polypeptide to selected cells by interaction with a cell surface protein and internalization of conjugate or MTSP10 polypeptide portion thereof.

These conjugates are used in a variety of methods and are particularly suited for use in methods of activation of prodrugs, such as prodrugs that upon
5 cleavage by the particular MTSP10, which is localized at or near the targeted cell or tissue, are cytotoxic. The prodrugs are administered prior to, or simultaneously with, or subsequently to the conjugate. Upon delivery to the targeted cells, the protease activates the prodrug, which then exhibits a therapeutic effect, such as a cytotoxic effect.

10 **1. Conjugation**

Conjugates with linked MTSP10 polypeptide domains can be prepared either by chemical conjugation, recombinant DNA technology, or combinations of recombinant expression and chemical conjugation. The MTSP10 polypeptide domains and the targeting agent can be linked in any orientation and more than
15 one targeting agent and/or MTSP10 polypeptide domain can be present in a conjugate.

a. Fusion proteins

Fusion proteins are provided herein. A fusion protein contains: a) one or a plurality of domains of an MTSP10 polypeptide and b) a targeting agent. The
20 fusion proteins are generally produced by recombinant expression of nucleic acids that encode the fusion protein.

b. Chemical conjugation

To effect chemical conjugation herein, the MTSP10 polypeptide domain is linked via one or more selected linkers or directly to the targeting agent.
25 Chemical conjugation must be used if the targeted agent is other than a peptide or protein, such as a nucleic acid or a non-peptide drug. Any means known to those of skill in the art for chemically conjugating selected moieties can be used.

2. Linkers

Linkers for two purposes are contemplated herein. The conjugates can
30 include one or more linkers between the MTSP10 polypeptide portion and the targeting agent. Additionally, linkers are used for facilitating or enhancing immobilization of an MTSP10 polypeptide or portion thereof on a solid support,

RECTIFIED SHEET (RULE 91)

-112-

such as a microtiter plate, silicon or silicon-coated chip, glass or plastic support, such as for high throughput solid phase screening protocols.

Any linker known to those of skill in the art for preparation of conjugates can be used herein. These linkers are typically used in the preparation of

5 chemical conjugates; peptide linkers can be incorporated into fusion proteins.

Linkers can be any moiety suitable to associate a domain of MTSP10 polypeptide and a targeting agent. Such linkers and linkages include, but are not limited to, peptidic linkages, amino acid and peptide linkages, typically containing between one and about 60 amino acids, more generally between about 10 and

10 30 amino acids, chemical linkers, such as heterobifunctional cleavable cross-linkers, including but are not limited to, N-succinimidyl (4-iodoacetyl)-aminobenzoate, sulfosuccinimidyl (4-iodoacetyl)-aminobenzoate, 4-succinimidyl-oxy carbonyl- α -(2-pyridyldithio)toluene, sulfosuccinimidyl-6-[α -methyl- α -(pyridyldithiol)-toluamido] hexanoate, N-succinimidyl-3-(-2-pyridyldithio) -

15 propionate, succinimidyl 6[3-(-2-pyridyldithio)-propionamido] hexanoate, sulfosuccinimidyl 6[3-(-2-pyridyldithio)-propionamido] hexanoate, 3-(2-pyridyldithio)-propionyl hydrazide, Ellman's reagent, dichlorotriazinic acid, and S-(2-thiopyridyl)-L-cysteine. Other linkers include, but are not limited to peptides and other moieties that reduce steric hindrance between the domain of MTSP10

20 polypeptide and the targeting agent, intracellular enzyme substrates, linkers that increase the flexibility of the conjugate, linkers that increase the solubility of the conjugate, linkers that increase the serum stability of the conjugate, photocleavable linkers and acid cleavable linkers.

Other exemplary linkers and linkages that are suitable for chemically

25 linked conjugates include, but are not limited to, disulfide bonds, thioether bonds, hindered disulfide bonds, and covalent bonds between free reactive groups, such as amine and thiol groups. These bonds are produced using heterobifunctional reagents to produce reactive thiol groups on one or both of the polypeptides and then reacting the thiol groups on one polypeptide with

30 reactive thiol groups or amine groups to which reactive maleimido groups or thiol groups can be attached on the other. Other linkers include, acid cleavable linkers, such as bismaleimideoxy propane, acid labile-transferrin conjugates

RECTIFIED SHEET (RULE 91)

-113-

and adipic acid dihydrazide, that would be cleaved in more acidic intracellular compartments; cross linkers that are cleaved upon exposure to UV or visible light; and linkers, such as various domains, such as C_H1, C_H2, and C_H3, from the constant region of human IgG₁ (see, Batra *et al. Molecular Immunol.*, 30:379-386 (1993)). In some embodiments, several linkers can be included in order to take advantage of desired properties of each linker.

Chemical linkers and peptide linkers can be inserted by covalently coupling the linker to the domain of MTSP10 polypeptide and the targeting agent. The heterobifunctional agents, described below, can be used to effect such covalent coupling. Peptide linkers can also be linked by expressing DNA encoding the linker and therapeutic agent (TA), linker and targeted agent, or linker, targeted agent and therapeutic agent (TA) as a fusion protein. Flexible linkers and linkers that increase solubility of the conjugates are contemplated for use, either alone or with other linkers are also contemplated herein.

a) Acid cleavable, photocleavable and heat sensitive linkers

Acid cleavable linkers, photocleavable and heat sensitive linkers can also be used, particularly where it can be necessary to cleave the domain of MTSP10 polypeptide to permit it to be more readily accessible to reaction. Acid cleavable linkers include, but are not limited to, bismaleimideoxy propane; and adipic acid dihydrazide linkers (see, *e.g.*, Fattom *et al.* (1992) *Infection & Immun.* 60:584-589) and acid labile transferrin conjugates that contain a sufficient portion of transferrin to permit entry into the intracellular transferrin cycling pathway (see, *e.g.*, Welhöner *et al.* (1991) *J. Biol. Chem.* 266:4309-4314).

Photocleavable linkers are linkers that are cleaved upon exposure to light (see, *e.g.*, Goldmacher *et al.* (1992) *Bioconj. Chem.* 3:104-107, which linkers are herein incorporated by reference), thereby releasing the targeted agent upon exposure to light. Photocleavable linkers that are cleaved upon exposure to light are known (see, *e.g.*, Hazum *et al.* (1981) in *Pept., Proc. Eur. Pept. Symp.*, 16th, Brunfeldt, K (Ed), pp. 105-110, which describes the use of a nitrobenzyl group as a photocleavable protective group for cysteine; Yen *et al.* (1989) *Makromol. Chem* 190:69-82, which describes water soluble photocleavable copolymers, including hydroxypropylmethacrylamide copolymer, glycine

-114-

copolymer, fluorescein copolymer and methylrhodamine copolymer; Goldmacher *et al.* (1992) *Bioconj. Chem.* 3:104-107, which describes a cross-linker and reagent that undergoes photolytic degradation upon exposure to near UV light (350 nm); and Senter *et al.* (1985) *Photochem. Photobiol* 42:231-237, which describes nitrobenzyloxycarbonyl chloride cross linking reagents that produce photocleavable linkages), thereby releasing the targeted agent upon exposure to light. Such linkers would have particular use in treating dermatological or ophthalmic conditions that can be exposed to light using fiber optics. After administration of the conjugate, the eye or skin or other body part can be exposed to light, resulting in release of the targeted moiety from the conjugate. Such photocleavable linkers are useful in connection with diagnostic protocols in which it is desirable to remove the targeting agent to permit rapid clearance from the body of the animal.

b) Other linkers for chemical conjugation

Other linkers, include trityl linkers, particularly, derivatized trityl groups to generate a genus of conjugates that provide for release of therapeutic agents at various degrees of acidity or alkalinity. The flexibility thus afforded by the ability to preselect the pH range at which the therapeutic agent is released allows selection of a linker based on the known physiological differences between tissues in need of delivery of a therapeutic agent (see, e.g., U.S. Patent No. 5,612,474). For example, the acidity of tumor tissues appears to be lower than that of normal tissues.

c) Peptide linkers

The linker moieties can be peptides. Peptide linkers can be employed in fusion proteins and also in chemically linked conjugates. The peptide typically has from about 2 to about 60 amino acid residues, for example from about 5 to about 40, or from about 10 to about 30 amino acid residues. The length selected depends upon factors, such as the use for which the linker is included.

Peptide linkers are advantageous when the targeting agent is proteinaceous. For example, the linker moiety can be a flexible spacer amino acid sequence, such as those known in single-chain antibody research. Examples of such known linker moieties include, but are not limited to,

-115-

peptides, such as $(\text{Gly}_m\text{Ser})_n$ and $(\text{Ser}_m\text{Gly})_n$, in which n is 1 to 6, including 1 to 4 and 2 to 4, and m is 1 to 6, including 1 to 4, and 2 to 4, enzyme cleavable linkers and others.

Additional linking moieties are described, for example, in Huston *et al.*,
5 *Proc. Natl. Acad. Sci. U.S.A.* 85:5879-5883, 1988; Whitlow, M., *et al.*, *Protein Engineering* 6:989-995, 1993; Newton *et al.*, *Biochemistry* 35:545-553, 1996; A. J. Cumber *et al.*, *Bioconj. Chem.* 3:397-401, 1992; Ladurner *et al.*, *J. Mol. Biol.* 273:330-337, 1997; and U.S. Patent. No. 4,894,443. In some
embodiments, several linkers can be included in order to take advantage of
10 desired properties of each linker.

3. Targeting agents

Any agent that facilitates detection, immobilization, or purification of the conjugate is contemplated for use herein. For chemical conjugates any moiety that has such properties is contemplated; for fusion proteins, the targeting agent
15 is a protein, peptide or fragment thereof that is sufficient to effects the targeting activity. Contemplated targeting agents include those that deliver the MTSP10 polypeptide or portion thereof to selected cells and tissues. Such agents include tumor specific monoclonal antibodies and portions thereof, growth factors, such as FGF, EGF, PDGF, VEGF, cytokines, including chemokines, and other such
20 agents.

4. Nucleic acids, plasmids and cells

Isolated nucleic acid fragments encoding fusion proteins are provided. The nucleic acid fragment that encodes the fusion protein includes: a) nucleic acid encoding a protease domain of an MTSP10 polypeptide; and b) nucleic acid
25 encoding a protein, peptide or effective fragment thereof that facilitates: i) affinity isolation or purification of the fusion protein; ii) attachment of the fusion protein to a surface; or iii) detection of the fusion protein. Generally, the nucleic acid is DNA.

Plasmids for replication and vectors for expression that contain the above
30 nucleic acid fragments are also provided. Cells containing the plasmids and vectors are also provided. The cells can be any suitable host including, but are not limited to, bacterial cells, yeast cells, fungal cells, plant cells, insect cell and

-116-

animal cells. The nucleic acids, plasmids, and cells containing the plasmids can be prepared according to methods known in the art including any described herein.

Also provided are methods for producing the above fusion proteins. An exemplary method includes the steps of growing, for example, culturing the cells so that they proliferate, cells containing a plasmid encoding the fusion protein under conditions whereby the fusion protein is expressed by the cell, and recovering the expressed fusion protein. Methods for expressing and recovering recombinant proteins are well known in the art (*See generally, Current Protocols in Molecular Biology* (1998) § 16, John Wiley & Sons, Inc.) and such methods can be used for expressing and recovering the expressed fusion proteins.

The recovered fusion proteins can be isolated or purified by methods known in the art such as centrifugation, filtration, chromatography, electrophoresis, immunoprecipitation, and other such methods, or by a combination thereof (*See generally, Current Protocols in Molecular Biology* (1998) § 10, John Wiley & Sons, Inc.). Generally the recovered fusion protein is isolated or purified through affinity binding between the protein or peptide fragment of the fusion protein and an affinity binding moiety. As discussed in the above sections regarding the construction of the fusion proteins, any affinity binding pairs can be constructed and used in the isolation or purification of the fusion proteins. For example, the affinity binding pairs can be protein binding sequences/protein, DNA binding sequences/DNA sequences, RNA binding sequences/RNA sequences, lipid binding sequences/lipid, polysaccharide binding sequences/polysaccharide, or metal binding sequences/metal.

5. Immobilization and supports or substrates therefor

In certain embodiments, where the targeting agents are designed for linkage to surfaces, the MTSP10 polypeptide can be attached by linkage such as ionic or covalent, non-covalent or other chemical interaction, to a surface of a support or matrix material. Immobilization can be effected directly or via a linker. The MTSP10 polypeptide can be immobilized on any suitable support, including, but are not limited to, silicon chips, and other supports described herein and known to those of skill in the art. A plurality of MTSP10 polypeptide

-117-

or protease domains thereof can be attached to a support, such as an array (*i.e.*, a pattern of two or more) of conjugates on the surface of a silicon chip or other chip for use in high throughput protocols and formats.

It is also noted that the domains of the MTSP10 polypeptide can be
5 linked directly to the surface or via a linker without a targeting agent linked thereto. Hence chips containing arrays of the domains of the MTSP10 polypeptide are also provided.

The matrix material or solid supports contemplated herein are generally any of the insoluble materials known to those of skill in the art to immobilize
10 ligands and other molecules, and are those that are used in many chemical syntheses and separations. Such supports are used, for example, in affinity chromatography, in the immobilization of biologically active materials, and during chemical syntheses of biomolecules, including proteins, amino acids and other organic molecules and polymers. The preparation of and use of supports is well
15 known to those of skill in this art; there are many such materials and preparations thereof known. For example, naturally-occurring support materials, such as agarose and cellulose, can be isolated from their respective sources, and processed according to known protocols, and synthetic materials can be prepared in accord with known protocols.

20 The supports are typically insoluble materials that are solid, porous, deformable, or hard, and have any required structure and geometry, including, but not limited to: beads, pellets, disks, capillaries, hollow fibers, needles, solid fibers, random shapes, thin films and membranes. Thus, the item can be fabricated from the matrix material or combined with it, such as by coating all or
25 part of the surface or impregnating particles.

Typically, when the matrix is particulate, the particles are at least about 10-2000 μm , but can be smaller or larger, depending upon the selected application. Selection of the matrices is governed, at least in part, by their physical and chemical properties, such as solubility, functional groups,
30 mechanical stability, surface area swelling propensity, hydrophobic or hydrophilic properties and intended use.

RECTIFIED SHEET (RULE 91)

-118-

If necessary, the support matrix material can be treated to contain an appropriate reactive moiety. In some cases, the support matrix material already containing the reactive moiety can be obtained commercially. The support matrix material containing the reactive moiety can thereby serve as the matrix

5 support upon which molecules are linked. Materials containing reactive surface moieties such as amino silane linkages, hydroxyl linkages or carboxysilane linkages can be produced by well established surface chemistry techniques involving silanization reactions, or the like. Examples of these materials are those having surface silicon oxide moieties, covalently linked to gamma-amino-

10 propylsilane, and other organic moieties; N-[3-(triethoxysilyl)propyl]phthelamic acid; and bis-(2-hydroxyethyl)aminopropyltriethoxysilane. Exemplary of readily available materials containing amino group reactive functionalities, include, but are not limited to, para-aminophenyltriethoxysilane. Also derivatized polystyrenes and other such polymers are well known and readily available to

15 those of skill in this art (e.g., the Tentagel® Resins are available with a multitude of functional groups, and are sold by Rapp Polymere, Tubingen, Germany; see, U.S. Patent No. 4,908,405 and U.S. Patent No. 5,292,814; see, also Butz et al., *Peptide Res.*, 7:20-23 (1994); and Kleine et al., *Immunobiol.*, 190:53-66 (1994)).

20 These matrix materials include any material that can act as a support matrix for attachment of the molecules of interest. Such materials are known to those of skill in this art, and include those that are used as a support matrix. These materials include, but are not limited to, inorganics, natural polymers, and synthetic polymers, including, but are not limited to: cellulose, cellulose

25 derivatives, acrylic resins, glass, silica gels, polystyrene, gelatin, polyvinyl pyrrolidone, co-polymers of vinyl and acrylamide, polystyrene cross-linked with divinylbenzene and others (see, Merrifield, *Biochemistry*, 3:1385-1390 (1964)), polyacrylamides, latex gels, polystyrene, dextran, polyacrylamides, rubber, silicon, plastics, nitrocellulose, celluloses, natural sponges. Of particular interest

30 herein, are highly porous glasses (see, e.g., U.S. Patent No. 4,244,721) and others prepared by mixing a borosilicate, alcohol and water.

- Synthetic supports include, but are not limited to: acrylamides, dextran-derivatives and dextran co-polymers, agarose-polyacrylamide blends, other polymers and co-polymers with various functional groups, methacrylate derivatives and co-polymers, polystyrene and polystyrene copolymers (*see, e.g.*,
- 5 Merrifield, *Biochemistry*, 3:1385-1390 (1964); Berg et al., in *Innovation Perspect. Solid Phase Synth. Collect. Pap.*, Int. Symp., 1st, Epton, Roger (Ed), pp. 453-459 (1990); Berg et al., *Pept., Proc. Eur. Pept. Symp.*, 20th, Jung, G. et al. (Eds), pp. 196-198 (1989); Berg et al., *J. Am. Chem. Soc.*, 111:8024-8026 (1989); Kent et al., *Isr. J. Chem.*, 17:243-247 (1979); Kent et
- 10 al., *J. Org. Chem.*, 43:2845-2852 (1978); Mitchell et al., *Tetrahedron Lett.*, 42:3795-3798 (1976); U.S. Patent No. 4,507,230; U.S. Patent No. 4,006,117; and U.S. Patent No. 5,389,449). Such materials include those made from polymers and co-polymers such as polyvinylalcohols, acrylates and acrylic acids such as polyethylene-co-acrylic acid, polyethylene-co-methacrylic acid, polyethy-
- 15 lene-co-ethylacrylate, polyethylene-co-methyl acrylate, polypropylene-co-acrylic acid, polypropylene-co-methyl-acrylic acid, polypropylene-co-ethylacrylate, polypropylene-co-methyl acrylate, polyethylene-co-vinyl acetate, polypropylene-co-vinyl acetate, and those containing acid anhydride groups such as polyethylene-co-maleic anhydride and polypropylene-co-maleic anhydride.
- 20 Liposomes have also been used as solid supports for affinity purifications (Powell et al. *Biotechnol. Bioeng.*, 33:173 (1989)).

- Numerous methods have been developed for the immobilization of proteins and other biomolecules onto solid or liquid supports (*see, e.g.*, Mosbach, *Methods in Enzymology*, 44 (1976); Weetall, *Immobilized Enzymes, Antigens, Antibodies, and Peptides*, (1975); Kennedy et al., *Solid Phase Biochemistry, Analytical and Synthetic Aspects*, Scouten, ed., pp. 253-391
- 25 (1983); *see, generally*, Affinity Techniques. Enzyme Purification: *Part B. Methods in Enzymology*, Vol. 34, ed. W. B. Jakoby, M. Wilchek, Acad. Press, N.Y. (1974); and Immobilized Biochemicals and Affinity Chromatography,
- 30 *Advances in Experimental Medicine and Biology*, vol. 42, ed. R. Dunlap, Plenum Press, N.Y. (1974)).

-120-

Among the most commonly used methods are absorption and adsorption or covalent binding to the support, either directly or via a linker, such as the numerous disulfide linkages, thioether bonds, hindered disulfide bonds, and covalent bonds between free reactive groups, such as amine and thiol groups, known to those of skill in art (see, e.g., the PIERCE CATALOG, ImmunoTechnology Catalog & Handbook, 1992-1993, which describes the preparation of and use of such reagents and provides a commercial source for such reagents; Wong, *Chemistry of Protein Conjugation and Cross Linking*, CRC Press (1993); see also DeWitt et al., *Proc. Natl. Acad. Sci. U.S.A.*, 90:6909 (1993); Zuckermann et al., *J. Am. Chem. Soc.*, 114:10646 (1992); Kurth et al., *J. Am. Chem. Soc.*, 116:2661 (1994); Ellman et al., *Proc. Natl. Acad. Sci. U.S.A.*, 91:4708 (1994); Sucholeiki, *Tetrahedron Lttrs.*, 35:7307 (1994); Su-Sun Wang, *J. Org. Chem.*, 41:3258 (1976); Padwa et al., *J. Org. Chem.*, 41:3550 (1971); and Vedejs et al., *J. Org. Chem.*, 49:575 (1984), which describe photosensitive linkers).

To effect immobilization, a composition containing the protein or other biomolecule is contacted with a support material such as alumina, carbon, an ion-exchange resin, cellulose, glass or a ceramic. Fluorocarbon polymers have been used as supports to which biomolecules have been attached by adsorption (see, U.S. Patent No. 3,843,443; Published International PCT Application WO/86 03840).

J. Prognosis and diagnosis

MTSP10 polypeptide proteins, domains, analogs, and derivatives thereof, and encoding nucleic acids (and sequences complementary thereto), and anti-MTSP10 polypeptide antibodies, can be used in diagnostics, particularly diagnosis of lung, head and neck, such as esophageal tumors, prostate, colon, ovary, cervix, breast and pancreas cancers. Such molecules can be used in assays, such as immunoassays, to detect, prognose, diagnose, or monitor various conditions, diseases, and disorders affecting MTSP10 polypeptide expression, or monitor the treatment thereof. For purposes herein, the presence of MTSP10s in body fluids or tumor tissues are of particular interest.

-121-

In particular, such an immunoassay is carried out by a method including contacting a sample derived from a patient with an anti-MTSP10 polypeptide antibody under conditions such that specific binding can occur, and detecting or measuring the amount of any specific binding by the antibody. Such binding of
5 antibody, in tissue sections, can be used to detect aberrant MTSP10 polypeptide localization or aberrant (*e.g.*, increased, decreased or absent) levels of MTSP10 polypeptide. In a specific embodiment, antibody to an MTSP10 polypeptide can be used to assay in a patient tissue or body fluid, such as serum, sample for the presence of MTSP10 polypeptide where an aberrant level of MTSP10
10 polypeptide is an indication of a diseased condition.

The immunoassays which can be used include but are not limited to competitive and non-competitive assay systems using techniques such as western blots, radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoprecipitation assays, precipitin
15 reactions, gel diffusion precipitin reactions, immunodiffusion assays, agglutination assays, complement-fixation assays, immunoradiometric assays, fluorescent immunoassays and protein A immunoassays.

MTSP10 polypeptide genes and related nucleic acid sequences and subsequences, including complementary sequences, also can be used in
20 hybridization assays. MTSP10 polypeptide nucleic acid sequences, or subsequences thereof containing about at least 8 nucleotides, generally 14 or 16 or 30 or more, generally less than 1000 or up to 100, contiguous nucleotides can be used as hybridization probes. Hybridization assays can be used to detect, prognose, diagnose, or monitor conditions, disorders, or disease states
25 associated with aberrant changes in MTSP10 polypeptide expression and/or activity as described herein. In particular, such a hybridization assay is carried out by a method by contacting a sample containing nucleic acid with a nucleic acid probe capable of hybridizing to MTSP10 polypeptide encoding DNA or RNA, under conditions such that hybridization can occur, and detecting or measuring
30 any resulting hybridization.

In a specific embodiment, a method of diagnosing a disease or disorder characterized by detecting an aberrant level of an MTSP10 polypeptide in a

RECTIFIED SHEET (RULE 91)

-122-

subject is provided herein by measuring the level of the DNA, RNA, protein or functional activity of the MTSP10 polypeptide in a sample derived from the subject, wherein an increase or decrease in the level of the DNA, RNA, protein or functional activity of the MTSP10 polypeptide, relative to the level of the DNA,
5 RNA, protein or functional activity found in an analogous sample not having the disease or disorder indicates the presence of the disease or disorder in the subject.

Kits for diagnostic use are also provided, that contain in one or more containers an anti-MTSP10 polypeptide antibody, and, optionally, a labeled
10 binding partner to the antibody. Alternatively, the anti-MTSP10 polypeptide antibody can be labeled (with a detectable marker, *e.g.*, a chemiluminescent, enzymatic, fluorescent, or radioactive moiety). A kit is also provided that includes in one or more containers a nucleic acid probe capable of hybridizing to the MTSP10 polypeptide-encoding nucleic acid. In a specific embodiment, a kit
15 can include in one or more containers a pair of primers (*e.g.*, each in the size range of 6-30 nucleotides) that are capable of priming amplification [*e.g.*, by polymerase chain reaction (see *e.g.*, Innis et al., 1990, PCR Protocols, Academic Press, Inc., San Diego, CA), ligase chain reaction (see EP 320,308) use of Q β replicase, cyclic probe reaction, or other methods known in the art under
20 appropriate reaction conditions of at least a portion of an MTSP10 polypeptide-encoding nucleic acid. A kit can optionally further include in a container a predetermined amount of a purified MTSP10 polypeptide or nucleic acid, *e.g.*, for use as a standard or control.

K. Pharmaceutical compositions and modes of administration

25 **1. Components of the compositions**

Pharmaceutical compositions containing the identified compounds that modulate the activity of an MTSP10 polypeptide are provided herein. Also provided are combinations of a compound that modulates the activity of an MTSP10 polypeptide and another treatment or compound for treatment of a
30 neoplastic disorder, such as a chemotherapeutic compound.

-123-

The MTSP10 polypeptide modulator and the anti-tumor agent can be packaged as separate compositions for administration together or sequentially or intermittently. Alternatively, they can be provided as a single composition for administration or as two compositions for administration
5 as a single composition. The combinations can be packaged as kits.

a. MTSP10 polypeptide inhibitors

Any MTSP10 polypeptide inhibitors, including those described herein when used alone or in combination with other compounds, that can alleviate, reduce, ameliorate, prevent, or place or maintain in a state of remission of
10 clinical symptoms or diagnostic markers associated with neoplastic diseases, including undesired and/or uncontrolled angiogenesis, can be used in the present combinations.

In one embodiment, the MTSP10 polypeptide inhibitor is an antibody or fragment thereof that specifically reacts with an MTSP10 polypeptide or the
15 protease domain thereof, an inhibitor of the MTSP10 polypeptide production, an inhibitor of MTSP10 polypeptide membrane-localization, or any inhibitor of the expression of or, especially, the activity of an MTSP10 polypeptide.

b. Anti-angiogenic agents and anti-tumor agents

Any anti-angiogenic agents and anti-tumor agents, including those
20 described herein, when used alone or in combination with other compounds, that can alleviate, reduce, ameliorate, prevent, or place or maintain in a state of remission of clinical symptoms or diagnostic markers associated with undesired and/or uncontrolled angiogenesis and/or tumor growth and metastasis, particularly solid neoplasms, vascular malformations and cardiovascular
25 disorders, chronic inflammatory diseases and aberrant wound repairs, circulatory disorders, crest syndromes, dermatological disorders, or ocular disorders, can be used in the combinations. Also contemplated are anti-tumor agents for use in combination with an inhibitor of an MTSP10 polypeptide.

-124-

c. Anti-tumor agents and anti-angiogenic agents

The compounds identified by the methods provided herein or provided herein can be used in combination with anti-tumor agents and/or anti-angiogenesis agents.

5 2. Formulations and route of administration

The compounds herein and agents can be formulated as pharmaceutical compositions, typically for single dosage administration. The concentrations of the compounds in the formulations are effective for delivery of an amount, upon administration, that is effective for the intended treatment. Typically, the
10 compositions are formulated for single dosage administration. To formulate a composition, the weight fraction of a compound or mixture thereof is dissolved, suspended, dispersed or otherwise mixed in a selected vehicle at an effective concentration such that the treated condition is relieved or ameliorated. Pharmaceutical carriers or vehicles suitable for administration of the compounds
15 provided herein include any such carriers known to those skilled in the art to be suitable for the particular mode of administration.

In addition, the compounds can be formulated as the sole pharmaceutically active ingredient in the composition or can be combined with other active ingredients. Liposomal suspensions, including tissue-targeted
20 liposomes, can also be suitable as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art. For example, liposome formulations can be prepared as described in U.S. Patent No. 4,522,811.

The active compound is included in the pharmaceutically acceptable
25 carrier in an amount sufficient to exert a therapeutically useful effect in the absence of undesirable side effects on the patient treated. The therapeutically effective concentration can be determined empirically by testing the compounds in known in vitro and in vivo systems, such as the assays provided herein.

The concentration of active compound in the drug composition depends
30 on absorption, inactivation and excretion rates of the active compound, the physicochemical characteristics of the compound, the dosage schedule, and amount administered as well as other factors known to those of skill in the art.

-125-

Typically a therapeutically effective dosage is contemplated. The amounts administered can be on the order of 0.001 to 1 mg/ml, including about 0.005-0.05 mg/ml and about 0.01 mg/ml, of blood volume. Pharmaceutical dosage unit forms are prepared to provide from about 1 mg to about 1000 mg, including from about 10 to about 500 mg, and including about 25-75 mg of the essential active ingredient or a combination of essential ingredients per dosage unit form. The precise dosage can be empirically determined.

The active ingredient can be administered at once, or can be divided into a number of smaller doses to be administered at intervals of time. It is understood that the precise dosage and duration of treatment is a function of the disease being treated and can be determined empirically using known testing protocols or by extrapolation from in vivo or in vitro test data. It is to be noted that concentrations and dosage values can also vary with the severity of the condition to be alleviated. It is to be further understood that for any particular subject, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions, and that the concentration ranges set forth herein are exemplary only and are not intended to limit the scope or use of the claimed compositions and combinations containing them.

Pharmaceutically acceptable derivatives include acids, salts, esters, hydrates, solvates and prodrug forms. The derivative is typically selected such that its pharmacokinetic properties are superior to the corresponding neutral compound.

Thus, effective concentrations or amounts of one or more of the compounds provided herein or pharmaceutically acceptable derivatives thereof are mixed with a suitable pharmaceutical carrier or vehicle for systemic, topical or local administration to form pharmaceutical compositions. Compounds are included in an amount effective for ameliorating or treating the disorder for which treatment is contemplated. The concentration of active compound in the composition depends on absorption, inactivation, excretion rates of the active compound, the dosage schedule, amount administered, particular formulation as well as other factors known to those of skill in the art.

-126-

Solutions or suspensions used for parenteral, intradermal, subcutaneous, or topical application can include any of the following components: a sterile diluent, such as water for injection, saline solution, fixed oil, polyethylene glycol, glycerine, propylene glycol or other synthetic solvent; antimicrobial agents, such as benzyl alcohol and methyl parabens; antioxidants, such as ascorbic acid and sodium bisulfite; chelating agents, such as ethylenediaminetetraacetic acid (EDTA); buffers, such as acetates, citrates and phosphates; and agents for the adjustment of tonicity such as sodium chloride or dextrose. Parenteral preparations can be enclosed in ampules, disposable syringes or single or multiple dose vials made of glass, plastic or other suitable material.

In instances in which the compounds exhibit insufficient solubility, methods for solubilizing compounds can be used. Such methods are known to those of skill in this art, and include, but are not limited to, using cosolvents, such as dimethylsulfoxide (DMSO), using surfactants, such as Tween®, or dissolution in aqueous sodium bicarbonate. Derivatives of the compounds, such as prodrugs of the compounds can also be used in formulating effective pharmaceutical compositions. For ophthalmic indications, the compositions are formulated in an ophthalmically acceptable carrier. For the ophthalmic uses herein, local administration, either by topical administration or by injection are contemplated. Time release formulations are also desirable. Typically, the compositions are formulated for single dosage administration, so that a single dose administers an effective amount.

Upon mixing or addition of the compound with the vehicle, the resulting mixture can be a solution, suspension, emulsion or other composition. The form of the resulting mixture depends upon a number of factors, including the intended mode of administration and the solubility of the compound in the selected carrier or vehicle. If necessary, pharmaceutically acceptable salts or other derivatives of the compounds are prepared.

The compound is included in the pharmaceutically acceptable carrier in an amount sufficient to exert a therapeutically useful effect in the absence of undesirable side effects on the patient treated. It is understood that number and degree of side effects depends upon the condition for which the compounds are

-127-

administered. For example, certain toxic and undesirable side effects are tolerated when treating life-threatening illnesses that would not be tolerated when treating disorders of lesser consequence.

The compounds also can be mixed with other active materials, that do
5 not impair the desired action, or with materials that supplement the desired action known to those of skill in the art. The formulations of the compounds and agents for use herein include those suitable for oral, rectal, topical, inhalational, buccal (*e.g.*, sublingual), parenteral (*e.g.*, subcutaneous, intramuscular, intradermal, or intravenous), transdermal administration or any
10 route. The most suitable route in any given case depends on the nature and severity of the condition being treated and on the nature of the particular active compound which is being used. The formulations are provided for administration to humans and animals in unit dosage forms, such as tablets, capsules, pills, powders, granules, sterile parenteral solutions or suspensions, and oral solutions
15 or suspensions, and oil-water emulsions containing suitable quantities of the compounds or pharmaceutically acceptable derivatives thereof. The pharmaceutically therapeutically active compounds and derivatives thereof are typically formulated and administered in unit-dosage forms or multiple-dosage forms. Unit-dose forms as used herein refers to physically discrete units suitable
20 for human and animal subjects and packaged individually as is known in the art. Each unit-dose contains a predetermined quantity of the therapeutically active compound sufficient to produce the desired therapeutic effect, in association with the required pharmaceutical carrier, vehicle or diluent. Examples of unit-dose forms include ampoules and syringes and individually packaged tablets
25 or capsules. Unit-dose forms can be administered in fractions or multiples thereof. A multiple-dose form is a plurality of identical unit-dosage forms packaged in a single container to be administered in segregated unit-dose form. Examples of multiple-dose forms include vials, bottles of tablets or capsules or bottles of pints or gallons. Hence, multiple dose form is a multiple of unit-doses
30 which are not segregated in packaging.

The composition can contain along with the active ingredient: a diluent such as lactose, sucrose, dicalcium phosphate, or carboxymethylcellulose; a

-128-

lubricant, such as magnesium stearate, calcium stearate and talc; and a binder such as starch, natural gums, such as gum acacia, gelatin, glucose, molasses, polyvinylpyrrolidone, celluloses and derivatives thereof, povidone, crospovidones and other such binders known to those of skill in the art. Liquid

5 pharmaceutically administrable compositions can, for example, be prepared by dissolving, dispersing, or otherwise mixing an active compound as defined above and optional pharmaceutical adjuvants in a carrier, such as, for example, water, saline, aqueous dextrose, glycerol, glycols, ethanol, and the like, to thereby form a solution or suspension. If desired, the pharmaceutical composition to be
10 administered can also contain minor amounts of nontoxic auxiliary substances such as wetting agents, emulsifying agents, or solubilizing agents, pH buffering agents and the like, for example, acetate, sodium citrate, cyclodextrine derivatives, sorbitan monolaurate, triethanolamine sodium acetate, triethanolamine oleate, and other such agents. Methods of preparing such
15 dosage forms are known, or will be apparent, to those skilled in this art (see, *e.g.*, Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, Pa., 15th Edition, 1975). The composition or formulation to be administered contains a quantity of the active compound in an amount sufficient to alleviate the symptoms of the treated subject.

20 Dosage forms or compositions containing active ingredient in the range of 0.005% to 100% with the balance made up from non-toxic carrier can be prepared. For oral administration, the pharmaceutical compositions can take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (*e.g.*,
25 pregelatinized maize starch, polyvinyl pyrrolidone or hydroxypropyl methylcellulose); fillers (*e.g.*, lactose, microcrystalline cellulose or calcium hydrogen phosphate); lubricants (*e.g.*, magnesium stearate, talc or silica); disintegrants (*e.g.*, potato starch or sodium starch glycolate); or wetting agents (*e.g.*, sodium lauryl sulphate). The tablets can be coated by methods well-
30 known in the art.

The pharmaceutical preparation can also be in liquid form, for example, solutions, syrups or suspensions, or can be presented as a drug product for

RECTIFIED SHEET (RULE 91)

-129-

reconstitution with water or other suitable vehicle before use. Such liquid preparations can be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (*e.g.*, sorbitol syrup, cellulose derivatives or hydrogenated edible fats); emulsifying agents (*e.g.*, lecithin or acacia); non-aqueous vehicles (*e.g.*, almond oil, oily esters, or fractionated vegetable oils); and preservatives (*e.g.*, methyl or propyl-p-hydroxybenzoates or sorbic acid).

Formulations suitable for rectal administration can be presented as unit dose suppositories. These can be prepared by admixing the active compound with one or more conventional solid carriers, for example, cocoa butter, and then shaping the resulting mixture.

Formulations suitable for topical application to the skin or to the eye generally are formulated as an ointment, cream, lotion, paste, gel, spray, aerosol and oil. Carriers which can be used include vaseline, lanoline, polyethylene glycols, alcohols, and combinations of two or more thereof. The topical formulations can further advantageously contain 0.05 to 15 percent by weight of thickeners selected from among hydroxypropyl methyl cellulose, methyl cellulose, polyvinylpyrrolidone, polyvinyl alcohol, poly (alkylene glycols), poly/hydroxyalkyl, (meth)acrylates or poly(meth)acrylamides. A topical formulation is often applied by instillation or as an ointment into the conjunctival sac. It also can be used for irrigation or lubrication of the eye, facial sinuses, and external auditory meatus. It can also be injected into the anterior eye chamber and other places. The topical formulations in the liquid state can be also present in a hydrophilic three-dimensional polymer matrix in the form of a strip, contact lens, and the like from which the active components are released.

For administration by inhalation, the compounds for use herein can be delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, *e.g.*, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol, the dosage unit can be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, *e.g.*, gelatin, for use in an inhaler or insufflator can be formulated

-130-

containing a powder mix of the compound and a suitable powder base such as lactose or starch.

Formulations suitable for buccal (sublingual) administration include, for example, lozenges containing the active compound in a flavored base, usually sucrose and acacia or tragacanth; and pastilles containing the compound in an inert base such as gelatin and glycerin or sucrose and acacia.

The compounds can be formulated for parenteral administration by injection, *e.g.*, by bolus injection or continuous infusion. Formulations for injection can be presented in unit dosage form, *e.g.*, in ampules or in multi-dose containers, with an added preservative. The compositions can be suspensions, solutions or emulsions in oily or aqueous vehicles, and can contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient can be in powder form for reconstitution with a suitable vehicle, *e.g.*, sterile pyrogen-free water or other solvents, before use.

Formulations suitable for transdermal administration can be presented as discrete patches adapted to remain in intimate contact with the epidermis of the recipient for a prolonged period of time. Such patches suitably contain the active compound as an optionally buffered aqueous solution of, for example, 0.1 to 0.2 M concentration with respect to the active compound. Formulations suitable for transdermal administration can also be delivered by iontophoresis (*see, e.g.*, Pharmaceutical Research 3 (6), 318 (1986)) and typically take the form of an optionally buffered aqueous solution of the active compound.

The pharmaceutical compositions can also be administered by controlled release means and/or delivery devices (*see, e.g.*, in U.S. Patent Nos. 3,536,809; 3,598,123; 3,630,200; 3,845,770; 3,847,770; 3,916,899; 4,008,719; 4,687,610; 4,769,027; 5,059,595; 5,073,543; 5,120,548; 5,354,566; 5,591,767; 5,639,476; 5,674,533 and 5,733,566).

Desirable blood levels can be maintained by a continuous infusion of the active agent as ascertained by plasma levels. It should be noted that the attending physician would know how to and when to terminate, interrupt or adjust therapy to lower dosage due to toxicity, or bone marrow, liver or kidney dysfunctions. Conversely, the attending physician would also know how to and

-131-

when to adjust treatment to higher levels if the clinical response is not adequate (precluding toxic side effects).

The efficacy and/or toxicity of the MTSP10 polypeptide inhibitor(s), alone or in combination with other agents also can be assessed by the methods known
5 in the art (See generally, O'Reilly, *Investigational New Drugs*, 15:5-13 (1997)).

The active compounds or pharmaceutically acceptable derivatives can be prepared with carriers that protect the compound against rapid elimination from the body, such as time release formulations or coatings.

Kits containing the compositions and/or the combinations with
10 instructions for administration thereof are provided. The kit can further include a needle or syringe, typically packaged in sterile form, for injecting the complex, and/or a packaged alcohol pad. Instructions are optionally included for administration of the active agent by a clinician or by the patient.

Finally, the compounds or MTSP10 polypeptides or protease domains
15 thereof or compositions containing any of the preceding agents can be packaged as articles of manufacture containing packaging material, a compound or suitable derivative thereof provided herein, which is effective for treatment of a diseases or disorders contemplated herein, within the packaging material, and a label that indicates that the compound or a suitable derivative thereof is for treating the
20 diseases or disorders contemplated herein. The label can optionally include the disorders for which the therapy is warranted.

L. Methods of treatment

The compounds identified by the methods herein are used for treating or preventing neoplastic diseases in an animal, particularly a mammal, including a
25 human, is provided herein. In one embodiment, the method includes administering to a mammal an effective amount of an inhibitor of an MTSP10 polypeptide, whereby the disease or disorder is treated or prevented.

In an embodiment, the MTSP10 polypeptide inhibitor used in the treatment or prevention is administered with a pharmaceutically acceptable
30 carrier or excipient. The mammal treated can be a human. The inhibitors provided herein are those identified by the screening assays. In addition,

-132-

antibodies and antisense nucleic acids or double-stranded RNA (dsRNA), such as RNAi, are contemplated.

The treatment or prevention method can further include administering an anti-angiogenic treatment or agent or anti-tumor agent simultaneously with, prior to or subsequent to the MTSP10 polypeptide inhibitor, which can be any compound identified that inhibits the activity of an MTSP10 polypeptide. Such compounds include small molecule modulators, an antibody or a fragment or derivative thereof containing a binding region thereof against the MTSP10 polypeptide, an antisense nucleic acid or double-stranded RNA (dsRNA), such as RNAi, encoding an a portion of the MTSP10 polypeptide or complementary to thereto, and a nucleic acid containing at least a portion of a gene encoding the MTSP10 polypeptide into which a heterologous nucleotide sequence has been inserted such that the heterologous sequence inactivates the biological activity of at least a portion of the gene encoding the MTSP10 polypeptide, in which the portion of the gene encoding the MTSP10 polypeptide flanks the heterologous sequence to promote homologous recombination with a genomic gene encoding the MTSP10 polypeptide. In addition, such molecules are generally less than about 1000 nt long.

1. Antisense treatment

In a specific embodiment, as described hereinabove, MTSP10 polypeptide function is reduced or inhibited by MTSP10 polypeptide antisense nucleic acids, to treat or prevent neoplastic disease. The therapeutic or prophylactic use of nucleic acids of at least six nucleotides, generally up to about 150 nucleotides, that are antisense to a gene or cDNA encoding MTSP10 polypeptide or a portion thereof is provided. An MTSP10 polypeptide "antisense" nucleic acid as used herein refers to a nucleic acid capable of hybridizing to a portion of an MTSP10 polypeptide RNA (generally mRNA) by virtue of some sequence complementarity, and generally under high stringency conditions. The antisense nucleic acid can be complementary to a coding and/or noncoding region of an MTSP10 polypeptide mRNA. Such antisense nucleic acids have utility as therapeutics that reduce or inhibit MTSP10 polypeptide function, and can be used in the treatment or prevention of disorders as described *supra*.

-133-

The MTSP10 polypeptide antisense nucleic acids are of at least six nucleotides and are generally oligonucleotides (ranging from 6 to about 150 nucleotides including 6 to 50 nucleotides). The antisense molecule can be complementary to all or a portion of the protease domain. For example, the
5 oligonucleotide is at least 10 nucleotides, at least 15 nucleotides, at least 100 nucleotides, or at least 125 nucleotides. The oligonucleotides can be DNA or RNA or chimeric mixtures or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone. The oligonucleotide can include
10 other appending groups such as peptides, or agents facilitating transport across the cell membrane (see, *e.g.*, Letsinger et al., *Proc. Natl. Acad. Sci. U.S.A.* 86:6553-6556 (1989); Lemaitre et al., *Proc. Natl. Acad. Sci. U.S.A.* 84:648-652 (1987); PCT Publication No. WO 88/09810, published December 15, 1988) or blood-brain barrier (see, *e.g.*, PCT Publication No. WO 89/10134, published April
15 25, 1988), hybridization-triggered cleavage agents (see, *e.g.*, Krol et al., *BioTechniques* 6:958-976 (1988)) or intercalating agents (see, *e.g.*, Zon, *Pharm. Res.* 5:539-549 (1988)).

The MTSP10 polypeptide antisense nucleic acid generally is an oligonucleotide, typically single-stranded DNA or RNA or an analog thereof or
20 mixtures thereof. For example, the oligonucleotide includes a sequence antisense to a portion of a nucleic acid that encodes a human MTSP10 polypeptide. The oligonucleotide can be modified at any position on its structure with substituents generally known in the art.

The MTSP10 polypeptide antisense oligonucleotide can include at least
25 one modified base moiety which is selected from the group including, but not limited to, 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine,
30 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-

RECTIFIED SHEET (RULE 91)

-134-

2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, 5 uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine.

In another embodiment, the oligonucleotide includes at least one modified sugar moiety selected from the group including but not limited to arabinose, 2-fluoroarabinose, xylulose, and hexose. The oligonucleotide can include at least one modified phosphate backbone selected from a phosphorothioate, a phosphorodithioate, a phosphoramidothioate, a phosphoramidate, a phosphordiamidate, a methylphosphonate, an alkyl phosphotriester, and a formacetal or analog thereof.

The oligonucleotide can be an α -anomeric oligonucleotide. An α -anomeric oligonucleotide forms specific double-stranded hybrids with complementary RNA in which the strands run parallel to each other (Gautier et al., *Nucl. Acids Res.* 15:6625-6641 (1987)).

The oligonucleotide can be conjugated to another molecule, such as, but are not limited to, a peptide, hybridization triggered cross-linking agent, transport agent or a hybridization-triggered cleavage agent. The oligonucleotides can be synthesized by standard methods known in the art, *e.g.* by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). As examples, phosphorothioate oligonucleotides can be synthesized by the method of Stein et al. (*Nucl. Acids Res.* 16:3209 (1988)), methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin et al., *Proc. Natl. Acad. Sci. U.S.A.* 85:7448-7451 (1988)), etc.

In a specific embodiment, the MTSP10 polypeptide antisense oligonucleotide includes catalytic RNA or a ribozyme (see, *e.g.*, PCT International Publication WO 90/11364, published October 4, 1990; Sarver et al., *Science* 247:1222-1225 (1990)). In another embodiment, the oligonucleotide is a 2'-O-methylribonucleotide (Inoue et al., *Nucl. Acids Res.* 15:6131-6148 (1987)), or a

-135-

chimeric RNA-DNA analogue (Inoue et al., *FEBS Lett.* 215:327-330 (1987)). Alternatively, the oligonucleotide can be double-stranded RNA (dsRNA) such as RNAi.

In an alternative embodiment, the MTSP10 polypeptide antisense nucleic acid is produced intracellularly by transcription from an exogenous sequence. For example, a vector can be introduced *in vivo* such that it is taken up by a cell, within which cell the vector or a portion thereof is transcribed, producing an antisense nucleic acid (RNA). Such a vector would contain a sequence encoding the MTSP10 polypeptide antisense nucleic acid. Such a vector can remain episomal or become chromosomally integrated, as long as it can be transcribed to produce the desired antisense RNA. Such vectors can be constructed by recombinant DNA technology methods standard in the art. Vectors can be plasmid, viral, or others known in the art, used for replication and expression in mammalian cells. Expression of the sequence encoding the MTSP10 polypeptide antisense RNA can be by any promoter known in the art to act in mammalian, including human, cells. Such promoters can be inducible or constitutive. Such promoters include but are not limited to: the SV40 early promoter region (Bernoist and Chambon, *Nature* 290:304-310 (1981), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto et al., *Cell* 22:787-797 (1980), the herpes thymidine kinase promoter (Wagner et al., *Proc. Natl. Acad. Sci. U.S.A.* 78:1441-1445 (1981), the regulatory sequences of the metallothionein gene (Brinster et al., *Nature* 296:39-42 (1982), etc.

The antisense nucleic acids include sequence complementary to at least a portion of an RNA transcript of an MTSP10 polypeptide gene, including a human MTSP10 polypeptide gene. Absolute complementarity is not required.

The amount of MTSP10 polypeptide antisense nucleic acid that is effective in the treatment or prevention of neoplastic disease depends on the nature of the disease, and can be determined empirically by standard clinical techniques. Where possible, it is desirable to determine the antisense cytotoxicity in cells *in vitro*, and then in useful animal model systems prior to testing and use in humans.

-136-

2. RNA interference

RNA interference (RNAi) (see, *e.g.* Chuang *et al.* (2000) *Proc. Natl. Acad. Sci. U.S.A.* 97:4985) can be employed to inhibit the expression of a gene encoding an MTSP10. Interfering RNA (RNAi) fragments, particularly double-stranded (ds) RNAi, can be used to generate loss-of-MTSP10 function. Methods relating to the use of RNAi to silence genes in organisms including, mammals, C. elegans, Drosophila and plants, and humans are known (see, *e.g.*, Fire *et al.* (1998) *Nature* 391:806-811; Fire (1999) *Trends Genet.* 15:358-363; Sharp (2001) *Genes Dev.* 15:485-490; Hammond, *et al.* (2001) *Nature Rev. Genet.* 2:110-1119; Tuschl (2001) *Chem. Biochem.* 2:239-245; Hamilton *et al.* (1999) *Science* 286:950-952; Hammond *et al.* (2000) *Nature* 404:293-296; Zamore *et al.* (2000) *Cell* 101:25-33; Bernstein *et al.* (2001) *Nature* 409: 363-366; Elbashir *et al.* (2001) *Genes Dev.* 15:188-200; Elbashir *et al.* (2001) *Nature* 411:494-498; International PCT application No. WO 01/29058; International PCT application No. WO 99/32619).

Double-stranded RNA (dsRNA)-expressing constructs are introduced into a host, such as an animal or plant using, a replicable vector that remains episomal or integrates into the genome. By selecting appropriate sequences, expression of dsRNA can interfere with accumulation of endogenous mRNA encoding an MTSP10. RNAi also can be used to inhibit expression *in vitro*. Regions include at least about 21 (or 21) nucleotides that are selective (*i.e.* unique) for MTSP10 are used to prepare the RNAi. Smaller fragments of about 21 nucleotides can be transformed directly (*i.e.*, *in vitro* or *in vivo*) into cells; larger RNAi dsRNA molecules are generally introduced using vectors that encode them. dsRNA molecules are at least about 21 bp long or longer, such as 50, 100, 150, 200 and longer. Methods, reagents and protocols for introducing nucleic acid molecules in to cells *in vitro* and *in vivo* are known to those of skill in the art.

-137-

3. Gene Therapy

In an exemplary embodiment, nucleic acids that include a sequence of nucleotides encoding an MTSP10 polypeptide or functional domains or derivative thereof, are administered to promote MTSP10 polypeptide function, by way of
5 gene therapy. Gene therapy refers to therapy performed by the administration of a nucleic acid to a subject. In this embodiment, the nucleic acid produces its encoded protein that mediates a therapeutic effect by promoting MTSP10 polypeptide function. Any of the methods for gene therapy available in the art can be used (see, Goldspiel et al., *Clinical Pharmacy* 12:488-505 (1993); Wu
10 and Wu, *Biotherapy* 3:87-95 (1991); Tolstoshev, *An. Rev. Pharmacol. Toxicol.* 32:573-596 (1993); Mulligan, *Science* 260:926-932 (1993); and Morgan and Anderson, *An. Rev. Biochem.* 62:191-217 (1993); *TIBTECH* 11(5):155-215 (1993).

For example, one therapeutic composition for gene therapy includes an
15 MTSP10 polypeptide-encoding nucleic acid that is part of an expression vector that expresses an MTSP10 polypeptide or domain, fragment or chimeric protein thereof in a suitable host. In particular, such a nucleic acid has a promoter operably linked to the MTSP10 polypeptide coding region, the promoter being inducible or constitutive, and, optionally, tissue-specific. In another particular
20 embodiment, a nucleic acid molecule is used in which the MTSP10 polypeptide coding sequences and any other desired sequences are flanked by regions that promote homologous recombination at a desired site in the genome, thus providing for intrachromosomal expression of the SP protein nucleic acid (Koller and Smithies, *Proc. Natl. Acad. Sci. USA* 86:8932-8935 (1989); Zijlstra et al.,
25 *Nature* 342:435-438 (1989)).

Delivery of the nucleic acid into a patient can be either direct, in which case the patient is directly exposed to the nucleic acid or nucleic acid-carrying vector, or indirect, in which case, cells are first transformed with the nucleic acid *in vitro*, then transplanted into the patient. These two approaches are known,
30 respectively, as *in vivo* or *ex vivo* gene therapy.

In a specific embodiment, the nucleic acid is directly administered *in vivo*, where it is expressed to produce the encoded product. This can be

-138-

accomplished by any of numerous methods known in the art, *e.g.*, by constructing it as part of an appropriate nucleic acid expression vector and administering it so that it becomes intracellular, *e.g.*, by infection using a defective or attenuated retroviral or other viral vector (see U.S. Patent No. 4,980,286), or by direct injection of naked DNA, or by use of microparticle bombardment (*e.g.*, a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, encapsulation in liposomes, microparticles, or microcapsules, or by administering it in linkage to a peptide which is known to enter the nucleus, by administering it in linkage to a ligand subject to receptor-mediated endocytosis (see *e.g.*, Wu and Wu, *J. Biol. Chem.* 262:4429-4432 (1987)) (which can be used to target cell types specifically expressing the receptors), etc. In another embodiment, a nucleic acid-ligand complex can be formed in which the ligand is a fusogenic viral peptide to disrupt endosomes, allowing the nucleic acid to avoid lysosomal degradation. In yet another embodiment, the nucleic acid can be targeted *in vivo* for cell specific uptake and expression, by targeting a specific receptor (see, *e.g.*, PCT Publications WO 92/06180 dated April 16, 1992 (Wu et al.); WO 92/22635 dated December 23, 1992 (Wilson et al.); WO92/20316 dated November 26, 1992 (Findeis et al.); WO93/14188 dated July 22, 1993 (Clarke et al.), WO 93/20221 dated October 14, 1993 (Young)). Alternatively, the nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination (Koller and Smithies, *Proc. Natl. Acad. Sci. USA* 86:8932-8935 (1989); Zijlstra et al., *Nature* 342:435-438 (1989)).

In a specific embodiment, a viral vector that contains the MTSP10 polypeptide nucleic acid is used. For example, a retroviral vector can be used (see Miller et al., *Meth. Enzymol.* 217:581-599 (1993)). These retroviral vectors have been modified to delete retroviral sequences that are not necessary for packaging of the viral genome and integration into host cell DNA. The MTSP10 polypeptide nucleic acid to be used in gene therapy is cloned into the vector, which facilitates delivery of the gene into a patient. More detail about retroviral vectors can be found in Boesen et al., *Biotherapy* 6:291-302 (1994), which describes the use of a retroviral vector to deliver the *mdr1* gene to hematopoietic

-139-

stem cells in order to make the stem cells more resistant to chemotherapy.

Other references illustrating the use of retroviral vectors in gene therapy are:

Clowes et al., *J. Clin. Invest.* 93:644-651 (1994); Kiem et al., *Blood* 83:1467-1473 (1994); Salmons and Gunzberg, *Human Gene Therapy* 4:129-141 (1993);

5 and Grossman and Wilson, *Curr. Opin. in Genetics and Devel.* 3:110-114 (1993).

Adenoviruses are other viral vectors that can be used in gene therapy.

Adenoviruses are especially attractive vehicles for delivering genes to respiratory epithelia. Adenoviruses naturally infect respiratory epithelia where they cause a

10 mild disease. Other targets for adenovirus-based delivery systems are liver, the central nervous system, endothelial cells, and muscle. Adenoviruses have the advantage of being capable of infecting non-dividing cells. Kozarsky and Wilson, *Current Opinion in Genetics and Development* 3:499-503 (1993) present a review of adenovirus-based gene therapy. Bout et al., *Human Gene Therapy*
15 5:3-10 (1994) demonstrated the use of adenovirus vectors to transfer genes to the respiratory epithelia of rhesus monkeys. Other instances of the use of adenoviruses in gene therapy can be found in Rosenfeld et al., *Science* 252:431-434 (1991); Rosenfeld et al., *Cell* 68:143-155 (1992); and Mastrangeli et al., *J. Clin. Invest.* 91:225-234 (1993).

20 Adeno-associated virus (AAV) has also been proposed for use in gene therapy (Walsh et al., *Proc. Soc. Exp. Biol. Med.* 204:289-300 (1993)).

Another approach to gene therapy involves transferring a gene to cells in tissue culture by such methods as electroporation, lipofection, calcium phosphate mediated transfection, or viral infection. Usually, the method of
25 transfer includes the transfer of a selectable marker to the cells. The cells are then placed under selection to isolate those cells that have taken up and are expressing the transferred gene. Those cells are then delivered to a patient.

In this embodiment, the nucleic acid is introduced into a cell prior to administration *in vivo* of the resulting recombinant cell. Such introduction can
30 be carried out by any method known in the art, including but not limited to transfection, electroporation, microinjection, infection with a viral or bacteriophage vector containing the nucleic acid sequences, cell fusion,

-140-

chromosome-mediated gene transfer, microcell-mediated gene transfer, spheroplast fusion, etc. Numerous techniques are known in the art for the introduction of foreign genes into cells (see *e.g.*, Loeffler and Behr, *Meth. Enzymol.* 217:599-618 (1993); Cohen et al., *Meth. Enzymol.* 217:618-644

5 (1993); Cline, *Pharmac. Ther.* 29:69-92 (1985)) and can be used, provided that the necessary developmental and physiological functions of the recipient cells are not disrupted. The technique should provide for the stable transfer of the nucleic acid to the cell, so that the nucleic acid is expressible by the cell and generally heritable and expressible by its cell progeny.

10 The resulting recombinant cells can be delivered to a patient by various methods known in the art. In an embodiment, epithelial cells are injected, *e.g.*, subcutaneously. In another embodiment, recombinant skin cells can be applied as a skin graft onto the patient. Recombinant blood cells (*e.g.*, hematopoietic stem or progenitor cells) can be administered intravenously. The amount of cells
15 envisioned for use depends on the desired effect, patient state, etc., and can be determined by one skilled in the art.

Cells into which a nucleic acid can be introduced for purposes of gene therapy encompass any desired, available cell type, and include but are not limited to epithelial cells, endothelial cells, keratinocytes, fibroblasts, muscle
20 cells, hepatocytes; blood cells such as T lymphocytes, B lymphocytes, monocytes, macrophages, neutrophils, eosinophils, megakaryocytes, granulocytes; various stem or progenitor cells, in particular hematopoietic stem or progenitor cells, *e.g.*, such as stem cells obtained from bone marrow, umbilical cord blood, peripheral blood, fetal liver, and other sources thereof.

25 For example, a cell used for gene therapy is autologous to the patient. In an embodiment in which recombinant cells are used in gene therapy, an MTSP10 polypeptide nucleic acid is introduced into the cells such that it is expressible by the cells or their progeny, and the recombinant cells are then administered *in vivo* for therapeutic effect. In a specific embodiment, stem or progenitor cells
30 are used. Any stem and/or progenitor cells which can be isolated and maintained *in vitro* can potentially be used in accordance with this embodiment. Such stem cells include but are not limited to hematopoietic stem cells (HSC),

-141-

stem cells of epithelial tissues such as the skin and the lining of the gut, embryonic heart muscle cells, liver stem cells (PCT Publication WO 94/08598, dated April 28, 1994), and neural stem cells (Stemple and Anderson, *Cell* 71:973-985 (1992)).

5 Epithelial stem cells (ESCs) or keratinocytes can be obtained from tissues such as the skin and the lining of the gut by known procedures (Rheinwald, *Meth. Cell Bio.* 21A:229 (1980)). In stratified epithelial tissue such as the skin, renewal occurs by mitosis of stem cells within the germinal layer, the layer closest to the basal lamina. Stem cells within the lining of the gut provide for a
10 rapid renewal rate of this tissue. ESCs or keratinocytes obtained from the skin or lining of the gut of a patient or donor can be grown in tissue culture (Rheinwald, *Meth. Cell Bio.* 21A:229 (1980); Pittelkow and Scott, *Cano Clinic Proc.* 61:771 (1986)). If the ESCs are provided by a donor, a method for
15 suppression of host versus graft reactivity (*e.g.*, irradiation, drug or antibody administration to promote moderate immunosuppression) also can be used.

 With respect to hematopoietic stem cells (HSC), any technique which provides for the isolation, propagation, and maintenance *in vitro* of HSC can be used in this embodiment. Techniques by which this can be accomplished include (a) the isolation and establishment of HSC cultures from bone marrow
20 cells isolated from the future host, or a donor, or (b) the use of previously established long-term HSC cultures, which can be allogeneic or xenogeneic. Non-autologous HSC generally are used with a method of suppressing transplantation immune reactions of the future host/patient. In a particular
25 embodiment, human bone marrow cells can be obtained from the posterior iliac crest by needle aspiration (see, *e.g.*, Kodo et al., *J. Clin. Invest.* 73:1377-1384 (1984)). For example, the HSCs can be made highly enriched or in substantially
pure form. This enrichment can be accomplished before, during, or after long-term culturing, and can be done by any techniques known in the art. Long-term
30 cultures of bone marrow cells can be established and maintained by using, for example, modified Dexter cell culture techniques (Dexter et al., *J. Cell Physiol.* 91:335 (1977) or Witlock-Witte culture techniques (Witlock and Witte, *Proc. Natl. Acad. Sci. USA* 79:3608-3612 (1982)).

-142-

In a specific embodiment, the nucleic acid to be introduced for purposes of gene therapy includes an inducible promoter operably linked to the coding region, such that expression of the nucleic acid is controllable by controlling the presence or absence of the appropriate inducer of transcription.

5 **3. Prodrugs**

A method for treating tumors is provided. The method is practiced by administering a prodrug that is cleaved at a specific site by an MTSP10 to release an active drug or precursor that can be converted to active drug *in vivo*. Upon contact with a cell that expresses MTSP10 activity, the prodrug is
10 converted into an active drug. The prodrug can be a conjugate that contains the active agent, such as an anti-tumor drug, such as a cytotoxic agent, or other therapeutic agent (TA), linked to a substrate for the targeted MTSP10, such that the drug or agent is inactive or unable to enter a cell, in the conjugate, but is activated upon cleavage. The prodrug, for example, can contain an oligopeptide,
15 typically a relatively short, less than about 10 amino acids peptide, that is proteolytically cleaved by the targeted MTSP10. Cytotoxic agents, include, but are not limited to, alkylating agents, antiproliferative agents and tubulin binding agents. Others include, vinca drugs, mitomycins, bleomycins and taxanes.

M. Animal models

20 Transgenic animal models and animals, such as rodents, including mice and rats, cows, chickens, pigs, goats, sheep, monkeys, including gorillas, and other primates, are provided herein. In particular, transgenic non-human animals that contain heterologous nucleic acid encoding an MTSP10 polypeptide or a transgenic animal in which expression of the polypeptide has been altered, such
25 as by replacing or modifying the promoter region or other regulatory region of the endogenous gene are provided. Such an animal can be produced by promoting recombination between endogenous nucleic acid and an exogenous MTSP10 gene that could be over-expressed or mis-expressed, such as by expression under a strong promoter, via homologous or other recombination
30 event.

-143-

Transgenic animals can be produced by introducing the nucleic acid using any known method of delivery, including, but not limited to, microinjection, lipofection and other modes of gene delivery into a germline cell or somatic cells, such as an embryonic stem cell. Typically the nucleic acid is introduced into a
5 cell, such as an embryonic stem cell (ES), followed by injecting the ES cells into a blastocyst, and implanting the blastocyst into a foster mother, which is followed by the birth of a transgenic animal. Generally, introduction of a heterologous nucleic acid molecule into a chromosome of the animal occurs by a recombination between the heterologous MTSP10-encoding nucleic acid and
10 endogenous nucleic acid. The heterologous nucleic acid can be targeted to a specific chromosome.

In some instances, knockout animals can be produced. Such an animal can be initially produced by promoting homologous recombination between an MTSP10 polypeptide gene in its chromosome and an exogenous MTSP10
15 polypeptide gene that has been rendered biologically inactive (typically by insertion of a heterologous sequence, *e.g.*, an antibiotic resistance gene). In one embodiment, this homologous recombination is performed by transforming embryo-derived stem (ES) cells with a vector containing the insertionally inactivated MTSP10 polypeptide gene, such that homologous recombination
20 occurs, followed by injecting the ES cells into a blastocyst, and implanting the blastocyst into a foster mother, followed by the birth of the chimeric animal ("knockout animal") in which an MTSP10 polypeptide gene has been inactivated (see Capecchi, *Science* 244:1288-1292 (1989)). The chimeric animal can be bred to produce homozygous knockout animals, which can then be used to
25 produce additional knockout animals. Knockout animals include, but are not limited to, mice, hamsters, sheep, pigs, cattle, and other non-human mammals. For example, a knockout mouse is produced. The resulting animals can serve as models of specific diseases, such as cancers, that exhibit under-expression of an MTSP10 polypeptide. Such knockout animals can be used as animal models
30 of such diseases *e.g.*, to screen for or test molecules for the ability to treat or prevent such diseases or disorders.

RECTIFIED SHEET (RULE 91)

-144-

Other types of transgenic animals also can be produced, including those that over-express the MTSP10 polypeptide. Such animals include "knock-in" animals that are animals in which the normal gene is replaced by a variant, such as a mutant, an over-expressed form, or other form. For example, one species',
5 such as a rodent's endogenous gene can be replaced by the gene from another species, such as from a human. Animals also can be produced by non-homologous recombination into other sites in a chromosome; including animals that have a plurality of integration events.

After production of the first generation transgenic animal, a chimeric
10 animal can be bred to produce additional animals with over-expressed or mis-expressed MTSP10 polypeptides. Such animals include, but are not limited to, mice, hamsters, sheep, pigs, cattle and other non-human mammals. The resulting animals can serve as models of specific diseases, such as cancers, that are exhibit over-expression or mis-expression of an MTSP10 polypeptide. Such
15 animals can be used as animal models of such diseases *e.g.*, to screen for or test molecules for the ability to treat or prevent such diseases or disorders. In a specific embodiment, a mouse with over-expressed or mis-expressed MTSP10 polypeptide is produced.

20 The following examples are included for illustrative purposes only and are not intended to limit the scope of the invention.

EXAMPLE 1

Identification of MTSP10

The protein sequence of the protease domain of matriptase (MTSP1; accession number AF118224) was used to search the human HTGS (High
25 Throughput Genomic Sequence) database using the tblastn search and alignment algorithm, which compares a protein query sequence against a nucleotide sequence database dynamically translated in all six reading frames (both strands). Several potential serine proteases were identified, including the one
30 designated herein as MTSP10.

Based on the incomplete and unordered human genome sequences, MTSP10 appears to be localized in either chromosome 3 (AC024887 and AC

RECTIFIED SHEET (RULE 91)

-145-

073522 clones) or chromosome 8 (AC024964 clone). A search of sequences deposited in GenBank and human EST database showed that no identical sequence has been deposited.

Cloning of cDNA Encoding the Protease Domain of MTSP10

5 Using the nucleotide sequence of MTSP10 derived from the genomic sequence, two gene specific oligonucleotide primers were designed. The sequence for the 5'-end primer (C8-N1A-1) was

5'-CGCATCATCGGAGGCACAGACACCCT-3' (SEQ ID No. 7)

and that of the 3'-end primer (C8-NSP1-3AS) was

10 5'-CCAGGGAACAAAGTTTGACACCCTTG TG -3' (SEQ ID No. 8)

A band of 690 bp was amplified from human pancreas Marathon cDNA (Clontech). Subsequent sequence analysis showed that the nucleotide sequence of this DNA fragment matched that of the genomic MTSP10 exon sequences and includes most of the MTSP10 protease domain. A stop codon was not
15 found in this cDNA sequence.

To obtain the 3'-end of the encoding DNA of the MTSP10 protease domain, 3'-RACE (rapid amplification of cDNA ends) reactions were performed on Marathon-Ready cDNA library from human pancreas (Clontech). Marathon-Ready cDNA are specifically made for RACE reactions.

20 The first RACE reactions were performed by PCR using Marathon cDNA adaptor primer 1 (AP1) with gene specific primers, C8-N1A-1, 5'-CGCATCATCGGAGGCACAGACACCCT-3' (SEQ ID No. 9). The PCR products were purified from agarose gel.

A second nested PCR was then performed using Marathon cDNA adaptor
25 primer 2 (AP2) with gene specific primer (Ch8-NSP1-4)
5'-CTCCCACTGGTCAGAGAGTTCGCAGTG-3' (SEQ ID No. 10; using first 3'-RACE product as template). PCR products from RACE reactions which were larger than 300 bp were cut out and purified from agarose gel and subcloned into pCR2.1-TOPO cloning vector (Invitrogen, Carlsbad, CA). Colony
30 hybridization was then performed to identify positive colonies containing MTSP10 sequence. Positive clones were identified by colony hybridization using a 690 bp DNA fragment containing MTSP10 protease domain sequence

RECTIFIED SHEET (RULE 91)

-146-

(obtained from PCR reaction with primers C8-N1A-1 and C8-NSP1-3AS) and by DNA sequencing. Sequence analysis of the 3'-RACE products indicated that an additional 36 bp sequence including the presumed stop codon was obtained.

Gene expression profile of MTSP10 in normal and tumor tissues

- 5 To obtain information regarding the gene expression profile of the MTSP10 transcript, PCR analysis was carried out on cDNA panels made from several human adult tissues (Clontech, Cat. #K1420-1) cDNA panel using MTSP10-specific primers, C8-N1A-1 (5'-
- 10 CGCATCATCGGAGGCACAGACACCCT-3'; SEQ ID No. 11) and C8-N1A-2AS (5'- CTCCCCACTGCGAACTCTCTGACCAGTG-3'; SEQ ID No. 12). MTSP10 transcript was detected in pancreas, lung and kidney. MTSP10 transcript was also detected in small intestine Marathon-Ready cDNA (Clontech). PCR of the MTSP10 transcript from cDNA libraries made from several human primary tumors xenografted in nude mice (human tumor multiple tissue cDNA panel,
- 15 catalog number K1522-1, CLONTECH) was also performed. The MTSP10 transcript was detected in breast carcinoma (GI-101), lung carcinoma (LX-1 and GI-117), ovarian carcinoma (GI-102), and pancreatic adenocarcinoma (GI-103). The MTSP10 transcript can be weakly detected in prostatic adenocarcinoma (PC3). No apparent signal was detected in two forms of colon adenocarcinomas
- 20 (GI-112 and CX-1). The MTSP10 transcript was also detected in a CWR22R prostate tumor grown on nude mice.

PCR amplification of cDNA encoding full-length protease domain of MTSP10

- To obtain the cDNA fragment encoding the protease domain of MTSP10, an end-to-end PCR amplification using gene-specific primers and the cDNA
- 25 library from human pancreas was used. The two primers used were: 5'-CGCATCATCGGAGGCACAGACACCCT-3' (SEQ ID No. 11) for the 5' end; and 5'-TTACAAAAGAGAAGGGACATATTTATGAATC-3' (SEQ ID No. 21) for the 3' end. The sequences for both primers were derived from the cDNA sequence of MTSP10. The 5' primer contains the sequence that encodes a
- 30 region immediately upstream of the start of the MTSP10 protease domain (RIIGGDTL). The 3' primer corresponds to the sequence immediately after the presumed stop codon. A 720-bp fragment was amplified from the human

-147-

pancreas cDNA library. The PCR product was isolated and purified using the QIAquick gel extraction kit (Qiagen, Valencia, CA; catalog no. 28704), and was confirmed by DNA sequencing analysis.

Serine protease domain of MTSP10 and homology to other proteases

- 5 Sequence analysis of the translated MTSP10 protease domain sequence indicated that MTSP10 contains a trypsin-like serine protease domain characterized by the presence of a protease activation cleavage site at the beginning of the domain and the catalytic triad residues (histidine, aspartate and serine) in 3 highly-conserved regions. Alignment of the protein sequence with
10 that of Matriptase (accession number AF118224; see SEQ ID Nos. 1 and 3) shows 47% identity in the protease domain.

Sequence analysis

- MTSP10 nucleic acid and protein sequences were analyzed using DNA Strider (version 1.2). The cDNA encoding the protease domain of MTSP10 is
15 714 bp in length, which translates into a 238-amino acid protein. The cDNA sequence for the protease domain and the translated protein sequence of MTSP10 is as follows (see, also SEQ ID Nos. 5 and 22 and 6 and 23):

Nucleotide and Amino Acid sequences of Human MTSP10 Protease Domain, and domain organization

20 cDNA/protein sequences

Sequence Range: 1 to 2130

	10	20	30	40	50	60
25	ATAAACCTGGTTTATACACATCTGCCTTCTCCAAATTTATGAGCAGTCTGTTGTTGCA					
	TATTTGGACCAATATGTTGTAGACGGAAGAGGTTTAAATACTCGTCAGACAACAACGT					
	I	N	L	V	Y	T
	T	S	A	F	S	K
	F	Y	E	Q	S	V
	V	A				
	70	80	90	100	110	120
30	GATGTCAGCAGCAACAACAAAGGCGGCCTCCTTGTCACCTTTTGGATTGTTTGTCTATG					
	CTACAGTCGTCGTTGTTGTTTCCGCCGAGGAACAGGTGAAAACCTAACAAAAACAGTAC					
	D	V	S	S	N	N
	K	G	G	L	L	V
	H	F	W	I	V	F
	V	M				
	130	140	150	160	170	180
35	CCACGTGCCAAAGGCCACATCTTCTGTGAAGACTGTGTTGCCGCCATCTTGAAGGACTCC					
	GGTGCACGGTTTCCGGTGTAGAAGACACTTCTGACACAACGGCGGTAGAACTTCTCTGAGG					
	P	R	A	K	G	H
	I	F	C	E	D	C
	V	A	A	I	L	K
	D	S				
	190	200	210	220	230	240
40	ATCCAGACAGCATATAAACCGGACCTCTGTGGGGAGCTTGCAGGGACTGGCTGTGGAC					
	TAGGTCGTGTTTCGTAGTATTTGGCCTGGAGACACCCCTCGAACGTCCTGACCGACACCTG					
	I	Q	T	S	I	I
	N	R	T	S	V	G
	S	L	Q	G	L	A
	V	D				
	250	260	270	280	290	300
45	ATGGACTCTGTGGTACTAAATGCTGGGCTTCGGTCAGATTACTCGTCAACCATAGGATCT					
	TACCTGAGACACCATGATTTACGACCCGAAGCCAGTCTAATGAGCAGTTGGTATCCTAGA					
	M	D	S	V	V	L
	N	A	G	L	R	S
	D	Y	S	S	T	I
	G	S				

-148-

5
10
15
20
25
30
35
40
45
50
55
60
65
70

310 320 330 340 350 360
GACAAAGGCTGCTCTCAGTACTTCTATGCAGAGCATCTGTCTCTCCACTACCCGCTGGAG
CTGTTTCCGACGAGAGTTCATGAAGATACGTCTCGTAGACAGAGAGGTGATGGGCGACCTC
D K G C S Q Y F Y A E H L S L H Y P L E

370 380 390 400 410 420
ATTCTGCAGCCTCAGGGAGGCTGATGTGTCACTTCAAGCTGGTGGCCATAGTGGGTAC
TAAAGACGTTCGGAGTCCCTCCGACTACACAGTGAAGTTCGACCACCGGTATCACCCGATG
I S A A S G R L M C H F K L V A I V G Y

430 440 450 460 470 480
CTGATTCTCTCTCAATCAAGTCCATCCAAATCGAAGCCGACAACTGTGTCACTGACTCC
GACTAGCAGAGAGTTCAGGTAGGTTTAGCTTCGGCTGTTGACACAGTGAAGTGAAGG
L I R L S I K S I Q I E A D N C V T D S

490 500 510 520 530 540
CTGACCATTACGACTCCCTTTTGCCCATCCGGAGCAGCATCTTGTACAGAATTTGTGA
GACTGGTAAATGCTGAGGGAAACGGGTAGGCTCGTGTAGAACATGTCTTAAACACTT
L T I Y D S L L P I R S S I L Y R I C E

550 560 570 580 590 600
CCCACAGAACATTAATGTCTTGTCTTCTACAAATAATCTCATGTGGTGACATTTAAG
GGGTGTTCTTGTAAATACAGTAAACAAAGATGTTTATTAGAGTACAACTGTAAATTC
P T R T L M S F V S T N N L M L V T F K

610 620 630 640 650 660
TCTCTCATATACGGAGGCTCTCAGGAATCCGGGCATATTTTGAGGTCAITCCAGAACAA
AGAGGAGTATATGCTCCGAGAGTCTTAGGCCGTATAAACTCCAGTAAAGTCTTGT
S P H I R R L S G I R A Y F E V I P E Q

670 680 690 700 710 720
AAGTGTGAAAACACAGTGTGGTCAAAGACATCACTGGCTTTGAAGGGAAAATTTCAAGC
TTCACACTTTTGTGTACACACAGTTCCTGTAGTGACCGAACTTCCCTTTTAAAGTTCG
K C E N T V L V K D I T G F E G K I S S

730 740 750 760 770 780
CCATATTACCCGAGCTACTATCCTCCAAATGCAAGTGTACCTGGAAATTTAGACTTCT
GGTATAATGGGCTCGATGATAGGAGGTTTACGTTACATGGACCTTTAAAGTCTGAAGA
P Y Y P S Y Y P P K C K C T W K F Q T S

790 800 810 820 830 840
CTATCAACTCTTGGCATAGCACTGAAATTCATAACTATTCAATAACCAAGAAGAGTATG
GATAGTTGAGAACCCTATCGTGACTTTAAGATATTGATAAGTATTGGTTCTTCTCATAC
L S T L G I A L K F Y N Y S I T K K S M

850 860 870 880 890 900
AAAGGCTGTGAGCATGGATGGTGGGAAATTTATGAGCACATGTACTGTGGCTCCTACATG
TTCCGACACTCGTACCTACCACCTTTAAATACTCGTGTACATGACACCGAGGATGTAC
K G C E H G W W E I Y E H M Y C G S Y M

910 920 930 940 950 960
GATCATCAGACAATTTTTCGAGTGCCCGAGCCCTCTGGTTACATTACAGTCCAGTGCAGT
CTAGTAGTCTGTAAAAAGCTCACGGGTCGGGAGACCAAGTGAAGTTCGAGGTCACGTCA
D H Q T I F R V P S P L V H I Q L Q C S

970 980 990 1000 1010 1020
TCAAGGCTTTTCAGGCAAGCCACTTTTGGCAGAATATGGCAGTTACAACATCAGTCAACCC
AGTTCCGAAAGTCCGTTTCGGTGAAAACCGTCTTATACCGTCAATGTTGTAGTCAAGTGGG
S R L S G K P L L A E Y G S Y N I S Q P

1030 1040 1050 1060 1070 1080
TGCCCTGTGGGATCTTTAGATGCTCTCCGGTTTATGTGTCCCTCAGGCCAGCGTGGT
ACGGGACACCCTAGAAAATCTACGAGGAGGCCAAATACACAGGGAGTCCGGGTCGACCA
C P V G S F R C S S G L C V P Q A Q R G

1090 1100 1110 1120 1130 1140
GATGGAGTAAATGACTGCTTTGATGAAAGTGAAGTGTGTTTTCGCTGAGCCCTCAACCT
CTACCTCATTTACTGACGAACTACTTCACTACTTGACAAAACGCACTCGGGAGTTGGA
D G V N D C F D E S D E L F C V S P Q P

1150 1160 1170 1180 1190 1200

-149-

GCCTGCAATACCAGCTCCTTCAGGCAGCATGGCCCTCTCATCTGTGATGGCTTCAGGGAC
CGGACGTTATGGTCGAGGAAGTCCGTCGTACCGGGAGAGTAGACACTACCGAAGTCCCTG
A C N T S S F R Q H G P L I C D G F R D

5 1210 1220 1230 1240 1250 1260
TGTGAGAATGGCCGGGATGAGCAAACTGCACTCAAAGTATTCATGCAACAACAGAACT
ACACTCTTACCGGCCCTACTCGTTTGGACGTGAGTTTCATAAGGTACGTTGTTGCTTGA
C E N G R D E Q N C T Q S I P C N N R T

10 1270 1280 1290 1300 1310 1320
TTTAAGTGTGGCAATGATATTTGCTTTAGGAAACAAAATGCAAAATGTGATGGGACAGTG
AAATTACACCGGTTACTATAAACGAAATCCTTTGTTTACGTTTACACTACCCTGTCAC
F K C G N D I C F R K Q N A K C D G T V

15 1330 1340 1350 1360 1370 1380
GATTGTCCAGATGGAAGTGATGAAGAAGGCTGCACCTGCAGCAGGAGTTCTCCGCCCTT
CTAACAGGTCTACCTTCACTACTTCTTCCGACGTGGACGTCGTCTCAAGGAGGCGGGAA
D C P D G S D E E G C T C S R S S S A L

20 1390 1400 1410 1420 1430 1440
CACCGCATCATCGGAGGCACAGACACCTGGAGGGGGTTGGCCGTGGCAGGTGAGCCTC
GTGGCGTAGTAGCCTCCGTGTCTGTGGGACCTCCCCCAACCGGCACCGTCCAGTCGGAG
H R I I G G T D T L E G G W P W Q V S L

25 1450 1460 1470 1480 1490 1500
CACTTTGTTGGATCTGCCCTACTGTGGTGCCTCAGTCATCTCCAGGGAGTGGCTTCTTTCT
GTGAAACAACCTAGACGGATGACACCACGGAGTCAGTAGAGGTCCCTCACCGAAGAAAGA
H F V G S A Y C G A S V I S R E W L L S

30 1510 1520 1530 1540 1550 1560
GCAGCCCACTGTTTTCATGGAACAGGCTGTGAGATCCACACCATGGACTGACACACCTC
CGTCGGGTGACAAAGTACCTTTGTCCGACAGTCTAGGGTGTGGTACCTGACGTGGAG
A A H C F H G N R L S D P T P W T A H L

35 1570 1580 1590 1600 1610 1620
GGGATGTATGTTGAGGGGAATGCCAAGTTTGTCTCCCGGTGAGAAGAATTGTGGTCCAC
CCCTACATACAAGTCCCCTTACGGTTCAAACAGAGGGGCCACTCTTCTTAACACCAGGTG
G M Y V Q G N A K F V S P V R R I V V H

40 1630 1640 1650 1660 1670 1680
GAGTACTATAACAGTCAGACTTTTGATTATGATATTGCTTTGCTACAGCTCAGTATTGCC
CTCATGATATTGTGAGTCTGAAACTAATACTATAACGAAACGATGTGAGTCATAACGG
E Y Y N S Q T F D Y D I A L L Q L S I A

45 1690 1700 1710 1720 1730 1740
TGGCCTGAGACCCCTGAACAGCTCATTGAGCCCAATATGCAATTCCTCCCACTGGTCAGAGA
ACCGGACTCTGGGACTTTGTGAGTAAGTCGGTTATACGTAAGGAGGGTGACCACTCTCT
W P E T L K Q L I Q P I C I P P T G Q R

50 1750 1760 1770 1780 1790 1800
GTTGCGAGTGGGGAGAAGTGTCTGGGTAAGTGGCTGGGGGCGAAGACAGCAAGCAGATAAT
CAAGCGTCACCCCTCTTACGACCCATTGACCGACCCCGCTTCTGTGCTTCTGTCTATTA
V R S G E K C W V T G W G R R H E A D N

55 1810 1820 1830 1840 1850 1860
AAAGGCTCCCTCGTTCTGCAGCAAGCGGAGGTAGAGCTCATTGATCAAACGCTCTGTGTT
TTCCGAGGGAGCAAGACGTCGTTTCGCTCCATCTCGAGTAAGTGTGCGAGACACAA
K G S L V L Q Q A E V E L I D Q T L C V

60 1870 1880 1890 1900 1910 1920
TCCACCTACGGGATCATCACTTCTCGGATGCTCTGTGACAGGCATAATGTGAGCAAGAGA
AGGTGGATGCCCTAGTAGTGAAGAGCCTACGAGACAGTCCGTATTACAGTCCGTTCTCT
S T Y G I I T S R M L C A G I M S G K R

65 1930 1940 1950 1960 1970 1980
GATGCTGCAAAAGGAGATTCCGGGTGGACCTTTATCTTGTGCAAGAAAAAGTGATGGAAAA
CTACGGACGTTTCTCTAAGCCACCTGGAAATAGAACAGCTTCTTTTCACTACCTTTT
D A C K G D S G G P L S C R R K S D G K

70 1990 2000 2010 2020 2030 2040
TGGATTTGACTGGCATTTGTTAGCTGGGGACATGGATGTGACGACCAAACTTCTCTGGT

-150-

ACCTAAACTGACCGTAACAATCGACCCCTGTACCTACACCTGCTGGTTTGAAAGGACCA
W I L T G I V S W G H G C G R P N F P G

2050 2060 2070 2080 2090 2100
5 GTTTACACAAGGGTGTCAAACCTTTGTTCCCTGGATTTCATAAATATGTCCTTCTCTTTTG
CAAATGTGTTCCACAGTTTGAACAAGGGACCTAAGTATTTATACAGGGAAGAGAAAAC
V Y T R V S N F V P W I H K Y V P S L L

2110 2120 2130
10 TAATTGCAAAAAAAAAAAAAAAAAAAAAA
ATTAACGTTTTTTTTTTTTTTTTTTTTTTT

Protein sequence

Sequence Range: 1 to 701

15 10 20 30 40 50 60
INLVYTTSAFSKFYEQSVVADVSSNNKGGLLVHFWIVFVMPRAKGHIFCEDCVAAILKDS

70 80 90 100 110 120
20 IQTSIINRTSVGSLQGLAVDMDSVVLNAGLRSDYSSSTIGSDKGCSQYFYAEHLSLHYPLE

130 140 150 160 170 180
ISAASGRLMCHFVLVAIVGYLIRLSIKSIQIEADNCVTDLSLTIYDSSLPIRSSILYRICE

190 200 210 220 230 240
25 PTRTLMFSFVSTNNMLLVTFKSPHIRLSGIRAYFEVIPEQKCENTVLVKDITGFEGKISS

250 260 270 280 290 300
PYYPSYYPKCKCTWKPFQTSLSLGLIALKFYNYSITKKSMMKGCEHGWWEIYEHMYCGSYM

30 310 320 330 340 350 360
DHQTI FRVPSPLVHIQLQCSSRLSGKPLLA EYGSYNISQPCPVGSFRCS SGLCVPQAQRG

370 380 390 400 410 420
35 DGVNDCFDESDELFCVSPQACNTSSFRQHGPLICDGFRCENGRDEQNCTQSIPCNNRT

430 440 450 460 470 480
FKCGNDICFRKQNAKCDGTVDGPDGSDEEGCTCSRSSSALHRIIGGTDLTLEGGWPWQVSL

490 500 510 520 530 540
40 HFVGSAYCGASVISREWLLSAAHCFHGNRLSDPTPWTALHGMVYQGNKRVSPVRRIVVH

550 560 570 580 590 600
EYNSQTFDYDIALQLLSIAWPETLKLQIPIPIPTGQVRSGEKCWVTGWGRRHEADN

610 620 630 640 650 660
45 KGSLVLQQAELIDQTL CVSTYGIITSRMLCAGIMSGKRDACKGDSGGPLSCRRKSDGK

670 680 690 700
50 WILTGIVSWGHGCGRPNFPGVYTRVSNFVPIHKYVPSLL*

Domain organization

55 CUB domain aa104-217
CUB domain aa222-335
LDLa domain aa340-377
LDLa domain aa381-412
LDLa domain aa415-453
Ser prot domain aa462-692

-151-

EXAMPLE 2**Expression of the protease MTSP domains**

Nucleic acid encoding the MTSP10 and protease domain thereof can be cloned into a derivative of the *Pichia pastoris* vector pPIC9K (available from

5 Invitrogen; see SEQ ID NO. 13). Plasmid pPIC9k features include the 5' AOX1 promoter fragment at 1-948; 5' AOX1 primer site at 855-875; alpha-factor secretion signal(s) at 949-1218; alpha-factor primer site at 1152-1172; multiple cloning site at 1192-1241; 3' AOX1 primer site at 1327-1347; 3' AOX1 transcription termination region at 1253-1586; HIS4 ORF at 4514-1980;

10 kanamycin resistance gene at 5743-4928; 3' AOX1 fragment at 6122-6879; ColE1 origin at 7961-7288; and the ampicillin resistance gene at 8966-8106. The plasmid used herein is derived from pPIC9K by eliminating the XhoI site in the kanamycin resistance gene and the resulting vector is herein designated pPIC9KX. Expression in *Pichia* can be performed using known methods (see,

15 *e.g.*, Zhang *et al.* (2000) *Biotechnology and Bioengineering* 70: No 1 October 5, 2000).

Mutagenesis of the Protease Domain of MTSP10 for Expression in Pichia

The gene encoding the protease domain of MTSP10 (residues 462-692 SEQ ID No. 23) was mutagenized by PCR SOE (PCR-based splicing by overlap

20 extension) to replace the unpaired cysteine at position 122 (chymotrypsin numbering system; Cys at residue 573, SEQ ID No. 23) with a serine. Two overlapping gene fragments, each containing the AGC codon for serine at position 122 were PCR amplified using the following primers: for the 5' gene fragment, AGTTAACTCGAGAAAAGATCATCGGAGGCACAGACACCCTG SEQ

25 ID No. 15 and ACCAGTGGGAGGAATGCTTATTGGCTGAATGAG SEQ ID No. 16; for the 3' gene fragment,

AGTTAAGAATTCCAAAAGAGAAGGGACATATTTATG SEQ ID No. 17 and CTCATTGAGCCAATAAGCATTCTCCCACTGGT SEQ ID No. 18. The amplified gene fragments were purified on a 1% agarose gel, mixed and

30 reamplified by PCR to produce the full length coding sequence for MTSP10 C122S. This sequence was then cut with restriction enzymes EcoRI and XhoI, and ligated into vector p9KXST2. Vector p9KXST2 was derived from pPic9KX

-152-

with bases 1228-1233 (CCTAGG) replaced with the sequence GGAGGTTGGTCTCATCCACAATTTGAAAAGTAA SEQ ID No. 19, which codes for a c-terminal streptagII sequence with a one glycine linker and stop codon (GGWSHPQFEK-Stop; SEQ ID No. 20).

5 Expression of MTSP10-ST2 C122S

P. pastoris clone GS115/pPIC9KXST2:MTSP10-ST2 C122S Sac MC5 expressing the C122S (chymotrypsin numbering; Cys₅₇₃ in SEQ ID No. 23) mutant and C-terminally Strep-tag II tagged form of MTSP10 was fermented at the 5 liter scale. An overnight culture of 200 ml (OD600 of approximately 24) was used to
10 inoculate 3.2 liters of fermentation medium in each of six Bioflo vessels (New Brunswick Scientific, Edison, NJ). The batch phase complex medium contained 10 g/l yeast extract, 20 g/l peptone, 40 g/l glycerol, 5 g/l ammonium sulfate, 0.2 g/l calcium sulfate(dihydrate), 2 g/l magnesium sulfate(heptahydrate), 2 g/l potassium sulfate, 25 g/l sodium hexametaphosphate, and 4.35 ml/l PTM1. The
15 culture was grown at a pH of 5.8 and a temperature of 28 °C in the batch phase. Concentrated ammonium hydroxide was used to maintain the pH of the culture. KFO 880 (KABO Chemicals, Cheyenne,WY) was used as needed to control foaming.

The batch phases of the fermentations lasted about 23-28 hours at which
20 time the culture had consumed all of the initial glycerol in the media. A substrate limited fed-batch of 50% (w/v) glycerol was initiated at 18 ml/l*hr at this point. One hour into the glycerol fed-batch the pH of the culture was linearly increased from 5.8 to 7.0 over a two hour period by addition of concentrated ammonium hydroxide. The glycerol fed-batch was from 3 to 4
25 hours in duration. The cultures reached densities of 204-234 g/l wet cell weight by this point.

Methanol induction was initiated following the end of the glycerol fed-batch phase. The culture was transitioned to methanol utilization by the method of Zhang et al. by adding 1.5 ml of methanol per liter of culture and linearly
30 decreasing the glycerol feed rate from 18 ml/l*hr to 0 ml/l*hr over a 3 hour period. The methanol addition served as an on-line calibration of the MeOH Sensor (Raven Biotech, Vancouver, BC, Canada) used to control the fermenter

-153-

throughout induction. After the initial amount of methanol was utilized, as indicated by the MeOH Sensor, another 1.5 ml/l was added to the culture and the MeOH sensor was used to control the methanol concentration in the fermenter at that level throughout the induction phase. The methanol fed to the
5 fermenter was supplemented with 2 ml/l PTM4 solution. The induction phase lasted about 40 to 44 hours.

Recovery and Purification of MTSP-10

The supernatant from each of the fermentations was harvested after removal of cells by centrifugation. Supernatants were pooled and concentrated
10 to about 1 liter using a 10 kDa ultrafiltration cartridge (A/G Technologies Corp., Needham, MA) on a SRT5 ultrafiltration system (North Carolina SRT, Cary, NC). The concentrate was diafiltered with 3 volumes of buffer A. The concentrate was drained from the system, then the system was rinsed with a volume of buffer A equal to the concentrated material. The concentrate and the rinse
15 material were combined to yield the final ultrafiltration product of about 1 liter. A final clarification of the supernatant was done with a SartoBran 300 0.45 + 0.2 μ m capsule filter (Sartorius Separations Div., Edgewood, NJ).

The diafiltered MTSP10-ST2 was slowly loaded overnight onto 90 ml of benzamidine sepharose (Amersham Pharmacia Biotech, Piscataway, NJ) which
20 had been equilibrated in buffer A. The column was then washed with 8 column volumes of Buffer B to remove contaminants. Elution of the MTSP10-ST2 was achieved using buffer C. Fractions of 45 ml were collected and analyzed by activity and SDS-PAGE before pooling the desired material to be further purified.

The MTSP10 post-benzamidine sample (usually about 150ml containing
25 2-5mg mtsp10 and 100mM benzamidine) was dialyzed into 100mM Tris, 150mM NaCl, 1mM EDTA, 0.001% tween80 pH8.0 binding buffer. Strep-Tactin Macroprep resin (Cat. No. 2-1505-010; IBA GmbH, Rudolf-Wissell-Str. 28, D-37079 Gottingen, Germany) was used to retain mtsp10 activity. The resin suspension was incubated in a cold room for 1 hour with shaking. The
30 completeness of mtsp10 binding to the resin was examined by monitoring the mtsp10 activity in the supernatant. Usually 50-70% of mtsp 10 activity can be retained by the resin. The resin was then packed into a column and washed with

-154-

5 column volumes of binding buffer. Mtsp10 activity was eluted using binding buffer containing 2.5mM D-desthiobiotin (Sigma product). Active fractions were identified by activity assay using substrate Spec-tPA (Chromogenics), pooled and dialyzed to remove D-desthiobiotin.

5 **PTM1:**

6.0 g/l $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 0.08 g/l NaI, 3.0 g/l $\text{MnSO}_4 \cdot \text{H}_2\text{O}$, 0.2 g/l $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$, 0.02 g/l H_3BO_3 , 0.5 g/l CoCl_2 , 20.0 g/l ZnCl_2 , 65.0 g/l $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.2 g/l biotin, 5.0 ml/l H_2SO_4 .

PTM4:

10 2.0 g/l $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 0.08 g/l NaI, 3.0 g/l $\text{MnSO}_4 \cdot \text{H}_2\text{O}$, 0.2 g/l $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$, 0.02 g/l H_3BO_3 , 0.5 g/l $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$, 7.0 g/l ZnCl_2 , 22.0 g/l $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.2 g/l biotin, 1.0 ml/l H_2SO_4

Buffer A: 50 mM Tris, pH 8.0, 50 mM NaCl, 0.005% Tw-80

Buffer B: 50 mM Tris, pH 8.0, 1.0 M NaCl, 0.005% Tw-80

15 Buffer C: 50 mM Tris, pH 8.0, 50 mM NaCl, 100 mM benzamidine, 0.005% Tw-80

EXAMPLE 3

20 **Assays for identification of candidate compounds that modulate that activity of an MTSP**

Assay for screening MTSP10 Inhibitors

The protease domain of MTSP10 expressed in *Pichia pastoris* was assayed for inhibition by various compounds as follows in Costar 96 well tissue culture plates (Corning NY). Approximately 1-10 nM MTSP10 is added without inhibitor, or with 100000 nM inhibitor and 7 1:6 dilutions to 1X direct buffer (29.2 mM Tris, pH 8.4, 29.2 mM Imidazole, 217 mM NaCl (100 μL final volume)), and allowed to incubate at room temperature for 30 minutes. 400 μM substrate Spectrozyme t-PA (American Diagnostica,, Greenwich, CT) is added and reaction is monitored in a SpectraMAX Plus microplate reader (Molecular Devices, Sunnyvale CA) by following change in absorbance at 405 nm for 20 minutes at 37°C. Spectrozyme UK can also be used as the substrate in this assay.

Identification of substrates

-155-

Other substrates for use in the assays can be identified empirically by testing substrates. The following list of substrates are exemplary of those that can be tested.

	Substrate name	Structure
5	S 2366	pyroGlu-Pro-Arg-pNA.HCl
	spectrozyme t-PA	CH ₃ SO ₂ -D-HHT-Gly-Arg-pNA.AcOH
	N-p-tosyl-Gly-Pro-Arg-pNA	N-p-tosyl-Gly-Pro-Arg-pNA
	Benzoyl-Val-Gly-Arg-pNA	Benzoyl-Val-Gly-Arg-pNA
	Pefachrome t-PA	CH ₃ SO ₂ -D-HHT-Gly-Arg-pNA
10	S 2765	N- α -Z-D-Arg-Gly-Arg-pNA.2HCl
	S 2444	pyroGlu-Gly-Arg-pNA.HCl
	S 2288	H-D-Ile-Pro-Arg-pNA.2HCl
	spectrozyme UK	Cbo-L-(γ)Glu(α -t-BuO)-Gly-Arg-pNA.2AcOH
	S 2302	H-D-Pro-Phe-Arg-pNA.2HCl
15	S 2266	H-D-Val-Leu-Arg-pNA.2HCl
	S 2222	Bz-Ile-Glu(g-OR)-Gly-Arg-pNA.HCl R = H(50%) and R = CH ₃ (50%)
	Chromozyme PK	Benzoyl-Pro-Phe-Arg-pNA
	S 2238	H-D-Phe-Pip-Arg-pNA.2HCl
	S 2251	H-D-Val-Leu-Lys-pNA.2HCl
20	Spectrozyme PI	H-D-Nle-HHT-Lys-pNA.2AcOH
		Pyr-Arg-Thr-Lys-Arg-AMC
		H-Arg-Gln-Arg-Arg-AMC
		Boc-Gln-Gly-Arg-AMC
		Z-Arg-Arg-AMC
25	Spectrozyme THE	H-D-HHT-Ala-Arg-pNA.2AcOH
	Spectrozyme fXIIa	H-D-CHT-Gly-Arg-pNA.2AcOH
		CVS 2081-6 (MeSO ₂ -dPhe-Pro-Arg-pNA)
		Pefachrome fVIIa (CH ₃ SO ₂ -D-CHA-But-Arg-pNA)

pNA = para-nitranilide (chromogenic)

30 AMC = amino methyl coumarin (fluorescent)

If none of the above substrates are cleaved, a coupled assay, can be used. Briefly, test the ability of the protease to activate and enzyme, such as plasminogen and trypsinogen. To perform these assays, the single chain protease is incubated with a zymogen, such as plasminogen or trypsinogen, in the presence of the a known substrate, such, lys-plasminogen, for the zymogen. If the single chain activates the zymogen, the activated enzyme, such as plasmin and trypsin, will degrade the substrate therefor.

EXAMPLE 4**Other Assays**

These assays are described with reference to MTSP1, but such assays can be readily adapted for use with MTSP10.

5 Amidolytic Assay for Determining Inhibition of Serine Protease Activity of Matriptase or MTSP1

The ability of test compounds to act as inhibitors of rMAP catalytic activity was assessed by determining the inhibitor-induced inhibition of amidolytic activity by the MAP, as measured by IC_{50} values. The assay buffer
10 was HBSA (10 mM Hepes, 150mM sodium chloride, pH 7.4, 0.1% bovine serum albumin). All reagents were from Sigma Chemical Co. (St. Louis, MO), unless otherwise indicated.

Two IC_{50} assays (a) one at either 30-minutes or 60-minutes (a 30-minute or a 60-minute preincubation of test compound and enzyme) and (b) one at
15 0-minutes (no preincubation of test compound and enzyme) were conducted. For the IC_{50} assay at either 30-minutes or 60-minutes, the following reagents were combined in appropriate wells of a Corning microtiter plate: 50 microliters of HBSA, 50 microliters of the test compound, diluted (covering a broad concentration range) in HBSA (or HBSA alone for uninhibited velocity
20 measurement), and 50 microliters of the rMAP (Corvas International) diluted in buffer, yielding a final enzyme concentration of 250 pM as determined by active site filtration. Following either a 30-minute or a 60-minute incubation at ambient temperature, the assay was initiated by the addition of 50 microliters of the substrate S-2765 (N- α -Benzyloxycarbonyl-D-arginyl-L-glycyl-L-arginine-p-
25 nitroaniline dihydrochloride; DiaPharma Group, Inc.; Franklin, OH) to each well, yielding a final assay volume of 200 microliters and a final substrate concentration of 100 μ M (about 4-times K_m). Before addition to the assay mixture, S-2765 was reconstituted in deionized water and diluted in HBSA. For the IC_{50} assay at 0 minutes; the same reagents were combined: 50 microliters of
30 HBSA, 50 microliters of the test compound, diluted (covering the identical concentration range) in HBSA (or HBSA alone for uninhibited velocity measurement), and 50 microliters of the substrate S-2765. The assay was

-157-

initiated by the addition of 50 microliters of rMAP. The final concentrations of all components were identical in both IC_{50} assays (at 30- or 60- and 0-minute).

The initial velocity of chromogenic substrate hydrolysis was measured in both assays by the change of absorbance at 405 nm using a Thermo Max®

- 5 Kinetic Microplate Reader (Molecular Devices) over a 5 minute period, in which less than 5% of the added substrate was used. The concentration of added inhibitor, which caused a 50% decrease in the initial rate of hydrolysis was defined as the respective IC_{50} value in each of the two assays (30- or 60-minutes and 0-minute).

10 ***In vitro* enzyme assays for specificity determination**

The ability of compounds to act as a selective inhibitor of matriptase activity was assessed by determining the concentration of test compound that inhibits the activity of matriptase by 50%, (IC_{50}) as described in the above Example, and comparing IC_{50} value for matriptase to that determined for all or

- 15 some of the following serine proteases: thrombin, recombinant tissue plasminogen activator (rt-PA), plasmin, activated protein C, chymotrypsin, factor Xa and trypsin.

The buffer used for all assays was HBSA (10 mM HEPES, pH 7.5, 150 mM sodium chloride, 0.1% bovine serum albumin).

- 20 The assay for IC_{50} determinations was conducted by combining in appropriate wells of a Corning microtiter plate, 50 microliters of HBSA, 50 microliters of the test compound at a specified concentration (covering a broad concentration range) diluted in HBSA (or HBSA alone for V_0 (uninhibited velocity) measurement), and 50 microliters of the enzyme diluted in HBSA. Following a
- 25 30 minute incubation at ambient temperature, 50 microliters of the substrate at the concentrations specified below were added to the wells, yielding a final total volume of 200 microliters. The initial velocity of chromogenic substrate hydrolysis was measured by the change in absorbance at 405 nm using a Thermo Max® Kinetic Microplate Reader over a 5 minute period in which less
- 30 than 5% of the added substrate was used. The concentration of added inhibitor which caused a 50% decrease in the initial rate of hydrolysis was defined as the IC_{50} value.

RECTIFIED SHEET (RULE 91)

-158-

Thrombin (fIIa) Assay

Enzyme activity was determined using the chromogenic substrate, Pefachrome t-PA (CH_3SO_2 -D-hexahydrotyrosine-glycyl-L-Arginine-p-nitroaniline, obtained from Pentapharm Ltd.). The substrate was reconstituted in deionized water prior to use. Purified human α -thrombin was obtained from Enzyme Research Laboratories, Inc. The buffer used for all assays was HBSA (10 mM HEPES, pH 7.5, 150 mM sodium chloride, 0.1% bovine serum albumin).

IC_{50} determinations were conducted where HBSA (50 μL), α -thrombin (50 μl) (the final enzyme concentration is 0.5 nM) and inhibitor (50 μl) (covering a broad concentration range), were combined in appropriate wells and incubated for 30 minutes at room temperature prior to the addition of substrate Pefachrome-t-PA (50 μl) (the final substrate concentration is 250 μM , about 5 times K_m). The initial velocity of Pefachrome t-PA hydrolysis was measured by the change in absorbance at 405 nm using a Thermo Max[®] Kinetic Microplate Reader over a 5 minute period in which less than 5% of the added substrate was used. The concentration of added inhibitor which caused a 50% decrease in the initial rate of hydrolysis was defined as the IC_{50} value.

Factor Xa

Factor Xa catalytic activity was determined using the chromogenic substrate S-2765 (N-benzyloxycarbonyl-D-arginine-L-glycine-L-arginine-p-nitroaniline), obtained from DiaPharma Group (Franklin, OH). All substrates were reconstituted in deionized water prior to use. The final concentration of S-2765 was 250 μM (about 5-times K_m). Purified human Factor X was obtained from Enzyme Research Laboratories, Inc. (South Bend, IN) and Factor Xa (FXa) was activated and prepared from it as described [Bock, P.E., Craig, P.A., Olson, S.T., and Singh, P. *Arch. Biochem. Biophys.* 273:375-388 (1989)]. The enzyme was diluted into HBSA prior to assay in which the final concentration was 0.25 nM.

Recombinant tissue plasminogen activator (rt-PA) Assay

rt-PA catalytic activity was determined using the substrate, Pefachrome t-PA (CH_3SO_2 -D-hexahydrotyrosine-glycyl-L-arginine-p-nitroaniline, obtained from Pentapharm Ltd.). The substrate was made up in deionized water followed by dilution in HBSA prior to the assay in which the final concentration was 500

-159-

micromolar (about 3-times K_m). Human rt-PA (Activase®) was obtained from Genentech Inc. The enzyme was reconstituted in deionized water and diluted into HBSA prior to the assay in which the final concentration was 1.0 nM.

Plasmin Assay

- 5 Plasmin catalytic activity was determined using the chromogenic substrate, S-2366 (L-pyroglutamyl-L-prolyl-L-arginine-p-nitroaniline hydrochloride), which was obtained from DiaPharma group. The substrate was made up in deionized water followed by dilution in HBSA prior to the assay in which the final concentration was 300 micromolar (about 2.5-times K_m).
- 10 Purified human plasmin was obtained from Enzyme Research Laboratories, Inc. The enzyme was diluted into HBSA prior to assay in which the final concentration was 1.0 nM.

Activated Protein C (aPC) Assay

- aPC catalytic activity was determined using the chromogenic substrate,
- 15 Pefachrome PC (delta-carbobenzloxy-D-lysine-L-prolyl-L-arginine-p-nitroaniline dihydrochloride), obtained from Pentapharm Ltd.). The substrate was made up in deionized water followed by dilution in HBSA prior to the assay in which the final concentration was 400 micromolar (about 3-times K_m). Purified human aPC was obtained from Hematologic Technologies, Inc. The enzyme was diluted
- 20 into HBSA prior to assay in which the final concentration was 1.0 nM.

Chymotrypsin Assay

- Chymotrypsin catalytic activity was determined using the chromogenic substrate, S-2586 (methoxy-succinyl-L-arginine-L-prolyl-L-tyrosyl-p-nitroanilide), which was obtained from DiaPharma Group. The substrate was made up in
- 25 deionized water followed by dilution in HBSA prior to the assay in which the final concentration was 100 micromolar (about 9-times K_m). Purified (3X-crystallized; CDI) bovine pancreatic alpha-chymotrypsin was obtained from Worthington Biochemical Corp. The enzyme was reconstituted in deionized water and diluted into HBSA prior to assay in which the final concentration was 0.5 nM.

-160-

Trypsin Assay

Trypsin catalytic activity was determined using the chromogenic substrate, S-2222 (benzoyl-L-isoleucine-L-glutamic acid-[gamma-methyl ester]-L-arginine-p-nitroanilide), which was obtained from DiaPharma Group. The

5 substrate was made up in deionized water followed by dilution in HBSA prior to the assay in which the final concentration was 250 micromolar (about 4-times K_m). Purified (3X-crystallized; TRL3) bovine pancreatic trypsin was obtained from Worthington Biochemical Corp. The enzyme was reconstituted in deionized water and diluted into HBSA prior to assay in which the final concentration was

10 0.5 nM.

Since modifications will be apparent to those of skill in this art, it is intended that this invention be limited only by the scope of the appended claims.

-161-

WHAT IS CLAIMED IS:

1. A substantially purified single or two chain polypeptide, comprising the protease domain of a type-II membrane-type serine protease 10 (MTSP10) or a catalytically active portion thereof.

5 2. The polypeptide of claim 1 that is an activated two chain polypeptide.

3. A polypeptide of claim 1 or claim 2 selected from the group consisting of

10 a polypeptide that comprises a sequence of amino acids encoded by the sequence that includes at least about 60%, 70%, 80% or 90% amino acid sequence identity with the sequence of amino acids set forth in SEQ ID No. 6;

15 a polypeptide that comprises a sequence of amino acids encoded by the sequence that includes at least about 60%, 70%, 80% or 90% amino acid sequence identity with the sequence of amino acids set forth in SEQ ID No. 23;

a polypeptide that comprises a sequence of amino acids encoded by the sequence of nucleotides set forth in SEQ ID No. 6 or SEQ ID No. 23;

20 a polypeptide that comprises a sequence of amino acids encoded by a sequence of nucleotides that hybridizes along at least 70% of its full-length under conditions of high stringency to the sequence of nucleotides set forth as SEQ ID No. 5 or SEQ ID No. 23;

25 a polypeptide that is encoded by a sequence of nucleotides that is a splice variant of the sequence of nucleotides that comprises the sequence set forth in SEQ ID No. 5 or SEQ ID No. 23.

4. The polypeptide of any of claims 1-3, wherein:

the MTSP10 portion of the polypeptide consists essentially of the protease domain of the MTSP10 or a catalytically active portion thereof.

30 5. The substantially purified polypeptide of any of claims 1-4, wherein the MTSP10 is a human polypeptide.

-162-

6. The substantially purified polypeptide of any of claims 1-5, that consists essentially of the protease domain of MTSP10 or a catalytically active portion of the protease domain of MTSP10.

7. The substantially purified polypeptide of any of claims 1-6 that has
5 more than about 60%, 70%, 80% or 90% sequence identity with a polypeptide that comprises the sequence of amino acids set forth as SEQ ID No. 6, wherein the polypeptide is a protease.

8. A polypeptide of any of claims 1-7, wherein the protease domain
10 portion is encoded by a nucleic acid molecule that hybridizes under conditions of high stringency along at least 70% of its full-length to a nucleic acid molecule comprising a sequence of nucleotides set forth in SEQ ID No. 5 or at least one domain thereof or a catalytically active portion of the domain.

9. The polypeptide of claim 8, wherein the domain is the protease domain.

15 10. The polypeptide of any of claims 1-4, that comprises residues 1-230 of SEQ ID No. 6.

11. A polypeptide of any of claims 10, that is a mutein, wherein:
up to about 50% of the amino acids are replaced with another amino acid;

20 and the resulting polypeptide is a single chain or two chain polypeptide that has catalytic activity of at least 1%, 5% or 10% of the unmutated polypeptide.

12. The polypeptide of claim 11, wherein up to about 10% of the amino acids are replaced with another amino acid.

25 13. The polypeptide of claim 11, wherein the resulting polypeptide is a single chain or two chain polypeptide and has catalytic activity of at least 50% of the unmutated polypeptide.

14. The polypeptide of claim 11, wherein:
a cysteine in the protease domain is replaced with another amino acid;
30 and the cysteine is one that is unpaired in a polypeptide that consists essentially of the protease domain.

-163-

15. The polypeptide of claim 14, wherein the replacing amino acid is a serine.
16. A nucleic acid molecule, comprising a sequence of nucleotides that encodes a polypeptide of any of claims 1-15.
- 5 17. The nucleic acid molecule of claim 16 that comprises a sequence of nucleotides selected from the group consisting of:
- (a) a sequence of nucleotides set forth in as nucleotides 1-690 in SEQ ID No. 5 or set forth in SEQ ID No. 23 or a portion thereof that encodes a proteolytically active polypeptide;
- 10 (b) a sequence of nucleotides that hybridizes under high stringency along its length or along at least about 70% of the full-length to the sequence of nucleotides in SEQ ID No. 5;
- (c) a sequence of nucleotides that encodes the polypeptide of SEQ ID No. 6;
- 15 (d) a sequence of nucleotides that is a splice variant of (a), (b) or (c);
- (e) a sequence of nucleotides that encodes the protease domain or a catalytically active portion thereof that includes a sequence of nucleotides having at least about 60%, 70%, 80%, 90% or 95% sequence identity the sequence set forth in SEQ ID No. 5; and
- 20 (f) a sequence of nucleotides comprising degenerate codons of (a), (b), (c), (d) or (e).
18. An isolated nucleic molecule that encodes a mutein polypeptide of claim 11.
19. A vector comprising the nucleic acid molecule of any of claims
- 25 16-18.
20. The vector of claim 19 that is an expression vector.
21. The vector of claim 20 that is a eukaryotic vector.
22. The vector of claim 20 or claim 21 that includes a sequence of nucleotides that directs secretion of any polypeptide encoded by a sequence of
- 30 nucleotides operatively linked thereto.
23. The vector of claim 22 that is a *Pichia* vector or an *E. coli* vector.

-164-

24. A cell, comprising the vector of any of claims 19-23.
25. The cell of claim 24 that is a prokaryotic cell.
26. The cell of claim 24 that is a eukaryotic cell.
27. The cell of claim 24 that is selected from among a bacterial cell, a
5 yeast cell, a plant cell, an insect cell and an animal cell.
28. The cell of claim 24 that is a mammalian cell.
29. A recombinant non-human animal, wherein an endogenous gene
that encodes a polypeptide of any of claims 1-15 has been deleted or inactivated
by homologous recombination or insertional mutagenesis of the animal or an
10 ancestor thereof.
30. A method for producing a polypeptide that contains a protease
domain of an MTSP10 polypeptide, comprising:
culturing the cell of any of claims 24-28 under conditions whereby the
encoded polypeptide is expressed by the cell; and
15 recovering the expressed polypeptide.
31. The method of claim 30, wherein the polypeptide is secreted into
the culture medium.
32. The method of claim 30, wherein the cell is a *Pichia* cell.
33. The method of claim 30, wherein the polypeptide is expressed in
20 the cytoplasm of the host cell.
34. An antisense nucleic acid molecule that comprises at least 14
contiguous nucleotides or modified nucleotides that are complementary to a
contiguous sequence of nucleotides encoding the protease domain of an
MTSP10 of any of claims 1-15; or
25 comprises at least 16 contiguous nucleotides or modified nucleotides that
are complementary to a contiguous sequence of nucleotides encoding the
protease domain of an MTSP10 any of claims 1-15; or
comprises at least 30 contiguous nucleotides or modified nucleotides that
are complementary to a contiguous sequence of nucleotides encoding the
30 protease domain of an MTSP10 of any of claims 1-15.

-165-

35. The antisense molecule of claim 34 that includes a contiguous sequence of nucleotides that is the complement of the sequence of nucleotides in SEQ ID No. 5 or SEQ ID No. 22.

36. A double-stranded RNA (dsRNA) molecule that comprises at least
5 about 21 contiguous nucleotides or modified nucleotides from a sequence of nucleotides encoding an MTSP10 of any of claims 1-15.

37. An antibody that specifically binds to the single chain form and/or two-chain form of a protease domain of the polypeptide of any of claims 1-15, or a fragment or derivative of the antibody containing a binding domain thereof,
10 wherein the antibody is a polyclonal antibody or a monoclonal antibody.

38. The antibody of claim 37 that inhibits the enzymatic activity of the polypeptide.

39. A conjugate, comprising:
a polypeptide of any of claims 1-15, and
15 a targeting agent linked to the polypeptide directly or via a linker.

40. The conjugate of claim 39, wherein the targeting agent permits
affinity isolation or purification of the conjugate;
attachment of the conjugate to a surface;
detection of the conjugate; or
20 targeted delivery to a selected tissue or cell.

41. A combination, comprising:
an agent or treatment that inhibits the catalytic activity of the polypeptide of any of claims 1-15; and
another treatment or agent selected from anti-tumor and anti-
25 angiogenic treatments and agents.

42. The combination of claim 41, wherein the inhibitor and the anti-tumor and/or anti-angiogenic agent are formulated in a single pharmaceutical composition or each is formulated in separate pharmaceutical compositions.

43. The combination of claim 41, wherein the inhibitor is selected from
30 antibodies and antisense oligonucleotides and double-stranded RNA (dsRNA).

44. A solid support comprising two or more polypeptides of any of claims 1-15 linked thereto either directly or via a linker.

-166-

45. The support of claim 44, wherein the polypeptides comprise an array.

46. The support of claim 44, wherein the polypeptides comprise a plurality of different protease domains.

5 47. A solid support comprising two or more nucleic acid molecules of any of claims 16-18 or oligonucleotide portions thereof linked thereto either directly or via a linker, wherein the oligonucleotides contain at least 16 nucleotides.

10 48. The support of claim 47, wherein the nucleic acid molecules comprise an array.

49. The support of claim 47 or claim 48, wherein the nucleic acid molecules comprise a plurality of molecules that encode different protease domains.

15 50. A method for identifying compounds that modulate the protease activity of a polypeptide, comprising:

contacting a polypeptide of any of claims 1-15 with a substrate that is proteolytically cleaved by the polypeptide, and, either simultaneously, before or after, adding a test compound or plurality thereof;

20 measuring the amount of substrate cleaved in the presence of the test compound; and

selecting compounds that change the amount of substrate cleaved compared to a control, whereby compounds that modulate the activity of the polypeptide are identified.

25 51. The method of claim 50, wherein the test compounds are small molecules, peptides, peptidomimetics, natural products, antibodies or fragments thereof that modulate the activity of the polypeptide.

52. The method of claim 50 or claim 51, wherein a plurality of the test substances are screened simultaneously.

30 53. The method of any of claims 50-52, wherein the polypeptide consists essentially of a polypeptide encoded by a sequence of nucleotides selected from the group consisting of a sequence of nucleotides that:

(a) a sequence of nucleotides in SEQ ID No. 5 or SEQ ID No. 22;

-167-

(b) a sequence of nucleotides that hybridizes under high stringency along its length or along at least about 70% of the full-length to the sequence of nucleotides in SEQ ID No. 5 or SEQ ID No. 23;

(c) a sequence of nucleotides that encodes the polypeptide of SEQ ID No. 6 or SEQ ID No. 23;

(d) a sequence of nucleotides that is a splice variant of (a), (b) or (c);

(e) a sequence of nucleotides that encodes the protease domain or a catalytically active portion thereof that includes a sequence of nucleotides having at least about 60%, 70%, 80%, 90% or 95% sequence identity with the sequence set forth in SEQ ID No. 5; and

(f) a sequence of nucleotides comprising degenerate codons of (a), (b), (c), (d) or (e).

54. The method of any of claims 50-52, wherein the polypeptide consists essentially of a polypeptide selected from the group consisting of:

- a polypeptide that comprises a sequence of amino acids encoded by the sequence of nucleotides in SEQ ID No. 5 as amino acids 1-230 or set forth in SEQ ID No. 23 and comprising amino acids 463-692 of SEQ ID No. 23;
- a polypeptide that comprises a sequence of amino acids encoded by a sequence of nucleotides that hybridizes under conditions of high stringency to the sequence of nucleotides in SEQ ID No. 5 or set forth in SEQ ID No. 23;
- a polypeptide that comprises the sequence of amino acids set forth in SEQ ID No. 6;
- a polypeptide that comprises a sequence of amino acids having at least about 60% sequence identity with the sequence of amino acids in SEQ ID No. 6 or the sequence of amino acids in SEQ ID No. 23; and
- a polypeptide that is encoded by a sequence of nucleotides that is a splice variant of the sequence set forth in SEQ ID No. 23.

55. The method of any of claims 50-54, wherein the change in the amount of substrate cleaved is assessed by comparing the amount of substrate cleaved in the presence of the test compound with the amount of substrate cleaved in the absence of the test compound.

-168-

56. The method of any of claims 50-55, wherein a plurality of the polypeptides are linked to a solid support, either directly or via a linker.

57. The method of claim 56, wherein the polypeptides comprise an array.

5 58. A method of identifying a compound that specifically binds to a single-chain and/or two-chain protease domain and/or to single or two-chain polypeptide and/or to a proteolytically active portion of the single or two chain form thereof of an MTSP10 polypeptide, comprising:

10 contacting an MTSP10 polypeptide of any of claims 1-15 or a proteolytically active portion thereof with a test compound or plurality thereof under conditions conducive to binding thereof; and either:

a) identifying test compounds that specifically bind to the single chain and/or two chain form of the polypeptide or to a proteolytically active portion of the single an/or two chain form thereof, or

15 b) identifying test compounds that inhibit binding of a compound known to bind a single chain and/or two chain form of the polypeptide or to a proteolytically active portion of the single and/or two chain form thereof, wherein the known compound is contacted with the polypeptide before, simultaneously with or after the test compound.

20 59. The method of claim 58, wherein the polypeptide is linked either directly or indirectly via a linker to a solid support.

60. The method of claim 58 or claim 59, wherein the test compounds are small molecules, peptides, peptidomimetics, natural products, antibodies or fragments thereof.

25 61. The method of any of claims 58-60, wherein a plurality of the test substances are screened simultaneously.

62. The method of any of claims 58-61, wherein a plurality of the polypeptides are linked to a solid support.

30 63. The method of any of claims 58-62, wherein the polypeptide consists essentially of a polypeptide encoded by:

(a) a sequence of nucleotides in SEQ ID No. 5;

-169-

(b) a sequence of nucleotides that hybridizes under high stringency along its length or along at least about 70% of the full-length to the sequence of nucleotides in SEQ ID No. 5;

(c) a sequence of nucleotides that encodes the polypeptide of SEQ ID No. 5
6 or the polypeptide containing amino acids 463-692 of SEQ ID No. 23;

(d) a sequence of nucleotides that is a splice variant of (a), (b) or (c);

(e) a sequence of nucleotides that encodes the protease domain or a catalytically active portion thereof that includes a sequence of nucleotides having at least about 60%, 70%, 80%, 90% or 95% sequence identity the sequence set forth in SEQ ID No. 5; and
10

(f) a sequence of nucleotides comprising degenerate codons of (a), (b), (c), (d) or (e).

64. A method for identifying activators of the zymogen form of an MTSP10, comprising:

15 contacting a zymogen form of an MTSP10 polypeptide of any of claims 1-15 or a potentially proteolytically active portion thereof with a substrate of the activated form of the polypeptide;

adding a test compound, wherein the test compound is added before, after or simultaneously with the addition of the substrate; and

20 detecting cleavage of the substrate, thereby identifying compounds that activate the zymogen.

65. The method of claim 64, wherein the substrate is a chromogenic substrate.

66. The method of claim 64 or claim 65, wherein the substrate is a
25 L-pyroglutamyl-L-prolyl-L-arginine-p-nitroaniline hydrochloride.

67. The method of any of claims 64-66, wherein the test compound is a small molecule, a nucleic acid or a polypeptide.

68. A method for treating or preventing a neoplastic disease, in a mammal, comprising administering to a mammal an effective amount of an
30 inhibitor of a polypeptide of any of claims 1-15.

-170-

69. The method of claim 68, wherein the inhibitor is an antibody that specifically binds to the polypeptide, or a fragment or derivative of the antibody containing a binding domain thereof, wherein the antibody is a polyclonal antibody or a monoclonal antibody.

5 70. The method of claim 68 or claim 69, wherein the polypeptide consists essentially of a polypeptide encoded by a sequence of nucleotides that:

(a) a sequence of nucleotides in SEQ ID No. 5;

(b) a sequence of nucleotides that hybridizes under high stringency along its length or along at least about 70% of the full-length to the sequence of

10 nucleotides in SEQ ID No. 5;

(c) a sequence of nucleotides that encodes the polypeptide of SEQ ID No. 6 or the polypeptide containing amino acids 463-692 of SEQ ID No. 23;

(d) a sequence of nucleotides that is a splice variant of (a), (b) or (c);

15 (e) a sequence of nucleotides that encodes the protease domain or a catalytically active portion thereof that includes a sequence of nucleotides having at least about 60%, 70%, 80%, 90% or 95% sequence identity the sequence set forth in SEQ ID Nos. 5; and

(f) a sequence of nucleotides comprising degenerate codons of (a), (b), (c), (d) or (e).

20 71. The method of claim 68 or claim 69, wherein the polypeptide comprises a polypeptide encoded by:

(a) a sequence of nucleotides in SEQ ID No. 5;

(b) a sequence of nucleotides that hybridizes under high stringency along its length to the sequence of nucleotides in SEQ ID No. 5;

25 (c) a sequence of nucleotides that encodes the polypeptide of SEQ ID No. 6 or the polypeptide containing amino acids 463-692 of SEQ ID No. 23;

(d) a sequence of nucleotides that is a splice variant of (a), (b) or (c);
and

(e) degenerate codons of (a), (b), (c) or (d).

-171-

72. A method of inhibiting tumor initiation, growth or progression or treating a malignant or pre-malignant condition, comprising administering an agent that inhibits activation cleavage of the zymogen form of an MTSP10 polypeptide of any of claims 1-15 or a potentially proteolytically active portion thereof or inhibits an activity of the activated form of MTSP10 or a proteolytically active portion thereof.

73. The method of claim 72, wherein the condition is a condition of the breast, cervix, prostate, lung, ovary or colon.

74. The method of claim 72 or claim 73, wherein the agent is an antisense oligonucleotide, double-stranded RNA (dsRNA) or an antibody.

75. The method of any of claims 72-74, further comprising administering another treatment or agent selected from anti-tumor and anti-angiogenic treatments or agents.

76. The method of any of claims 72-74, wherein the polypeptide consists essentially of a polypeptide encoded by:

(a) a sequence of nucleotides in SEQ ID No. 5;

(b) a sequence of nucleotides that hybridizes under high stringency along its length or along at least about 70% of the full-length to the sequence of nucleotides in SEQ ID No. 5;

(c) a sequence of nucleotides that encodes the polypeptide of SEQ ID No. 6;

(d) a sequence of nucleotides that is a splice variant of (a), (b) or (c);

(e) a sequence of nucleotides that encodes the protease domain or a catalytically active portion thereof that includes a sequence of nucleotides having at least about 60%, 70%, 80%, 90% or 95% sequence identity with the sequence set forth in SEQ ID Nos. 5; and

(f) a sequence of nucleotides comprising degenerate codons of (a), (b), (c), (d) or (e).

77. The method of any of claims 72-74, wherein the polypeptide comprises a polypeptide encoded by a sequence of nucleotides that:

(a) a sequence of nucleotides in SEQ ID No. 5;

-172-

(b) a sequence of nucleotides that hybridizes under high stringency along its length to the sequence of nucleotides in SEQ ID No. 5;

(c) a sequence of nucleotides that encodes the polypeptide of SEQ ID No. 6 or the polypeptide containing amino acids 463-692 of SEQ ID No. 23;

5 (d) a sequence of nucleotides that is a splice variant of (a), (b) or (c);
and

(e) degenerate codons of (a), (b), (c) or (d).

78. A method of identifying a compound that binds to the single-chain and/or two-chain form of an MTSP10 polypeptide and/or to a proteolytically
10 active portion of a single-chain and/or two-chain form of an MTSP10 polypeptide of any of claims 1-15, comprising:

contacting a test compound with both forms;

determining to which form the compound binds; and

15 if it binds to a form of polypeptide, further determining whether the
compound has at least one of the following properties:

(i) inhibits activation cleavage of the single-chain zymogen form of polypeptide;

(ii) inhibits activity of the two-chain or single-chain form; and

(iii) inhibits dimerization of the polypeptide.

20 79. The method of claim 78, wherein the biological sample is selected from the group consisting of blood, urine, saliva, tears, synovial fluid, sweat, interstitial fluid, cerebrospinal fluid, a sperm sample, ascites fluid, tumor tissue biopsy and circulating tumor cells.

80. The method of claim 78 or claim 79, wherein both forms consist
25 essentially of the protease domain.

81. A method of detecting neoplastic disease, comprising: detecting a polypeptide that comprises a polypeptide of any of claims 1-15 in a biological sample, wherein the amount, the form and/or activity detected differs from the amount the form and/or activity of polypeptide detected from a subject who
30 does not have neoplastic disease.

82. The method of claim 81, wherein the biological sample is selected from the group consisting of blood, urine, saliva, tears, synovial fluid, sweat,

-173-

interstitial fluid, sperm, cerebrospinal fluid, ascites fluid, tumor tissue biopsy and circulating tumor cells.

83. A polypeptide of any of claims 1-15 that is a two chain form of MTSP that comprises from Cys₂₉₆ up to and including Cys₅₇₃ of SEQ ID No. 23.

5 84. A method of diagnosing the presence of a pre-malignant lesion, a malignancy, or other pathologic condition in a subject, comprising:

obtaining a biological sample from the subject; and

exposing it to a detectable agent that binds to a two-chain and/or single-chain form of an MTSP10 polypeptide, wherein the pathological condition is

10 characterized by the presence or absence of the two-chain or single-chain form.

85. A method of monitoring tumor progression and/or therapeutic effectiveness, comprising detecting and/or quantifying the level, the form and/or activity of an MTSP10 polypeptide in a body tissue or fluid sample.

86. The method of claim 84 or claim 85, wherein the tumor is a tumor
15 of the breast, cervix, prostate, lung, ovary or colon.

87. The method of any of claims 84-86, wherein the body fluid is blood, urine, sweat, saliva, cerebrospinal fluid and synovial fluid.

88. An isolated substantially pure polypeptide that consists essentially of the protease domain of MTSP10.

20 89. A method for identifying compounds that modulate the protease activity of an MTSP10 polypeptide, comprising:

contacting a polypeptide of any of claims 1-15 or a proteolytically active portion thereof with a substrate that is proteolytically cleaved by the polypeptide, and, either simultaneously, before or after, adding a test compound
25 or plurality thereof;

measuring the amount of substrate cleaved in the presence of the test compound; and

selecting compounds that change the amount of substrate cleaved compared to a control, whereby compounds that modulate the activity of the
30 polypeptide are identified.

-174-

90. The method of claim 89, wherein the polypeptide comprises a polypeptide encoded by a sequence of nucleotides that:

- (a) a sequence of nucleotides in SEQ ID No. 5;
- (b) a sequence of nucleotides that hybridizes under high stringency along
5 its length or along at least about 70% of the full-length to the sequence of nucleotides in SEQ ID No. 5;
- (c) a sequence of nucleotides that encodes the polypeptide of SEQ ID No. 6 or the polypeptide containing amino acids 463-692 of SEQ ID No. 23;
- (d) a sequence of nucleotides that is a splice variant of (a), (b) or (c);
- 10 (e) a sequence of nucleotides that encodes the protease domain or a catalytically active portion thereof that includes a sequence of nucleotides having at least about 60%, 70%, 80%, 90% or 95% sequence identity the sequence set forth in SEQ ID Nos. 5; and
- (f) a sequence of nucleotides comprising degenerate codons of (a),
15 (b), (c), (d) or (e).

91. The method of claim 89, wherein the polypeptide consists essentially of a polypeptide encoded by a sequence of nucleotides that:

- (a) a sequence of nucleotides in SEQ ID No. 5;
- (b) a sequence of nucleotides that hybridizes under high stringency along
20 its length or along at least about 70% of the full-length to the sequence of nucleotides in SEQ ID No. 5;
- (c) a sequence of nucleotides that encodes the polypeptide of SEQ ID No. 6 or the polypeptide containing amino acids 463-692 of SEQ ID No. 23;
- (d) a sequence of nucleotides that is a splice variant of (a), (b) or (c);
- 25 (e) a sequence of nucleotides that encodes the protease domain or a catalytically active portion thereof that includes a sequence of nucleotides having at least about 60%, 70%, 80%, 90% or 95% sequence identity the sequence set forth in SEQ ID Nos. 5; and
- (f) a sequence of nucleotides comprising degenerate codons of (a),
30 (b), (c), (d) or (e).

-175-

92. The polypeptide of any of claims 89-91, wherein the protease domain comprises the sequence of amino acids set forth in SEQ ID No. 6.

93. A transgenic non-human animal, comprising heterologous nucleic acid encoding a polypeptide of any of claims 1-15.

5 94. A probe or primer that comprises at least 14 contiguous nucleotides or modified nucleotides that are identical to a contiguous sequence of nucleotides encoding the protease domain of an MTSP10 of any of claims 1-15; or

 comprises at least 16 contiguous nucleotides or modified nucleotides that
10 are identical to a contiguous sequence of nucleotides encoding the protease domain of an MTSP10 of any of claims 1-15; or

 comprises at least 30 contiguous nucleotides or modified nucleotides that
are identical to a contiguous sequence of nucleotides encoding the protease
domain of an MTSP10 of any of claims 1-15, wherein the antisense molecule
15 does include nucleotides 1162-1262 of SEQ ID No. 18.

95. The polypeptide of any of claims 1-15, wherein:
the polypeptide includes at least amino acids 85-87 and/or 160-165 of
SEQ ID No. 18.

96. A polypeptide of any of claims 1-15 that comprises a sequence of
20 amino acids that includes one or one or more polypeptides that contain residues
104-217, 222-335, 340-377, 381-412 or 415-453 of SEQ ID No. 23.

-1-

SEQUENCE LISTING

<110> Corvas International, Inc.
 Madison, Edward
 Yeh, Jiunn-Chern

<120> NUCLEIC ACID MOLECULES ENCODING A TRANSMEMBRANE SERINE PROTEASE 10, THE
 ENCODED POLYPEPTIDES AND METHODS BASED THEREON

<130> 24745-1616PC

<140> Not Yet Assigned
 <141> Herewith

<150> 60/291,001
 <151> 14-MAY-2002

<160> 23

<170> FastSEQ for Windows Version 4.0

<210> 1
 <211> 3147
 <212> DNA
 <213> Homo Sapien

<220>
 <221> CDS
 <222> (23)...(2589)
 <223> Nucleotide sequence encoding MTSP1

<300>
 <301> O'Brien, T.J. and Tanimoto, H.
 <308> GenBank #AR081724
 <309> 2000-08-31
 <310> 5,972,616
 <311> 1998-02-20
 <312> 1999-10-26

<400> 1
 tcaagagcgg cctcggggta cc atg ggg agc gat cgg gcc cgc aag ggc gga 52
 Met Gly Ser Asp Arg Ala Arg Lys Gly Gly
 1 5 10

ggg ggc ccg aag gac ttc ggc gcg gga ctc aag tac aac tcc cgg cac 100
 Gly Gly Pro Lys Asp Phe Gly Ala Gly Leu Lys Tyr Asn Ser Arg His
 15 20 25

gag aaa gtg aat ggc ttg gag gaa ggc gtg gag ttc ctg cca gtc aac 148
 Glu Lys Val Asn Gly Leu Glu Glu Gly Val Glu Phe Leu Pro Val Asn
 30 35 40

aac gtc aag aag gtg gaa aag cat ggc ccg ggg cgc tgg gtg gtg ctg 196
 Asn Val Lys Lys Val Glu Lys His Gly Pro Gly Arg Trp Val Val Leu
 45 50 55

gca gcc gtg ctg atc ggc ctc ctc ttg gtc ttg ctg ggg atc ggc ttc 244
 Ala Ala Val Leu Ile Gly Leu Leu Leu Val Leu Gly Ile Gly Phe
 60 65 70

-2-

ctg gtg tgg cat ttg cag tac cgg gac gtg cgt gtc cag aag gtc ttc	292
Leu Val Trp His Leu Gln Tyr Arg Asp Val Arg Val Gln Lys Val Phe	
75 80 85 90	
aat ggc tac atg agg atc aca aat gag aat ttt gtg gat gcc tac gag	340
Asn Gly Tyr Met Arg Ile Thr Asn Glu Asn Phe Val Asp Ala Tyr Glu	
95 100 105	
aac tcc aac tcc act gag ttt gta agc ctg gcc agc aag gtg aag gac	388
Asn Ser Asn Ser Thr Glu Phe Val Ser Leu Ala Ser Lys Val Lys Asp	
110 115 120	
gcg ctg aag ctg ctg tac agc gga gtc cca ttc ctg ggc ccc tac cac	436
Ala Leu Lys Leu Leu Tyr Ser Gly Val Pro Phe Leu Gly Pro Tyr His	
125 130 135	
aag gag tgg gct gtg acg gcc ttc agc gag ggc agc gtc atc gcc tac	484
Lys Glu Ser Ala Val Thr Ala Phe Ser Glu Gly Ser Val Ile Ala Tyr	
140 145 150	
tac tgg tct gag ttc agc atc ccg cag cac ctg gtg gag gag gcc gag	532
Tyr Trp Ser Glu Phe Ser Ile Pro Gln His Leu Val Glu Glu Ala Glu	
155 160 165 170	
cgc gtc atg gcc gag gag cgc gta gtc atg ctg ccc ccg cgg gcg cgc	580
Arg Val Met Ala Glu Glu Arg Val Val Met Leu Pro Pro Arg Ala Arg	
175 180 185	
tcc ctg aag tcc ttt gtg gtc acc tca gtg gtg gct ttc ccc acg gac	628
Ser Leu Lys Ser Phe Val Val Thr Ser Val Val Ala Phe Pro Thr Asp	
190 195 200	
tcc aaa aca gta cag agg acc cag gac aac agc tgc agc ttt ggc ctg	676
Ser Lys Thr Val Gln Arg Thr Gln Asp Asn Ser Cys Ser Phe Gly Leu	
205 210 215	
cac gcc cgc ggt gtg gag ctg atg cgc ttc acc acg ccc ggc ttc cct	724
His Ala Arg Gly Val Glu Leu Met Arg Phe Thr Thr Pro Gly Phe Pro	
220 225 230	
gac agc ccc tac ccc gct cat gcc cgc tgc cag tgg gcc ctg cgg ggg	772
Asp Ser Pro Tyr Pro Ala His Ala Arg Cys Gln Trp Ala Leu Arg Gly	
235 240 245 250	
gac gcc gac tca gtg ctg agc ctc acc ttc cgc agc ttt gac ctt gcg	820
Asp Ala Asp Ser Val Leu Ser Leu Thr Phe Arg Ser Phe Asp Leu Ala	
255 260 265	
tcc tgc gac gag cgc ggc agc gac ctg gtg acg gtg tac aac acc ctg	868
Ser Cys Asp Arg Gly Ser Asp Leu Val Thr Val Tyr Asn Thr Leu	
270 275 280	
agc ccc atg gag ccc cac gcc ctg gtg cag ttg tgt ggc acc tac cct	916
Ser Pro Met Glu Pro His Ala Leu Val Gln Leu Cys Gly Thr Tyr Pro	
285 290 295	
ccc tcc tac aac ctg acc ttc cac tcc tcc cag aac gtc ctg ctc atc	964
Pro Ser Tyr Asn Leu Thr Phe His Ser Ser Gln Asn Val Leu Leu Ile	

-3-

300	305	310	
aca ctg ata acc aac act gag cgg cgg cat ccc ggc ttt gag gcc acc Thr Leu Ile Thr Asn Thr Glu Arg Arg His Pro Gly Phe Glu Ala Thr 315 320 325 330			1012
ttc ttc cag ctg cct agg atg agc agc tgt gga ggc cgc tta cgt aaa Phe Phe Gln Leu Pro Arg Met Ser Ser Cys Gly Gly Arg Leu Arg Lys 335 340 345			1060
gcc cag ggg aca ttc aac agc ccc tac tac cca ggc cac tac cca ccc Ala Gln Gly Thr Phe Asn Ser Pro Tyr Tyr Pro Gly His Tyr Pro Pro 350 355 360			1108
aac att gac tgc aca tgg aac att gag gtg ccc aac aac cag cat gtg Asn Ile Asp Cys Thr Trp Asn Ile Glu Val Pro Asn Asn Gln His Val 365 370 375			1156
aag gtg agc ttc aaa ttc ttc tac ctg ctg gag ccc ggc gtg cct gcg Lys Val Ser Phe Lys Phe Phe Tyr Leu Leu Glu Pro Gly Val Pro Ala 380 385 390			1204
ggc acc tgc ccc aag gac tac gtg gag atc aat ggg gag aaa tac tgc Gly Thr Cys Pro Lys Asp Tyr Val Glu Ile Asn Gly Glu Lys Tyr Cys 395 400 405 410			1252
gga gag agg tcc cag ttc gtc gtc acc agc aac agc aac aag atc aca Gly Glu Arg Ser Gln Phe Val Val Thr Ser Asn Ser Asn Lys Ile Thr 415 420 425			1300
gtt cgc ttc cac tca gat cag tcc tac acc gac acc ggc ttc tta gct Val Arg Phe His Ser Asp Gln Ser Tyr Thr Asp Thr Gly Phe Leu Ala 430 435 440			1348
gaa tac ctc tcc tac gac tcc agt gac cca tgc ccg ggg cag ttc acg Glu Tyr Leu Ser Tyr Asp Ser Ser Asp Pro Cys Pro Gly Gln Phe Thr 445 450 455			1396
tgc cgc acg ggg cgg tgt atc cgg aag gag ctg cgc tgt gat ggc tgg Cys Arg Thr Gly Arg Cys Ile Arg Lys Glu Leu Arg Cys Asp Gly Trp 460 465 470			1444
gcc gac tgc acc gac cac agc gat gag ctc aac tgc agt tgc gac gcc Ala Asp Cys Thr Asp His Ser Asp Glu Leu Asn Cys Ser Cys Asp Ala 475 480 485 490			1492
ggc cac cag ttc acg tgc aag aac aag ttc tgc aag ccc ctc ttc tgg Gly His Gln Phe Thr Cys Lys Asn Lys Phe Cys Lys Pro Leu Phe Trp 495 500 505			1540
gtc tgc gac agt gtg aac gac tgc gga gac aac agc gac gag cag ggg Val Cys Asp Ser Val Asn Asp Cys Gly Asp Asn Ser Asp Glu Gln Gly 510 515 520			1588
tgc agt tgt ccg gcc cag acc ttc agg tgt tcc aat ggg aag tgc ctc Cys Ser Cys Pro Ala Gln Thr Phe Arg Cys Ser Asn Gly Lys Cys Leu 525 530 535			1636
tcg aaa agc cag cag tgc aat ggg aag gac gac tgt ggg gac ggg tcc			1684

-4-

Ser	Lys	Ser	Gln	Gln	Cys	Asn	Gly	Lys	Asp	Asp	Cys	Gly	Asp	Gly	Ser		
540						545					550						
gac	gag	gcc	tcc	tgc	ccc	aag	gtg	aac	gtc	gtc	act	tgt	acc	aaa	cac	1732	
Asp	Glu	Ala	Ser	Cys	Pro	Lys	Val	Asn	Val	Val	Thr	Cys	Thr	Lys	His		
555					560					565					570		
acc	tac	cgc	tgc	ctc	aat	ggg	ctc	tgc	ttg	agc	aag	ggc	aac	cct	gag	1780	
Thr	Tyr	Arg	Cys	Leu	Asn	Gly	Leu	Cys	Leu	Ser	Lys	Gly	Asn	Pro	Glu		
				575					580					585			
tgt	gac	ggg	aag	gag	gac	tgt	agc	gac	ggc	tca	gat	gag	aag	gac	tgc	1828	
Cys	Asp	Gly	Lys	Glu	Asp	Cys	Ser	Asp	Gly	Ser	Asp	Glu	Lys	Asp	Cys		
			590					595					600				
gac	tgt	ggg	ctg	cgg	tca	ttc	acg	aga	cag	gct	cgt	gtt	gtt	ggg	ggc	1876	
Asp	Cys	Gly	Leu	Arg	Ser	Phe	Thr	Arg	Gln	Ala	Arg	Val	Val	Gly	Gly		
		605					610					615					
acg	gat	gcg	gat	gag	ggc	gag	tgg	ccc	tgg	cag	gta	agc	ctg	cat	gct	1924	
Thr	Asp	Ala	Asp	Glu	Gly	Glu	Trp	Pro	Trp	Gln	Val	Ser	Leu	His	Ala		
		620				625					630						
ctg	ggc	cag	ggc	cac	atc	tgc	ggg	gct	tcc	ctc	atc	tct	ccc	aac	tgg	1972	
Leu	Gly	Gln	Gly	His	Ile	Cys	Gly	Ala	Ser	Leu	Ile	Ser	Pro	Asn	Trp		
635					640					645					650		
ctg	gtc	tct	gcc	gca	cac	tgc	tac	atc	gat	gac	aga	gga	ttc	agg	tac	2020	
Leu	Val	Ser	Ala	Ala	His	Cys	Tyr	Ile	Asp	Asp	Arg	Gly	Phe	Arg	Tyr		
				655					660					665			
tca	gac	ccc	acg	cag	tgg	acg	gcc	ttc	ctg	ggc	ttg	cac	gac	cag	agc	2068	
Ser	Asp	Pro	Thr	Gln	Trp	Thr	Ala	Phe	Leu	Gly	Leu	His	Asp	Gln	Ser		
			670					675					680				
cag	cgc	agc	gcc	cct	ggg	gtg	cag	gag	cgc	agg	ctc	aag	cgc	atc	atc	2116	
Gln	Arg	Ser	Ala	Pro	Gly	Val	Gln	Glu	Arg	Arg	Leu	Lys	Arg	Ile	Ile		
		685				690						695					
tcc	cac	ccc	ttc	ttc	aat	gac	ttc	acc	ttc	gac	tat	gac	atc	gcg	ctg	2164	
Ser	His	Pro	Phe	Phe	Asn	Asp	Phe	Thr	Phe	Asp	Tyr	Asp	Ile	Ala	Leu		
		700				705					710						
ctg	gag	ctg	gag	aaa	cgc	gca	gag	tac	agc	tcc	atg	gtg	cgg	ccc	atc	2212	
Leu	Glu	Leu	Glu	Lys	Pro	Ala	Glu	Tyr	Ser	Ser	Met	Val	Arg	Pro	Ile		
715					720					725				730			
tgc	ctg	ccg	gac	gcc	tcc	cat	gtc	ttc	cct	gcc	ggc	aag	gcc	atc	tgg	2260	
Cys	Leu	Pro	Asp	Ala	Ser	His	Val	Phe	Pro	Ala	Gly	Lys	Ala	Ile	Trp		
				735					740					745			
gtc	acg	ggc	tgg	gga	cac	acc	cag	tat	gga	ggc	act	ggc	gcg	ctg	atc	2308	
Val	Thr	Gly	Trp	Gly	His	Thr	Gln	Tyr	Gly	Gly	Thr	Gly	Ala	Leu	Ile		
			750					755					760				
ctg	caa	aag	ggg	gag	atc	cgc	gtc	atc	aac	cag	acc	acc	tgc	gag	aac	2356	
Leu	Gln	Lys	Gly	Glu	Ile	Arg	Val	Ile	Asn	Gln	Thr	Thr	Cys	Glu	Asn		
		765					770						775				

-5-

```

ctc ctg ccg cag cag atc acg ccg cgc atg atg tgc gtg ggc ttc ctc      2404
Leu Leu Pro Gln Gln Ile Thr Pro Arg Met Met Cys Val Gly Phe Leu
780                               785                               790

agc ggc ggc gtg gac tcc tgc cag ggt gat tcc ggg gga ccc ctg tcc      2452
Ser Gly Gly Val Asp Ser Cys Gln Gly Asp Ser Gly Gly Pro Leu Ser
795                               800                               805                               810

agc gtg gag gcg gat ggg cgg atc ttc cag gcc ggt gtg gtg agc tgg      2500
Ser Val Glu Ala Asp Gly Arg Ile Phe Gln Ala Gly Val Val Ser Trp
815                               820                               825

gga gac ggc tgc gct cag agg aac aag cca ggc gtg tac aca agg ctc      2548
Gly Asp Gly Cys Ala Gln Arg Asn Lys Pro Gly Val Tyr Thr Arg Leu
830                               835                               840

cct ctg ttt cgg gac tgg atc aaa gag aac act ggg gta ta ggggccgggg      2599
Pro Leu Phe Arg Asp Trp Ile Lys Glu Asn Thr Gly Val
845                               850                               855

ccacccaaat gtgtacacct gcggggccac ccctcgtcca ccccagtgtg cacgcctgca      2659
ggctggagac tggaccgctg actgcaccag cgccccaga acatacactg tgaactcaat      2719
ctccagggct ccaaactctgc ctagaaaacc tctcgcttcc tcagcctcca aagtggagct      2779
gggaggtaga aggggaggac actggtggtt ctactgaccc aactgggggc aaaggtttga      2839
agacacagcc tcccccgcca gccccaagct gggccgaggc gcgtttgtgt atatctgcct      2899
cccctgtctg taaggagcag cgggaacgga gcttcggagc ctcctcagtg aaggtggtgg      2959
ggctgccgga tctgggctgt ggggcccttg ggccacgctc ttgaggaagc ccaggctcgg      3019
aggaccctgg aaaacagacg ggtctgagac tgaattgtt ttaccagctc ccagggtgga      3079
cttcagtgtg tgtatttgtg taaatgggta aaacaattta tttcttttta aaaaaaaaaa      3139
aaaaaaaaa                                     3147

<210> 2
<211> 855
<212> PRT
<213> Homo Sapien

<400> 2
Met Gly Ser Asp Arg Ala Arg Lys Gly Gly Gly Gly Pro Lys Asp Phe
1 5 10 15
Gly Ala Gly Leu Lys Tyr Asn Ser Arg His Glu Lys Val Asn Gly Leu
20 25 30
Glu Glu Gly Val Glu Phe Leu Pro Val Asn Asn Val Lys Lys Val Glu
35 40 45
Lys His Gly Pro Gly Arg Trp Val Val Leu Ala Ala Val Leu Ile Gly
50 55 60
Leu Leu Leu Val Leu Leu Gly Ile Gly Phe Leu Val Trp His Leu Gln
65 70 75 80
Tyr Arg Asp Val Arg Val Gln Lys Val Phe Asn Gly Tyr Met Arg Ile
85 90 95
Thr Asn Glu Asn Phe Val Asp Ala Tyr Glu Asn Ser Asn Ser Thr Glu
100 105 110
Phe Val Ser Leu Ala Ser Lys Val Lys Asp Ala Leu Lys Leu Leu Tyr
115 120 125
Ser Gly Val Pro Phe Leu Gly Pro Tyr His Lys Glu Ser Ala Val Thr
130 135 140
Ala Phe Ser Glu Gly Ser Val Ile Ala Tyr Tyr Trp Ser Glu Phe Ser
145 150 155 160
Ile Pro Gln His Leu Val Glu Glu Ala Glu Arg Val Met Ala Glu Glu
165 170 175

```

-6-

Arg Val Val Met Leu Pro Pro Arg Ala Arg Ser Leu Lys Ser Phe Val
 180 185 190
 Val Thr Ser Val Val Ala Phe Pro Thr Asp Ser Lys Thr Val Gln Arg
 195 200 205
 Thr Gln Asp Asn Ser Cys Ser Phe Gly Leu His Ala Arg Gly Val Glu
 210 215 220
 Leu Met Arg Phe Thr Thr Pro Gly Phe Pro Asp Ser Pro Tyr Pro Ala
 225 230 235 240
 His Ala Arg Cys Gln Trp Ala Leu Arg Gly Asp Ala Asp Ser Val Leu
 245 250 255
 Ser Leu Thr Phe Arg Ser Phe Asp Leu Ala Ser Cys Asp Glu Arg Gly
 260 265 270
 Ser Asp Leu Val Thr Val Tyr Asn Thr Leu Ser Pro Met Glu Pro His
 275 280 285
 Ala Leu Val Gln Leu Cys Gly Thr Tyr Pro Pro Ser Tyr Asn Leu Thr
 290 295 300
 Phe His Ser Ser Gln Asn Val Leu Leu Ile Thr Leu Ile Thr Asn Thr
 305 310 315 320
 Glu Arg Arg His Pro Gly Phe Glu Ala Thr Phe Phe Gln Leu Pro Arg
 325 330 335
 Met Ser Ser Cys Gly Gly Arg Leu Arg Lys Ala Gln Gly Thr Phe Asn
 340 345 350
 Ser Pro Tyr Tyr Pro Gly His Tyr Pro Pro Asn Ile Asp Cys Thr Trp
 355 360 365
 Asn Ile Glu Val Pro Asn Asn Gln His Val Lys Val Ser Phe Lys Phe
 370 375 380
 Phe Tyr Leu Leu Glu Pro Gly Val Pro Ala Gly Thr Cys Pro Lys Asp
 385 390 395 400
 Tyr Val Glu Ile Asn Gly Glu Lys Tyr Cys Gly Glu Arg Ser Gln Phe
 405 410 415
 Val Val Thr Ser Asn Ser Asn Lys Ile Thr Val Arg Phe His Ser Asp
 420 425 430
 Gln Ser Tyr Thr Asp Thr Gly Phe Leu Ala Glu Tyr Leu Ser Tyr Asp
 435 440 445
 Ser Ser Asp Pro Cys Pro Gly Gln Phe Thr Cys Arg Thr Gly Arg Cys
 450 455 460
 Ile Arg Lys Glu Leu Arg Cys Asp Gly Trp Ala Asp Cys Thr Asp His
 465 470 475 480
 Ser Asp Glu Leu Asn Cys Ser Cys Asp Ala Gly His Gln Phe Thr Cys
 485 490 495
 Lys Asn Lys Phe Cys Lys Pro Leu Phe Trp Val Cys Asp Ser Val Asn
 500 505 510
 Asp Cys Gly Asp Asn Ser Asp Glu Gln Gly Cys Ser Cys Pro Ala Gln
 515 520 525
 Thr Phe Arg Cys Ser Asn Gly Lys Cys Leu Ser Lys Ser Gln Gln Cys
 530 535 540
 Asn Gly Lys Asp Asp Cys Gly Asp Gly Ser Asp Glu Ala Ser Cys Pro
 545 550 555 560
 Lys Val Asn Val Val Thr Cys Thr Lys His Thr Tyr Arg Cys Leu Asn
 565 570 575
 Gly Leu Cys Leu Ser Lys Gly Asn Pro Glu Cys Asp Gly Lys Glu Asp
 580 585 590
 Cys Ser Asp Gly Ser Asp Glu Lys Asp Cys Asp Cys Gly Leu Arg Ser
 595 600 605
 Phe Thr Arg Gln Ala Arg Val Val Gly Gly Thr Asp Ala Asp Glu Gly
 610 615 620
 Glu Trp Pro Trp Gln Val Ser Leu His Ala Leu Gly Gln Gly His Ile
 625 630 635 640
 Cys Gly Ala Ser Leu Ile Ser Pro Asn Trp Leu Val Ser Ala Ala His

Cys	Tyr	Ile	Asp	645	Arg	Gly	Phe	Arg	650	Ser	Asp	Pro	Thr	655	Gln	Trp
Thr	Ala	Phe	660	Gly	Leu	His	Asp	665	Gln	Ser	Gln	Arg	Ser	Ala	Pro	Gly
Val	Gln	Glu	675	Arg	Arg	Leu	Lys	680	Ile	Ile	Ser	His	Pro	Phe	Phe	Asn
Asp	Phe	Thr	690	Phe	Asp	Tyr	Asp	695	Ile	Ala	Leu	Leu	Glu	Leu	Glu	Lys
705	Ala	Glu	Tyr	710	Ser	Met	Val	715	Pro	Ile	Cys	Leu	Pro	Asp	Ala	720
Ala	Glu	Tyr	Ser	725	Ser	Met	Val	730	Pro	Ile	Cys	Leu	Pro	Asp	Ala	735
His	Val	Phe	740	Pro	Ala	Gly	Lys	745	Trp	Val	Thr	Gly	Trp	Gly	His	750
Thr	Gln	Tyr	755	Gly	Gly	Thr	Gly	760	Leu	Ile	Leu	Gln	Lys	Gly	Glu	765
Arg	Val	Ile	770	Asn	Gln	Thr	Thr	775	Cys	Glu	Asn	Leu	Leu	Pro	Gln	780
Thr	Pro	Arg	785	Met	Met	Cys	Val	790	Gly	Phe	Leu	Ser	Gly	Gly	Val	795
Cys	Gln	Gly	800	Asp	Ser	Gly	Gly	805	Pro	Leu	Ser	Ser	Val	Glu	Ala	810
Arg	Ile	Phe	820	Gln	Ala	Gly	Val	825	Val	Ser	Trp	Gly	Asp	Gly	Cys	830
Arg	Asn	Lys	835	Pro	Gly	Val	Tyr	840	Thr	Arg	Leu	Pro	Leu	Phe	Arg	845
Ile	Lys	Glu	850	Asn	Thr	Gly	Val	855								

```
<210> 3
<211> 3147
<212> DNA
<213> Homo Sapien

<220>
<221> CDS
<222> (1865)...(2590)
<223> Nucleic acid sequence of protease domain of MTSP1
```

<400> 3						
tcaagagcgg	cctcggggta	ccatgggggag	cgatcgggcc	cgcaagggcg	gagggggccc	60
gaaggacttc	ggcgggggac	tcaagtacaa	ctcccgccac	gagaaagtga	atggcttggga	120
ggaaggcgtg	gagttctctg	cagtcaacaa	cgtcaagaag	gtggaaaaagc	atggcccggg	180
gogctcgggtg	gtgctcggcag	ccggtctgat	cggcctctct	ttggtcttgc	tggggatcgg	240
cttctcggtg	tggcatttgc	agtaaccggga	cgtgcgtgtc	cagaaggtct	tcaatggcta	300
catgagatc	acaaatgaga	attttgtgga	tgctcacgag	acactccaact	ccactgagtt	360
tgtaagcctg	gccagcaagg	tgaaggacgc	gctgaagctg	ctgtacagcg	gagtcaccatt	420
cctgggcccc	taccacaagg	agtcggctgt	gacggccttc	agcgaggggca	cgctcatcgc	480
ctactactgg	tctgagtcca	gcatacccgca	gcacctgggtg	gcaggaggccg	agcgcgtcat	540
ggcccgaggag	cgcgtagtca	tgctgcccc	gcggggcgcg	taccgtgaagt	ccttttgtggt	600
cacctcagtg	gtggtctttc	ccacggactc	caaaaacagta	cagaggacc	aggacaacag	660
ctgcagcttt	ggcctgcacc	cccgcggtgt	ggagctgatg	cgcttccaca	cgccccgctt	720
cctgcagacc	cctacaccgg	ctcactgcgc	ctgcagatgg	gcctcgcggg	gggacgcga	780
ctcagtgctg	agcctcacct	tccgcagctt	tgaccttgcg	tctcgcgacg	agcgcggcag	840
cgacctgggtg	acggtgtata	acacctgtgag	ccccatggag	ccccacgcc	tggtgcagtt	900
gtgtggcacc	taccctccct	cctacaacct	gaccttcac	tctcccaga	acgtctctgt	960
catcacactg	ataaccaaca	ctgagcggcg	gcatcccggc	tttgaggcca	ccttcttcca	1020
gctgcctagg	atgagcagct	gtggaggccg	cttacgtaaa	gcccagggga	cattacaacg	1080
ccccactac	ccaggccact	acccacccaa	cattgactgc	acatgtgaaca	ttaggtggcc	1140
caacaacac	catgtgaagg	tgagcttcaa	attctcttac	ctgctggagc	ccggcgtgcc	1200

-8-

tgcggggcacc	tgccccaagg	actacgtgga	gatcaatggg	gagaaataact	gcggagagag	1260
gtcccagttc	gtcgtcacca	gcaacagcaa	caagatcaca	gttcgcttcc	actcagatca	1320
gtccctacacc	gacaccggct	tcttagctga	atacctctcc	tacgactcca	gtgacccatg	1380
cccggggcag	ttcacgtgcc	gcacggggcg	gtgtatccgg	aaggagctgc	gctgtgatgg	1440
ctgggcccag	tgccaccgacc	acagcgatga	gctcaactgc	agttgcgacg	ccggccacca	1500
gttcacgtgc	aagaacaagt	tctgcaagcc	cctcttctgg	gtctgcgaca	gtgtgaacga	1560
ctgcggagac	aacagcgacg	agcaggggtg	cagttgtccg	gcccagacct	tcaggtgttc	1620
caatgggaag	tgccctctga	aaagccagca	gtgcaatggg	aaggacgact	gtggggacgg	1680
gtccgacgag	gcctcctgcc	ccaaggtgaa	cgtcgtcact	tgtaccaaac	acacctaccg	1740
ctgcctcaat	gggctctgct	tgagcaaggg	caaccctgag	tgtgacggga	aggaggactg	1800
tagcgacggc	tcagatgaga	aggactgcga	ctgtgggctg	cggtcattca	cgagacaggg	1860
tcgt gtt gtt ggg ggc acg gat gcg gat gag ggc gag tgg ccc tgg cag						1909
Val Val Gly Gly Thr Asp Ala Asp Glu Gly Glu Trp Pro Trp Gln						
1 5 10 15						
gta agc ctg cat gct ctg ggc cag ggc cac atc tgc ggt gct tcc ctc						1957
Val Ser Leu His Ala Leu Gly Gln Gly His Ile Cys Gly Ala Ser Leu						
20 25 30						
atc tct ccc aac tgg ctg gtc tct gcc gca cac tgc tac atc gat gac						2005
Ile Ser Pro Asn Trp Leu Val Ser Ala Ala His Cys Tyr Ile Asp Asp						
35 40 45						
aga gga ttc agg tac tca gac ccc acg cag tgg acg gcc ttc ctg ggc						2053
Arg Gly Phe Arg Tyr Ser Asp Pro Thr Gln Trp Thr Ala Phe Leu Gly						
50 55 60						
ttg cac gac cag agc cag cgc agc gcc cct ggg gtg cag gag cgc agg						2101
Leu His Asp Gln Ser Gln Arg Ser Ala Pro Gly Val Gln Glu Arg Arg						
65 70 75						
ctc aag cgc atc atc tcc cac ccc ttc ttc aat gac ttc acc ttc gac						2149
Leu Lys Arg Ile Ile Ser His Pro Phe Phe Asn Asp Phe Thr Phe Asp						
80 85 90 95						
tat gac atc gcg ctg ctg gag ctg gag aaa ccg gca gag tac agc tcc						2197
Tyr Asp Ile Ala Leu Leu Glu Leu Glu Lys Pro Ala Glu Tyr Ser Ser						
100 105 110						
atg gtg cgg ccc atc tgc ctg ccg gac gcc tcc cat gtc ttc cct gcc						2245
Met Val Arg Pro Ile Cys Leu Pro Asp Ala Ser His Val Phe Pro Ala						
115 120 125						
ggc aag gcc atc tgg gtc acg ggc tgg gga cac acc cag tat gga ggc						2293
Gly Lys Ala Ile Trp Val Thr Gly Trp Gly His Thr Gln Tyr Gly Gly						
130 135 140						
act ggc gcg ctg atc ctg caa aag ggt gag atc cgc gtc atc aac cag						2341
Thr Gly Ala Leu Ile Leu Gln Lys Gly Glu Ile Arg Val Ile Asn Gln						
145 150 155						
acc acc tgc gag aac ctc ctg ccg cag cag atc acg ccg cgc atg atg						2389
Thr Thr Cys Glu Asn Leu Leu Pro Gln Gln Ile Thr Pro Arg Met Met						
160 165 170 175						
tgc gtg ggc ttc ctc agc ggc ggc gtg gac tcc tgc cag ggt gat tcc						2437
Cys Val Gly Phe Leu Ser Gly Gly Val Asp Ser Cys Gln Gly Asp Ser						
180 185 190						

-9-

```

ggg gga ccc ctg tcc agc gtg gag gcg gat ggg cgg atc ttc cag gcc      2485
Gly Gly Pro Leu Ser Ser Val Glu Ala Asp Gly Arg Ile Phe Gln Ala
                195                        200                        205

ggt gtg gtg agc tgg gga gac ggc tgc gct cag agg aac aag cca ggc      2533
Gly Val Val Ser Trp Gly Asp Gly Cys Ala Gln Arg Asn Lys Pro Gly
                210                        215                        220

gtg tac aca agg ctc cct ctg ttt cgg gac tgg atc aaa gag aac act      2581
Val Tyr Thr Arg Leu Pro Leu Phe Arg Asp Trp Ile Lys Glu Asn Thr
                225                        230                        235

ggg gta tag gggccggggc caccctaatg tgtacacctg cggggccacc      2630
Gly Val *
240

catcgtccac cccagtgtgc acgcctgcag gctggagact ggaccgctga ctgcaccagc      2690
gccccagaaa catacactgt gaactcaatc tccagggtc caaatctgcc tagaaaacct      2750
ctcgttctct cagcctccaa agtggagctg ggaggtagaa ggggaggaca ctggtggttc      2810

tactgaccca actggggggca aagggttgaa gacacagcct ccccgccag ccccaagctg      2870
ggccgaggcg cgtttgtgta tatctgcctc cctgtctgt aaggagcagc gggaacggag      2930
cttcggagcc tcctcagtga aggtggtggg gctgccgat ctgggctgtg gggcccttgg      2990
gccacgctct tgaggaagcc caggctcgga ggaccctgga aaacagacgg gtctgagact      3050
gaaattgttt taccagctcc cagggtggac ttcagtgtgt gtatttgtgt aaatgggtaa      3110
aacaatttat ttctttttaa aaaaaaaaaa aaaaaaaa      3147

<210> 4
<211> 241
<212> PRT
<213> Homo Sapien

<400> 4
Val Val Gly Gly Thr Asp Ala Asp Glu Gly Glu Trp Pro Trp Gln Val
 1          5          10          15
Ser Leu His Ala Leu Gly Gln Gly His Ile Cys Gly Ala Ser Leu Ile
 20          25          30
Ser Pro Asn Trp Leu Val Ser Ala Ala His Cys Tyr Ile Asp Asp Arg
 35          40          45
Gly Phe Arg Tyr Ser Asp Pro Thr Gln Trp Thr Ala Phe Leu Gly Leu
 50          55          60
His Asp Gln Ser Gln Arg Ser Ala Pro Gly Val Gln Glu Arg Arg Leu
 65          70          75          80
Lys Arg Ile Ile Ser His Pro Phe Phe Asn Asp Phe Thr Phe Asp Tyr
 85          90          95
Asp Ile Ala Leu Leu Glu Leu Glu Lys Pro Ala Glu Tyr Ser Ser Met
100          105          110
Val Arg Pro Ile Cys Leu Pro Asp Ala Ser His Val Phe Pro Ala Gly
115          120          125
Lys Ala Ile Trp Val Thr Gly Trp Gly His Thr Gln Tyr Gly Gly Thr
130          135          140
Gly Ala Leu Ile Leu Gln Lys Gly Glu Ile Arg Val Ile Asn Gln Thr
145          150          155          160
Thr Cys Glu Asn Leu Leu Pro Gln Gln Ile Thr Pro Arg Met Met Cys
165          170          175
Val Gly Phe Leu Ser Gly Gly Val Asp Ser Cys Gln Gly Asp Ser Gly
180          185          190
Gly Pro Leu Ser Ser Val Glu Ala Asp Gly Arg Ile Phe Gln Ala Gly
195          200          205

```

-10-

Val Val Ser Trp Gly Asp Gly Cys Ala Gln Arg Asn Lys Pro Gly Val
 210 215 220
 Tyr Thr Arg Leu Pro Leu Phe Arg Asp Trp Ile Lys Glu Asn Thr Gly
 225 230 235 240
 Val

<210> 5
 <211> 717
 <212> DNA
 <213> Homo sapien

<220>
 <221> CDS
 <222> (1)...(717)
 <223> Nucleotide sequence encoding MTSP10, including
 MTSP10 protease domain

<400> 5
 atc atc gga ggc aca gac acc ctg gag ggg ggt tgg ccg tgg cag gtc 48
 Ile Ile Gly Gly Thr Asp Thr Leu Glu Gly Gly Trp Pro Trp Gln Val
 1 5 10 15
 agc ctc cac ttt gtt gga tct gcc tac tgt ggt gcc tca gtc atc tcc 96
 Ser Leu His Phe Val Gly Ser Ala Tyr Cys Gly Ala Ser Val Ile Ser
 20 25 30
 agg gag tgg ctt ctt tct gca gcc cac tgt ttt cat gga aac agg ctg 144
 Arg Glu Trp Leu Leu Ser Ala Ala His Cys Phe His Gly Asn Arg Leu
 35 40 45
 tca gat ccc aca cca tgg act gca cac ctc ggg atg tat gtt cag ggg 192
 Ser Asp Pro Thr Pro Trp Thr Ala His Leu Gly Met Tyr Val Gln Gly
 50 55 60
 aat gcc aag ttt gtc tcc ccg gtg aga aga att gtg gtc cac gag tac 240
 Asn Ala Lys Phe Val Ser Pro Val Arg Arg Ile Val Val His Glu Tyr
 65 70 75 80
 tat aac agt cag act ttt gat tat gat att gct ttg cta cag ctc agt 288
 Tyr Asn Ser Gln Thr Phe Asp Tyr Asp Ile Ala Leu Leu Gln Leu Ser
 85 90 95
 att gcc tgg cct gag acc ctg aaa cag ctc att cag cca ata tgc att 336
 Ile Ala Trp Pro Glu Thr Leu Lys Gln Leu Ile Gln Pro Ile Cys Ile
 100 105 110
 cct ccc act ggt cag aga gtt cgc agt ggg gag aag tgc tgg gta act 384
 Pro Pro Thr Gly Gln Arg Val Arg Ser Gly Glu Lys Cys Trp Val Thr
 115 120 125
 ggc tgg ggg cga aga cac gaa gca gat aat aaa ggc tcc ctc gtt ctg 432
 Gly Trp Gly Arg Arg His Glu Ala Asp Asn Lys Gly Ser Leu Val Leu
 130 135 140
 cag caa gcg gag gta gag ctc att gat caa acg ctc tgt gtt tcc acc 480
 Gln Gln Ala Glu Val Glu Leu Ile Asp Gln Thr Leu Cys Val Ser Thr
 145 150 155 160
 tac ggg atc atc act tct cgg atg ctc tgt gca ggc ata atg tca ggc 528

-11-

```

Tyr Gly Ile Ile Thr Ser Arg Met Leu Cys Ala Gly Ile Met Ser Gly
      165                      170                      175

aag aga gat gcc tgc aaa gga gat tgc ggt gga cct tta tct tgt cga      576
Lys Arg Asp Ala Cys Lys Gly Asp Ser Gly Gly Pro Leu Ser Cys Arg
      180                      185                      190

aga aaa agt gat gga aaa tgg att ttg act ggc att gtt agc tgg gga      624
Arg Lys Ser Asp Gly Lys Trp Ile Leu Thr Gly Ile Val Ser Trp Gly
      195                      200                      205

cat gga agt gga cga cca aac ttt cct ggt gtt tac aca agg gtg tca      672
His Gly Ser Gly Arg Pro Asn Phe Pro Gly Val Tyr Thr Arg Val Ser
      210                      215                      220

aac ttt gtt ccc tgg att cat aaa tat gtc cct tct ctt ttg taa      717
Asn Phe Val Pro Trp Ile His Lys Tyr Val Pro Ser Leu Leu *
      225                      230                      235

```

<210> 6
 <211> 238
 <212> PRT
 <213> Homo sapien

```

<400> 6
Ile Ile Gly Gly Thr Asp Thr Leu Glu Gly Gly Trp Pro Trp Gln Val
  1      5      10      15
Ser Leu His Phe Val Gly Ser Ala Tyr Cys Gly Ala Ser Val Ile Ser
      20      25      30
Arg Glu Trp Leu Leu Ser Ala Ala His Cys Phe His Gly Asn Arg Leu
      35      40      45
Ser Asp Pro Thr Pro Trp Thr Ala His Leu Gly Met Tyr Val Gln Gly
      50      55      60
Asn Ala Lys Phe Val Ser Pro Val Arg Arg Ile Val Val His Glu Tyr
      65      70      75      80
Tyr Asn Ser Gln Thr Phe Asp Tyr Asp Ile Ala Leu Leu Gln Leu Ser
      85      90      95
Ile Ala Trp Pro Glu Thr Leu Lys Gln Leu Ile Gln Pro Ile Cys Ile
      100      105      110
Pro Pro Thr Gly Gln Arg Val Arg Ser Gly Glu Lys Cys Trp Val Thr
      115      120      125
Gly Trp Gly Arg Arg His Glu Ala Asp Asn Lys Gly Ser Leu Val Leu
      130      135      140
Gln Gln Ala Glu Val Glu Leu Ile Asp Gln Thr Leu Cys Val Ser Thr
      145      150      155      160
Tyr Gly Ile Ile Thr Ser Arg Met Leu Cys Ala Gly Ile Met Ser Gly
      165      170      175
Lys Arg Asp Ala Cys Lys Gly Asp Ser Gly Gly Pro Leu Ser Cys Arg
      180      185      190
Arg Lys Ser Asp Gly Lys Trp Ile Leu Thr Gly Ile Val Ser Trp Gly
      195      200      205
His Gly Ser Gly Arg Pro Asn Phe Pro Gly Val Tyr Thr Arg Val Ser
      210      215      220
Asn Phe Val Pro Trp Ile His Lys Tyr Val Pro Ser Leu Leu
      225      230      235

```

<210> 7
 <211> 26

-12-

<212> DNA
<213> Artificial Sequence

<220>
<223> 5' end primer
<400> 7
cgcatcatcg gaggcacaga caccct 26

<210> 8
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> 3' end primer
<400> 8
ccaggaaca aagtttgaca cccttggtg 28

<210> 9
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR 5' end primer
<400> 9
cgcatcatcg gaggcacaga caccct 25

<210> 10
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR 3' end primer

<400> 10
ctcccactgg tcagagagtt cgcagtg 27

<210> 11
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR 5' end primer

<400> 11
cgcatcatcg gaggcacaga caccct 26

<210> 12
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR 3' end primer

<400> 12

-13-

ctccccactg cgaactctct gaccagtg

28

<210> 13

<211> 9276

<212> DNA

<213> *Pichia pastoris*

<400> 13

agatctaaca	tccaaagacg	aaagggtgaa	tgaaaccttt	ttgccatccg	acatccacag	60
gtccattctc	acacataagt	gccaaacgca	acaggagggg	atacactagc	agcagaccgt	120
tgcaaacgca	ggacctccac	tcctcttctc	ctcaacaccc	acttttgcca	tcgaaaaacc	180
agcccagtta	ttgggcttga	ttggagctcg	ctcattccaa	ttccttctat	taggctacta	240
acaccatgac	tttattagcc	tgtctatect	ggccccctcg	gcgaggttca	tgtttgttta	300
tttcggaatg	caacaagctc	cgcattacac	ccgaacatca	ctccagatga	gggctttctg	360
agtgtggggg	caaatagttt	catgttcccc	aaatggccca	aaactgacag	tttaaaccgt	420
gtcttgggaac	ctaatatgac	aaaagcggtg	tctcatocaa	gatgaactaa	gtttgggttcg	480
ttgaaatgct	aacggccagt	tggtaaaaaa	gaaacttcca	aaagtcgcca	taccgtttgt	540
cttgtttggg	attgattgac	gaatgctcaa	aaataatctc	attaatgctt	agcgcagtct	600
ctctatcgct	tctgaacccc	ggtgcacctg	tgccgaaacg	caaatgggga	aacacccgct	660
ttttggatga	ttatgcattg	tctccacatt	gtatgcttcc	aagattctgg	tggaataact	720
gctgatagcc	taacgttcat	gatcaaaatt	taactgttct	aacccctact	tgacagcaat	780
atataaacag	aaggaagctg	ccctgtctta	aacctttttt	tttatcatca	ttattagctt	840
actttcataa	ttgcgactgg	ttccaattga	caagcttttg	attttaacga	ctttaacga	900
caacttgaga	agatcaaaaa	acaactaatt	attcgaagga	tccaaacgat	gagatttcct	960
tcaatttttta	ctgcagtttt	attcgcagca	tcctccgcat	tagctgctcc	agtcaacact	1020
acaacagaag	atgaaacggc	acaaattccg	gctgaagctg	tcacgggtta	ctcagattta	1080
gaaggggatt	togatgttgc	tgttttgcca	ttttccaaca	gcacaaataa	cggtttattg	1140
tttataaata	ctactattgc	cagcattgct	gctaagaag	aaggggtatc	tctcgagaaa	1200
agagaggctg	aagcttacgt	agaattccct	agggcgcccg	cgaattaatt	cgcttagac	1260
atgactgttc	ctcagttcaa	gttgggcact	taogagaaga	ccggtcttgc	tagattctaa	1320
tcaagaggat	gtcagaatgc	catttgcctg	agagatgcag	gcttcatctt	tgatactttt	1380
ttattttgta	cctatatagt	ataggatttt	ttttgtcatt	ttgtttcttc	tcgtacgagc	1440
ttgtcctga	tcagcctatc	tcgcagctga	tgaatatctt	gtggtagggg	tttgggaaaa	1500
tcattcgagt	ttgatgtttt	tcttggtatt	tcctactcct	cttcagagta	cagaagatta	1560
agtgagaagt	tcgtttgtgc	aagcttateg	ataagcttta	atgcggtagt	ttatcacagt	1620
taaatttgcta	acgcagtcag	gcaccgtgta	tgaaatctaa	caatgcgctc	atcgtcatcc	1680
tcggcacccg	caccctggat	gctgtaggca	taggcttggt	tatgcgggta	ctgccggggc	1740
tcttgcggga	tatcgtccat	tcogacagca	tcgccagtc	ctatggcggtg	ctgctagcgc	1800
tatatgcgtt	gatgcaattt	ctatgcgcac	ccgttctctg	agcactgtcc	gaccgctttg	1860
gccgcgcgcc	agtcctgttc	gcttcgctac	ttggagccac	tatcgactac	gcgatcatgg	1920
cgaccacacc	cgtcctgtgg	atctatcgaa	tctaaatgta	agttaaaatc	tctaaataat	1980
taaataagtc	ccagttttct	catacgaacc	ttaacagcat	tgcggtgagc	atctagacct	2040
tcaacagcag	ccagatccat	cactgcttgg	ccaatatggt	tcagtccctc	aggagttaacg	2100
tcttgtgaag	tgatgaactt	ctggaagggt	gcagtgttaa	ctccgctgta	ttgacgggca	2160
tatccgtacg	ttggcaaagt	gtgggttggt	ccggaggagt	aatctccaca	actctctgga	2220
gagtaggcac	caacaacac	agatccagcg	tggtgtactt	gatcaacata	agaagaagca	2280
ttctcgattt	gcaggatcaa	gtgttcagga	gcgtactgat	tggaactttc	caaagcctgc	2340
tcgtagggtg	caaccgatag	gggtgtagag	tgtgcaatac	acttgcgtac	aatttcaacc	2400
cttggcaact	gcacagcttg	gttgtgaaca	gcattctcaa	ttctggcaag	ctccttgtct	2460
gtcatatcga	cagccaacag	aatcacctgg	gaatcaatac	catgttcagc	ttgagacaga	2520
aggctctgag	caacgaaatc	tggtacagcg	tatttatcag	caataactag	aacttcagaa	2580
ggcccagcag	gcattgtcaat	actacacagg	gctgatgtgt	cattttgaac	catcatcttg	2640
gcagcagtaa	cgaactgggt	tcctggacca	aatattttgt	cacacttagg	aacagtttct	2700
gtcccgtaag	catagcagc	tactgcctgg	gcgcctcctg	ctagcacgat	acacttagca	2760
ccaaccttgt	gggcaacgta	gatgacttct	ggggtaaggg	taccatcctt	cttaggtgga	2820
gatgcaaaaa	caatttcttt	gcaaccagca	actttggcag	gaacacccag	catcagggaa	2880
gtggaaggca	gaattgcggg	tccaccagga	atatagaggc	caactttctc	aataggtctt	2940
gcaaaacgag	agcagactac	accagggcaa	gtctcaactt	gcaacgtctc	cgttagttag	3000
gcttcatgga	atttctctgac	gttatctata	gagagatcaa	tggtctctct	aacgttatct	3060

-14-

ggcaattgca	taagttcctc	tgggaaagga	gcttctaaca	caggtgtctt	caaagcgact	3120
ccatcaaact	tggcagtttag	ttctaaaagg	gctttgtcac	cattttgacg	aacattgtcg	3180
acaattgggt	tgactaattc	cataatctgt	tccgttttct	ggataggacg	acgaagggca	3240
tcttcaattt	cttgtgagga	ggccttagaa	acgtcaattt	tgacacaattc	aatacgacct	3300
tcagaaggga	cttctttagg	tttggattct	tcttttaggt	gttccttggt	gtatcctggc	3360
ttggcatctc	ctttccttct	agtgcctttt	agggacttca	tatccaggtt	tctctccacc	3420
tctgtccaacg	tcacaccgta	cttggcacat	ctaactaatg	caaaataaaa	taagtcagca	3480
cattcccagg	ctatatcttc	cttggattta	gcttctgcaa	gttcattcagc	ttcctcccta	3540
attttagcgt	tcaacaaaac	ttcgtcgtca	aataaccggt	tggtataaga	accttctgga	3600
gcattgtctc	tacgatccca	caaggtggct	tccatggctc	taagaccctt	tgattggcca	3660
aaacaggaag	tgcgttccaa	gtgacagaaa	ccaacacctg	tttgttcaac	cacaaatttc	3720
aagcagtcct	catcacatc	caattcgata	cccagcaact	tttgagttgc	tccagatgta	3780
gcacctttat	accacaaaac	gtgacgacga	gattggtaga	ctccagtttg	tgctcttata	3840
gctcctggaa	tagacttttt	ggacgagtag	accaggccca	acgagtaatt	agaagagtag	3900
gccaccaaaag	tagtgaatag	accatcgggg	cggtcagtag	tcaaagacgc	caacaaaatt	3960
tacttgacag	ggaacttttt	gacatcttca	gaaagtctgt	attcagtagt	caattgccga	4020
gcatacaata	tggggattat	accagaagca	acagtggaaag	tcacatctac	caactttgcy	4080
gtctcagaaa	aagcataaac	agtctacta	ccgccattag	tgaaactttt	caaatcgccc	4140
agtgaggaga	aaaaaggcac	agcgatacta	gcattagcgg	gcaaggatgc	aactttatca	4200
accaggggtcc	tatagataac	cctagcgcct	gggatcatcc	tttggaacaac	tctttctgcc	4260
aaatctaggt	ccaaaatcac	ttcattgata	ccattattgt	acaacttgag	caagttgtcg	4320
atcagctcct	caaatgggtc	ctctgtaacg	gatgactcaa	cttgacacatt	aacttgaagc	4380
tcagtcgatt	gagtgaaact	gatcaggttg	tgagctgggt	cagcagcata	gggaaacacg	4440
gcttttccta	ccaaactcaa	ggaattatca	aactctgcaa	cacttgctgta	tgagggtagc	4500
aagggaaatg	tcatacttga	agtcggacag	tgagtgtagt	cttgagaaat	tctgaagccg	4560
tatttttatt	atcagttagt	cagtcacacg	gagatcctct	acgccggacg	catcggtggc	4620
gacctgcagg	gggggggggg	gcgctgaggt	ctgcctcgtg	aagaaggtgt	tgctgactca	4680
taccaggcct	gaatcgcccc	atcatccagc	cagaaagtga	gggagccacg	gttgatgaga	4740
gctttgttgt	aggtggacca	gttggtgatt	ttgaactttt	gctttgccac	ggaacgggtc	4800
gcgtttgtcg	gaagatgcgt	gatctgatcc	ttcaactcag	caaaagtctg	atttattcaa	4860
caaagccgcc	gtcccgtcaa	gtcagcgtaa	tgctctgcca	gtgttacaac	caattaacca	4920
attctgatta	gaaaaactca	tcgagcatca	aatgaaactg	caatttatcc	atatcaggat	4980
tatcaatacc	atatttttga	aaaagccggt	tctgtaatga	aggagaaaac	tcaccgagcg	5040
agttccatag	gatggcaaga	tcctgggtatc	ggtctgcgat	tccgactcgt	ccaacatcaa	5100
tacaacctat	taatttcccc	tcgtcaaaaa	taagggttatc	aagtgaagaaa	tcaccatgag	5160
tgacgactga	atccggtagg	aatggcaaaa	gcttatgcat	ttctttccag	acttgttcaa	5220
caggccagcc	attacgctcg	tcatacaaat	cactcgcatc	aaccaaacccg	ttattcattc	5280
gtgattgcgc	ctgagcgaga	cgaaatacgc	gatcgctggt	aaaaggacaa	ttacaaacag	5340
gaatcgaatg	caaccggcgc	aggaacactg	ccagcgcac	aacaatattt	tcacctgaat	5400
caggatattc	ttctaatacc	tggaatgctg	ttttccgggg	gatcgcagtg	gtgagtaacc	5460
atgcatcatc	aggagtacgg	ataaaatgct	tgatggtcgg	aagaggcata	aattccgtca	5520
gccagtttag	tctgaccatc	tcattctgtaa	catcattggc	aacgctacct	ttgccatggt	5580
tcagaaacaa	ctctggcgca	tcgggcttcc	catacaatcg	atagattgtc	gcacctgatt	5640
gcccgacatt	atcgcgagcc	catttatacc	catataaatc	agcatccatg	ttggaattta	5700
atcgcggcct	cgagcaagac	gtttcccggt	gaatatggct	cataacaccc	cttgatttac	5760
tgtttatgta	agcagacagt	tttatgttcc	atgatgatat	atttttatct	tgtgcaatgt	5820
aacatcagag	attttgagac	acaacgtggc	tttccccccc	ccccctgcag	gtcggcatca	5880
ccggcgccac	aggtgcggtt	gctggcgcc	atctcgccga	catcaccgat	ggggaagatc	5940
gggctcgcca	cttcgggctc	atgagcgctt	gtttcgccgt	gggtatggtg	gcaggccccg	6000
tggccggggg	actgttgggc	gccatctcct	tgcatgcacc	attccttgcg	gcggcggtgc	6060
tcaacggcct	caacctacta	ctgggctgct	tcctaagca	ggagtgcgat	aagggagagc	6120
gtcagatgta	tatgattgga	agtaggggaa	tggtgatacc	cgcattcttc	agtgcttga	6180
gggtctctat	cagattatgc	ccaactaaag	caaccggagg	aggagatttc	atggtaaatt	6240
tctctgactt	ttggtcatca	gtagactcga	actgtgagac	tatctcggtt	atgacagcag	6300
aaatgtcctt	cttgagagaca	gtaaatgaag	tcccaccaat	aaagaaatcc	ttgttatcag	6360
gaacaaactt	cttggttccga	actttttcgg	tgccctgaac	tataaaatgt	agagtggata	6420
tgctcggtag	gaatggagcg	ggcaaatgct	taccttctgg	accttcaaga	ggtatgtagg	6480
gtttgtagat	actgatgcc	acttcagtga	caacgttgct	atttcgttca	aaccattccg	6540
aatccagaga	aatcaaaagt	gtttgtctac	tattgatcca	agccagtgcg	gtcttgaaac	6600

-15-

tgacaatagt	gtgctcgtgt	tttgagggtca	tctttgtatg	aataaatcta	gtctttgatc	6660
taaaataatct	tgacgagcca	aggcgataaa	tacccaaatac	taaaactctt	ttaaaacggt	6720
aaaaggacaa	gtatgtctgc	ctgtattaaa	ccccaaatca	gctcgtagtc	tgatcctcat	6780
caactttgagg	ggcactatct	tgtttttagag	aaattttgcgg	agatgcgata	tcgagaaaaa	6840
ggtacgctga	ttttaaacgt	gaaatttatc	tcaagatctc	tgccctcgcgc	gtttcgggtga	6900
tgacgggtgaa	aacctctgac	acatgcagct	cccgagagacg	gtcacagctt	gtctgtaagc	6960
ggatgccggg	agcagacaag	ccggtcaggg	cgctcagcg	ggtgttggcg	ggtgtcgggg	7020
cgcagccatg	acccagtcac	gtagcgatag	cggagtgtat	actggcttaa	ctatgcccga	7080
tcagagcaga	ttgtactgag	agtgcacat	atgcgggtgtg	aaataccgca	cagatgcgta	7140
aggagaaaaat	accgcatcag	gcgctcttcc	gcttcctcgc	tactgactc	gctgcgctcg	7200
gtcgttcggc	tgccgcgagc	ggtatcagct	cactcaaagg	cggtataacg	gttatccaca	7260
gaatcagggg	ataacgcagg	aaagaacatg	tgagcaaaag	gccagcaaaa	ggccaggaac	7320
cgtaaaaagg	ccgcgttgct	ggcgttttcc	catagggtcc	gccccctga	cgagcatcac	7380
aaaaatcgcg	gctcaagtca	gaggtggcga	aaccgcagacg	gactataaag	ataccaggcg	7440
tttccccctg	gaagctccct	cgtgcgctct	cctgttccga	ccctgcgcgt	taccggatag	7500
ctgtccgcct	ttctcccttc	gggaagcgtg	gcgctttctc	aatgctcagc	ctgtagggtat	7560
ctcagttcgg	tgtaggtcgt	tcgctccaag	ctgggctgtg	tgacgaacc	ccccgttcag	7620
cccgaccgct	gcgccttatc	cggtaactat	cgtcttgagt	ccaaccgggt	aagacacgac	7680
ttatcgccac	tgccagcagc	cactggtaac	aggattagca	gagcgaggta	tgtaggcggg	7740
gctacagagt	tcttgaagtg	gtggcctaac	tacggctaca	ctagaaggac	agtatttggt	7800
atctgcgctc	tgctgaagcc	agttaccttc	ggaaaaagag	ttggtagctc	ttgatccggc	7860
aaacaaacca	ccgctggtag	cgggtggttt	ttgtttgca	agcagcagat	tacgcgcaga	7920
aaaaaaggat	ctcaagaaga	tcctttgatc	ttttctacgg	ggtctgacgc	tcagtggaaac	7980
gaaaactcac	gttaagggat	tttggtcatg	agattatcaa	aaaggatott	cacctagatc	8040
cttttaaat	aaaaatgaag	ttttaaatca	atctaaagta	tatatgagta	aacttggtct	8100
gacagttacc	aatgcttaat	cagtgaggca	cctatctcag	cgatctgtct	atttcgttca	8160
tcctatagttg	cctgactccc	cgctcgtgtag	ataactacga	tacgggaggg	cttaccatct	8220
ggccccagtg	ctgcaatgat	accgcgagac	ccacgctcac	cggctccaga	tttatcagca	8280
ataaaccagc	cagccggaag	ggccgagcgc	agaagtggtc	ctgcaacttt	atccgcctcc	8340
atccagctcta	ttaattgttg	ccgggaagct	agagtaagta	gttcgccagt	taatagtgtg	8400
cgcaacgttg	ttgccattgc	tgccagcatc	gtgggtgcac	gctcgtcggt	tggtatggct	8460
tcattcagct	ccggttccca	acgatcaagg	cgagttacat	gatcccccat	gttgtgcaaa	8520
aaagcgggta	gctccttcgg	tcctccgac	gttgtcagaa	gtaagttggc	cgcagtggtta	8580
tcactcatgg	ttatggcagc	actgcataat	tctcttactg	tcattgccatc	cgtaagatgc	8640
ttttctgtga	ctggtgagta	ctcaaccaag	tcattctgag	aatagtgtat	gcggcgaccg	8700
agttgtctct	gcccggcgctc	aacacgggat	aataccgcgc	cacatagcag	aactttaaaa	8760
gtgctcatca	ttggaaaacg	ttcttcgggg	cgaaaactct	caaggatctt	accgctgttg	8820
agatccagtt	cgatgtaacc	cactcgtgca	cccaactgat	cttcagcatc	ttttactttc	8880
accagcgttt	ctgggtgagc	aaaaacagga	aggcaaaatg	ccgcaaaaaa	gggaataagg	8940
gcgacacgga	aatgttgaat	actcatactc	ttcctttttc	aatattattg	aagcatttat	9000
cagggttatt	gtctcatgag	cggatacata	tttgaatgta	tttagaaaaa	taaacaataa	9060
gggggttcgc	gcacatttcc	ccgaaaagtg	ccacctgacg	tctaagaaac	cattattatc	9120
atgacattaa	cctataaaaa	taggcgtatc	acgaggccct	ttcgtcttca	agaattaatt	9180
ctcatgtttg	acagcttatc	atcgataagc	tgactcatgt	tggtattgtg	aaatagacgc	9240
agatcgggaa	cactgaaaaa	taacagttat	tattcgc			9276

<210> 14

<211> 11

<212> PRT

<213> Pichia protease cleavage site

<400> 14

Lys Arg Ile Ala Ser Gly Val Ile Ala Pro Lys

1

5

10

<210> 15

<211> 40

<212> DNA

<213> Artificial Sequence

-16-

```

<220>
<223> 5' end primer
<400> 15
agttaactcg agaaaagatc atcggaggca cagacaccctg          40

<210> 16
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> 3' end primer
<400> 16
accagtggga ggaatgctta ttggctgaat gag                  33

<210> 17
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> 5' end primer
<400> 17
agttaagaat tccaaaagag aaggacata tttatg                36

<210> 18
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> 3' end primer
<400> 18
ctcattcagc caataagcat tctcccact ggt                    33

<210> 19
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> encodes streptagII sequence with a one glycine linker
<400> 19
ggagggtggt ctcattccaca atttgaaaag taa                  33

<210> 20
<211> 10
<212> PRT
<213> streptagII sequence with a one glycine linker
<400> 20
Gly Gly Trp, Ser His Pro Gln Phe Glu Lys
  1             5             10

<210> 21
<211> 31
<212> DNA
<213> Artificial Sequence

<220>

```


-17-

<223> PCR 3' end primer

<400> 21

ttacaaaaga gaaggacat atttatgaat c

31

<210> 22

<211> 2130

<212> DNA

<213> Homo Sapien

<220>

<221> CDS

<222> (0)...(2104)

<223> MTSP 10 cDNA and domain organization

<400> 22

ata aac ctg gtt tat aca aca tct gcc ttc tcc aaa ttt tat gag cag	48
Ile Asn Leu Val Tyr Thr Thr Ser Ala Phe Ser Lys Phe Tyr Glu Gln	
1 5 10 15	

tct gtt gtt gca gat gtc agc agc aac aac aaa ggc ggc ctc ctt gtc	96
Ser Val Val Ala Asp Val Ser Ser Asn Asn Lys Gly Gly Leu Leu Val	
20 25 30	

cac ttt tgg att gtt ttt gtc atg cca cgt gcc aaa ggc cac atc ttc	144
His Phe Trp Ile Val Phe Val Met Pro Arg Ala Lys Gly His Ile Phe	
35 40 45	

tgt gaa gac tgt gtt gcc gcc atc ttg aag gac tcc atc cag aca agc	192
Cys Glu Asp Cys Val Ala Ala Ile Leu Lys Asp Ser Ile Gln Thr Ser	
50 55 60	

atc ata aac cgg acc tct gtg ggg agc ttg cag gga ctg gct gtg gac	240
Ile Ile Asn Arg Thr Ser Val Gly Ser Leu Gln Gly Leu Ala Val Asp	
65 70 75 80	

atg gac tct gtg gta cta aat gct ggg ctt cgg tca gat tac tcg tca	288
Met Asp Ser Val Val Leu Asn Ala Gly Leu Arg Ser Asp Tyr Ser Ser	
85 90 95	

acc ata gga tct gac aaa ggc tgc tct cag tac ttc tat gca gag cat	336
Thr Ile Gly Ser Asp Lys Gly Cys Ser Gln Tyr Phe Tyr Ala Glu His	
100 105 110	

ctg tct ctc cac tac ccg ctg gag att tct gca gcc tca ggg agg ctg	384
Leu Ser Leu His Tyr Pro Leu Glu Ile Ser Ala Ala Ser Gly Arg Leu	
115 120 125	

atg tgt cac ttc aag ctg gtg gcc ata gtg ggc tac ctg att cgt ctc	432
Met Cys His Phe Lys Leu Val Ala Ile Val Gly Tyr Leu Ile Arg Leu	
130 135 140	

tca atc aag tcc atc caa atc gaa gcc gac aac tgt gtc act gac tcc	480
Ser Ile Lys Ser Ile Gln Ile Glu Ala Asp Asn Cys Val Thr Asp Ser	
145 150 155 160	

ctg acc att tac gac tcc ctt ttg ccc atc cgg agc agc atc ttg tac	528
Leu Thr Ile Tyr Asp Ser Leu Leu Pro Ile Arg Ser Ser Ile Leu Tyr	
165 170 175	

-18-

aga att tgt gaa ccc aca aga aca tta atg tca ttt gtt tct aca aat Arg Ile Cys Glu Pro Thr Arg Thr Leu Met Ser Phe Val Ser Thr Asn 180 185 190	576
aat ctc atg ttg gtg aca ttt aag tct cct cat ata cgg agg ctc tca Asn Leu Met Leu Val Thr Phe Lys Ser Pro His Ile Arg Arg Leu Ser 195 200 205	624
gga atc cgg gca tat ttt gag gtc att cca gaa caa aag tgt gaa aac Gly Ile Arg Ala Tyr Phe Glu Val Ile Pro Glu Gln Lys Cys Glu Asn 210 215 220	672
aca gtg ttg gtc aaa gac atc act ggc ttt gaa ggg aaa att tca agc Thr Val Leu Val Lys Asp Ile Thr Gly Phe Glu Gly Lys Ile Ser Ser 225 230 235 240	720
cca tat tac ccg agc tac tat cct cca aaa tgc aag tgt acc tgg aaa Pro Tyr Tyr Pro Ser Tyr Tyr Pro Pro Lys Cys Lys Cys Thr Trp Lys 245 250 255	768
ttt cag act tct cta tca act ctt ggc ata gca ctg aaa ttc tat aac Phe Gln Thr Ser Leu Ser Thr Leu Gly Ile Ala Leu Lys Phe Tyr Asn 260 265 270	816
tat tca ata acc aag aag agt atg aaa ggc tgt gag cat gga tgg tgg Tyr Ser Ile Thr Lys Lys Ser Met Lys Gly Cys Glu His Gly Trp Trp 275 280 285	864
gaa att tat gag cac atg tac tgt ggc tcc tac atg gat cat cag aca Glu Ile Tyr Glu His Met Tyr Cys Gly Ser Tyr Met Asp His Gln Thr 290 295 300	912
att ttt cga gtg ccc agc cct ctg gtt cac att cag ctc cag tgc agt Ile Phe Arg Val Pro Ser Pro Leu Val His Ile Gln Leu Gln Cys Ser 305 310 315 320	960
tca agg ctt tca ggc aag cca ctt ttg gca gaa tat ggc agt tac aac Ser Arg Leu Ser Gly Lys Pro Leu Leu Ala Glu Tyr Gly Ser Tyr Asn 325 330 335	1008
atc agt caa ccc tgc cct gtg gga tct ttt aga tgc tcc tcc ggt tta Ile Ser Gln Pro Cys Pro Val Gly Ser Phe Arg Cys Ser Ser Gly Leu 340 345 350	1056
tgt gtc cct cag gcc cag cgt ggt gat gga gta aat gac tgc ttt gat Cys Val Pro Gln Ala Gln Arg Gly Asp Gly Val Asn Asp Cys Phe Asp 355 360 365	1104
gaa agt gat gaa ctg ttt tgc gtg agc cct caa cct gcc tgc aat acc Glu Ser Asp Glu Leu Phe Cys Val Ser Pro Gln Pro Ala Cys Asn Thr 370 375 380	1152
agc tcc ttc agg cag cat ggc cct ctc atc tgt gat ggc ttc agg gac Ser Ser Phe Arg Gln His Gly Pro Leu Ile Cys Asp Gly Phe Arg Asp 385 390 395 400	1200
tgt gag aat ggc cgg gat gag caa aac tgc act caa agt att cca tgc Cys Glu Asn Gly Arg Asp Glu Gln Asn Cys Thr Gln Ser Ile Pro Cys	1248

-19-

405	410	415	
aac aac aga act ttt aag tgt ggc aat gat att tgc ttt agg aaa caa Asn Asn Arg Thr Phe Lys Cys Gly Asn Asp Ile Cys Phe Arg Lys Gln 420 425 430			1296
aat gca aaa tgt gat ggg aca gtg gat tgt cca gat gga agt gat gaa Asn Ala Lys Cys Asp Gly Thr Val Asp Cys Pro Asp Gly Ser Asp Glu 435 440 445			1344
gaa ggc tgc acc tgc agc agg agt tcc tcc gcc ctt cac cgc atc atc Glu Gly Cys Thr Cys Ser Arg Ser Ser Ser Ala Leu His Arg Ile Ile 450 455 460			1392
gga ggc aca gac acc ctg gag ggg ggt tgg ccg tgg cag gtc agc ctc Gly Gly Thr Asp Thr Leu Glu Gly Gly Trp Pro Trp Gln Val Ser Leu 465 470 475 480			1440
cac ttt gtt gga tct gcc tac tgt ggt gcc tca gtc atc tcc agg gag His Phe Val Gly Ser Ala Tyr Cys Gly Ala Ser Val Ile Ser Arg Glu 485 490 495			1488
tgg ctt ctt tct gca gcc cac tgt ttt cat gga aac agg ctg tca gat Trp Leu Leu Ser Ala Ala His Cys Phe His Gly Asn Arg Leu Ser Asp 500 505 510			1536
ccc aca cca tgg act gca cac ctc ggg atg tat gtt cag ggg aat gcc Pro Thr Pro Trp Thr Ala His Leu Gly Met Tyr Val Gln Gly Asn Ala 515 520 525			1584
aag ttt gtc tcc ccg gtg aga aga att gtg gtc cac gag tac tat aac Lys Phe Val Ser Pro Val Arg Arg Ile Val Val His Glu Tyr Tyr Asn 530 535 540			1632
agt cag act ttt gat tat gat att gct ttg cta cag ctc agt att gcc Ser Gln Thr Phe Asp Tyr Asp Ile Ala Leu Leu Gln Leu Ser Ile Ala 545 550 555 560			1680
tgg cct gag acc ctg aaa cag ctc att cag cca ata tgc att cct ccc Trp Pro Glu Thr Leu Lys Gln Leu Ile Gln Pro Ile Cys Ile Pro Pro 565 570 575			1728
act ggt cag aga gtt cgc agt ggg gag aag tgc tgg gta act ggc tgg Thr Gly Gln Arg Val Arg Ser Gly Glu Lys Cys Trp Val Thr Gly Trp 580 585 590			1776
ggg cga aga cac gaa gca gat aat aaa ggc tcc ctc gtt ctg cag caa Gly Arg Arg His Glu Ala Asp Asn Lys Gly Ser Leu Val Leu Gln Gln 595 600 605			1824
gcg gag gta gag ctc att gat caa acg ctc tgt gtt tcc acc tac ggg Ala Glu Val Glu Leu Ile Asp Gln Thr Leu Cys Val Ser Thr Tyr Gly 610 615 620			1872
atc atc act tct cgg atg ctc tgt gca ggc ata atg tca ggc aag aga Ile Ile Thr Ser Arg Met Leu Cys Ala Gly Ile Met Ser Gly Lys Arg 625 630 635 640			1920
gat gcc tgc aaa gga gat tcg ggt gga cct tta tct tgt cga aga aaa			1968

-20-

Asp Ala Cys Lys Gly Asp Ser Gly Gly Pro Leu Ser Cys Arg Arg Lys
 645 650 655
 agt gat gga aaa tgg att ttg act ggc att gtt agc tgg gga cat gga 2016
 Ser Asp Gly Lys Trp Ile Leu Thr Gly Ile Val Ser Trp Gly His Gly
 660 665 670
 tgt gga cga cca aac ttt cct ggt gtt tac aca agg gtg tca aac ttt 2064
 Cys Gly Arg Pro Asn Phe Pro Gly Val Tyr Thr Arg Val Ser Asn Phe
 675 680 685
 gtt ccc tgg att cat aaa tat gtc cct tct ctt ttg taa t tgcaaaaaaa 2114
 Val Pro Trp Ile His Lys Tyr Val Pro Ser Leu Leu *
 690 695 700
 aaaaaaaaaa aaaaaa 2130
 <210> 23
 <211> 700
 <212> PRT
 <213> Homo Sapien
 <400> 23
 Ile Asn Leu Val Tyr Thr Thr Ser Ala Phe Ser Lys Phe Tyr Glu Gln
 1 5 10 15
 Ser Val Val Ala Asp Val Ser Ser Asn Asn Lys Gly Gly Leu Leu Val
 20 25 30
 His Phe Trp Ile Val Phe Val Met Pro Arg Ala Lys Gly His Ile Phe
 35 40 45
 Cys Glu Asp Cys Val Ala Ala Ile Leu Lys Asp Ser Ile Gln Thr Ser
 50 55 60
 Ile Ile Asn Arg Thr Ser Val Gly Ser Leu Gln Gly Leu Ala Val Asp
 65 70 75 80
 Met Asp Ser Val Val Leu Asn Ala Gly Leu Arg Ser Asp Tyr Ser Ser
 85 90 95
 Thr Ile Gly Ser Asp Lys Gly Cys Ser Gln Tyr Phe Tyr Ala Glu His
 100 105 110
 Leu Ser Leu His Tyr Pro Leu Glu Ile Ser Ala Ala Ser Gly Arg Leu
 115 120 125
 Met Cys His Phe Lys Leu Val Ala Ile Val Gly Tyr Leu Ile Arg Leu
 130 135 140
 Ser Ile Lys Ser Ile Gln Ile Glu Ala Asp Asn Cys Val Thr Asp Ser
 145 150 155 160
 Leu Thr Ile Tyr Asp Ser Leu Leu Pro Ile Arg Ser Ser Ile Leu Tyr
 165 170 175
 Arg Ile Cys Glu Pro Thr Arg Thr Leu Met Ser Phe Val Ser Thr Asn
 180 185 190
 Asn Leu Met Leu Val Thr Phe Lys Ser Pro His Ile Arg Arg Leu Ser
 195 200 205
 Gly Ile Arg Ala Tyr Phe Glu Val Ile Pro Glu Gln Lys Cys Glu Asn
 210 215 220
 Thr Val Leu Val Lys Asp Ile Thr Gly Phe Glu Gly Lys Ile Ser Ser
 225 230 235 240
 Pro Tyr Tyr Pro Ser Tyr Tyr Pro Pro Lys Cys Lys Cys Thr Trp Lys
 245 250 255
 Phe Gln Thr Ser Leu Ser Thr Leu Gly Ile Ala Leu Lys Phe Tyr Asn
 260 265 270
 Tyr Ser Ile Thr Lys Lys Ser Met Lys Gly Cys Glu His Gly Trp Trp
 275 280 285

-21-

Glu Ile Tyr Glu His Met Tyr Cys Gly Ser Tyr Met Asp His Gln Thr
 290 295 300
 Ile Phe Arg Val Pro Ser Pro Leu Val His Ile Gln Leu Gln Cys Ser
 305 310 315 320
 Ser Arg Leu Ser Gly Lys Pro Leu Leu Ala Glu Tyr Gly Ser Tyr Asn
 325 330 335
 Ile Ser Gln Pro Cys Pro Val Gly Ser Phe Arg Cys Ser Ser Gly Leu
 340 345 350
 Cys Val Pro Gln Ala Gln Arg Gly Asp Gly Val Asn Asp Cys Phe Asp
 355 360 365
 Glu Ser Asp Glu Leu Phe Cys Val Ser Pro Gln Pro Ala Cys Asn Thr
 370 375 380
 Ser Ser Phe Arg Gln His Gly Pro Leu Ile Cys Asp Gly Phe Arg Asp
 385 390 395 400
 Cys Glu Asn Gly Arg Asp Glu Gln Asn Cys Thr Gln Ser Ile Pro Cys
 405 410 415
 Asn Asn Arg Thr Phe Lys Cys Gly Asn Asp Ile Cys Phe Arg Lys Gln
 420 425 430
 Asn Ala Lys Cys Asp Gly Thr Val Asp Cys Pro Asp Gly Ser Asp Glu
 435 440 445
 Glu Gly Cys Thr Cys Ser Arg Ser Ser Ser Ala Leu His Arg Ile Ile
 450 455 460
 Gly Gly Thr Asp Thr Leu Glu Gly Gly Trp Pro Trp Gln Val Ser Leu
 465 470 475 480
 His Phe Val Gly Ser Ala Tyr Cys Gly Ala Ser Val Ile Ser Arg Glu
 485 490 495
 Trp Leu Leu Ser Ala Ala His Cys Phe His Gly Asn Arg Leu Ser Asp
 500 505 510
 Pro Thr Pro Trp Thr Ala His Leu Gly Met Tyr Val Gln Gly Asn Ala
 515 520 525
 Lys Phe Val Ser Pro Val Arg Arg Ile Val Val His Glu Tyr Tyr Asn
 530 535 540
 Ser Gln Thr Phe Asp Tyr Asp Ile Ala Leu Leu Gln Leu Ser Ile Ala
 545 550 555 560
 Trp Pro Glu Thr Leu Lys Gln Leu Ile Gln Pro Ile Cys Ile Pro Pro
 565 570 575
 Thr Gly Gln Arg Val Arg Ser Gly Glu Lys Cys Trp Val Thr Gly Trp
 580 585 590
 Gly Arg Arg His Glu Ala Asp Asn Lys Gly Ser Leu Val Leu Gln Gln
 595 600 605
 Ala Glu Val Glu Leu Ile Asp Gln Thr Leu Cys Val Ser Thr Tyr Gly
 610 615 620
 Ile Ile Thr Ser Arg Met Leu Cys Ala Gly Ile Met Ser Gly Lys Arg
 625 630 635 640
 Asp Ala Cys Lys Gly Asp Ser Gly Gly Pro Leu Ser Cys Arg Arg Lys
 645 650 655
 Ser Asp Gly Lys Trp Ile Leu Thr Gly Ile Val Ser Trp Gly His Gly
 660 665 670
 Cys Gly Arg Pro Asn Phe Pro Gly Val Tyr Thr Arg Val Ser Asn Phe
 675 680 685
 Val Pro Trp Ile His Lys Tyr Val Pro Ser Leu Leu
 690 695 700